

1 Isolation and characterization of antagonistic bacteria with the potential 2 for biocontrol of soil-borne wheat diseases

3
4 **Running title:** Biocontrol of soil-borne wheat diseases by XH-9

5
6 **Xiaohui Wang¹, Changdong Wang², Chao Ji², Qian Li², Jiamiao Zhang², Xin Song², Sui
7 Jun Kang¹, Zhaoyang Liu², Jianfeng Du², Xunli Liu^{2*}**

8
9 ¹ College of Life Science, Shandong Agriculture University, No. 61, Daizong Street, Taian,
10 Shandong 271018, China

11 ² College of Forestry, Shandong Agriculture University, No. 61, Daizong Street, Taian,
12 Shandong 271018, China

13
14 * Corresponding author, E-mail: xunliliu@163.com

15 Tel.: 0086 0538 8249131

16 Fax: 0086 0538 8249164

17

18

19

20

21

22

23

24

25

Abstract

Bacillus amyloliquefaciens subsp. *plantarum* XH-9 is a plant-beneficial rhizobacterium that shows good antagonistic potential against phytopathogens by releasing diffusible and volatile antibiotics, and secreting hydrolytic enzymes. Furthermore, the XH-9 strain possesses important plant growth-promoting characteristics, including nitrogen fixation (7.92 ± 1.05 mg/g), phosphate solubilization (58.67 ± 4.20 µg/L), potassium solubilization (10.07 ± 1.26 µg/mL), and the presence of siderophores (4.92 ± 0.46 µg/mL), indole-3-acetic acid (IAA) (7.76 ± 0.51 µg/mL) and 1-aminocyclopropane-1-carboxylic acid deaminase (ACC-deaminase) (4.67 ± 1.21 nmol/[mg•h]). Moreover, the XH-9 strain showed good capacities for wheat, corn, and chili root colonization, which are critical prerequisites for controlling soil-borne diseases as a bio-control agent. Real-time quantitative polymerase chain reaction experiments showed that the amount of *Fusarium oxysporum* DNA associated with the XH-9 strain after treatment significantly decreased compared with control group. Accordingly, wheat plants inoculated with the XH-9 strain showed significant increases in the plant shoot heights (14.20%), root lengths (32.25%), dry biomass levels (11.93%), and fresh biomass levels (16.28%) relative to the un-inoculated plants. The results obtained in this study suggest that the XH-9 strain has potential as plant-growth promoter and biocontrol agent when applied in local arable land to prevent damage caused by *F. oxysporum* and other phytopathogens.

Importance

Plant diseases, particularly soilborne pathogens, play a significant role in the destruction of agricultural resources. Although these diseases can be controlled to some extent with crop and fungicides, while these measures increase the cost of production, promote resistance, and

lead to environmental contamination, so they are being phased out. Plant growth-promoting rhizobacteria are an alternative to chemical pesticides that can play a key role in crop production by means of siderophore and indole-3-acetic acid production, antagonism to soilborne root pathogens, phosphate and potassium solubilization, and nitrogen fixation. These rhizobacteria can also promote a beneficial change in the microorganism community by significantly reducing its pathogenic fungi component. Their use is fully in accord with the principles of sustainability.

Key words: Antagonistic bacteria, Biological control, Plant-growth promotion, Real-time quantitative PCR, *Bacillus amyloliquefaciens plantarum*, Root colonization

Introduction

Plant diseases induced by soil-borne plant pathogens are among the most difficult to control and create challenging problems in agriculture. The *Fusarium* genus comprises several fungal species, many of which may cause crown rot, stem rot, and head blight in wheat (1). Especially, *Fusarium oxysporum* causes vascular wilt and root rot in over 100 plant species (2). In the past few decades, measures for preventing and controlling soil-borne diseases have usually relied on the use of resistant varieties and fungicides. However, developing resistant varieties can be difficult in the absence of a dominant gene (3), and chemical fungicides represent a potential risk to both the environment and human health (4). Furthermore, previous reports have shown that chemical-control measures create imbalances in the microbial community, which may reduce the activity of beneficial organisms and may also lead to the development of resistant pathogen strains (3). In this regard, biological control by using plant growth-promoting rhizobacteria (PGPR) is a potentially viable alternative strategy for disease management (5).

PGPR have been reported to improve plant growth and suppress the growth of plant pathogens (6). Several mechanisms have been suggested whereby bacteria can promote plant growth (phytohormone production, nitrogen fixation, and stimulating nutrient uptake) and reduce plant diseases (competition for space, nutrients, and ecological niches; production of hydrocyanic acid, siderophores, and antifungal compounds; or the induction of systemic resistance [ISR] (7, 8). Furthermore, the ability to colonize the rhizosphere has been considered a relevant trait for biocontrol (9).

To date, several microorganisms have been successfully used to control plant pathogens, of which *Pseudomonas* spp. and *Bacillus* spp. are important they are aggressive colonizers of the rhizosphere of various crops and show a broad spectrum of antagonistic activities against many pathogens (10). In addition, nonpathogenic bacteria, fungi, and actinomycetes belonging to the genera *Trichoderma*, *Fusarium*, *Streptomyces*, and *Serratia* have been used as potential biocontrol agents against damping-off and root rot pathogens, and other soil-borne pathogens (11-13).

Therefore, the aims of this study were to: (1) isolate and evaluate indigenous soil bacteria for their biocontrol potentials against fungal pathogens; (2) identify and characterize select antagonistic bacteria based on the 16S rDNA gene sequences; (3) reveal the mechanism for promoting plant growth and suppressing plant pathogens in vitro; (4) test the ability of bacteria to colonize the rhizosphere of different crops; and (5) determine the impact of selected strain on the *F. oxysporum* content in the wheat rhizosphere by performing real-time quantitative polymerase chain reaction (qPCR) experiments.

Results

Isolation, characterization, and identification of antagonistic bacteria

Numerous bacterial colonies with obviously variable morphological features were grown on Potato dextrose agar (PDA) medium. Several isolates of effective bacterial

antagonists belonged to the *Bacillus* genus. Subsequently, these bacterial isolates were evaluated for their plant growth-promoting traits. Among them, the XH-9 strain was selected from our collection in this work, which not only showed wide broad-spectrum activity against phytopathogens (Fig. 1), but also showed evidence of nitrogen fixation (7.92 ± 1.05 mg/g); phosphate solubilization (58.67 ± 4.20 μ g/L) and potassium solubilization (10.07 ± 1.26 μ g/mL); and siderophore (4.92 ± 0.46 μ g/mL), IAA (7.76 ± 0.51 μ g/mL), and ACC deaminase (4.67 ± 1.21 nmol/[mg•h]) production. Thus, the XH-9 strain possessed several key plant growth-promoting characteristics. Next, the XH-9 strain was identified as *Bacillus amyloliquefaciens subsp. Plantarum*, based on phylogenetic analysis of the 16S rRNA sequence and was deposited in the China General Microbiological Culture Collection Center (CGMCC) under accession number no.13151.

Antagonistic mechanisms

Both diffusible antibiotics and volatile metabolites of the XH-9 strain inhibited the growth of *F. oxysporum*, *B. sorokiniana*, and *R. solani*. The colony diameters of these pathogens in the treatment and control groups are presented in Table 1. Compared with the control group, diffusible antibiotics and volatile metabolites efficiently reduced the growth of *F. oxysporum* to 6.59% and 47.07%, respectively, and the growth of *R. solani* to 14.29% and 54.29%, respectively. Both diffusible antibiotics and volatile metabolites completely inhibited the growth of *B. sorokiniana*. Additionally, the XH-9 strain grew in solid media containing sodium carboxymethyl cellulose, yeast glucans, or skimmed milk as the sole carbon source and produced hyaline zones around the bacterial colonies (Fig. 2). These data indicated that the XH-9 strain produced several hydrolases. Quantification of the produced enzymes showed that the cellulase, glucanase, and protease levels reached 22.42 ± 2.63 U/mL, 624.19 ± 52.42 U/mL, and 13.40 ± 1.28 U/mL, respectively. Interestingly, the level of

chitinase secreted by the XH-9 strain was up to 769.21 ± 27.90 U/mL, although no semicircular hyaline zone formed around the colonies on the plate containing colloidal chitin.

Root-colonization assay

Pot experiments were performed to test the colonization capability of the XH-9 strain. We assessed bacterial population densities in the rhizosphere of different crops at 10-d intervals after bacterial inoculation. The results shown in Table 2 indicated that the XH-9 strain could colonize chili corn and whea. Almost 10^4 – 10^5 (CFUs)/g of root were detected in root rhizosphere soil samples from different crops. The result indicated that the XH-9 strain had an exceptionally high ability to stably colonize the root system of different crops, which is a critical prerequisite for the successful use of PGPR.

Biomass accumulation in a pot experiment

The performance of the XH-9 strain was evaluated after challenging it by inoculation with a suspension of 10^7 conidia mL⁻¹ from *F. oxysporum*. As shown in Table 3, the XH-9 strain promoted the growth of wheat seedlings. Compared with the control group, the XH-9 strain significantly increased wheat growth in terms of the plant shoot height (14.20%), root length (32.25%), dry biomass (11.93%), and fresh biomass (16.28%), relative to the pathogen control.

Plasmid construction and extraction

Genomic DNA was extracted from *F. oxysporum*, *F. pseudograminearum*, *F. graminearum*, *B. sorokiniana*, and *R. solani* and observed under ultraviolet (UV) light and after running it on a 2% agarose gel (Fig. 3a). Next, the JB-F and JB-R primer pair was tested in amplification reactions against all prepared fungal genome DNA samples and water

(as a negative control). As expected, the primers only amplifying the target sequence of *F. oxysporum* with a specific PCR product of 323 base pairs (bp) (Fig. 3b), which was used to construct recombinant plasmid. Subsequently, the plasmids were transformed into phage-resistant, chemically competent Trans1-T1 cells and grown in PDA medium containing 8 μ L isopropyl β -D-1-thiogalac topyranoside (IPTG) and 40 μ L X-gal at 30°C. After 24 h, a large number of white and blue colonieals appeared on the plate (Fig. 3c)everal white colonies were selected and sequenced at a commercial sequencing company (Sangon Biotech, Shanghai, China) using the M13 forward and M13 reverse primer. The specific sequence of *F. oxysporum* is shown in additional file. Transformants that were successfully identified were expanded in culture to extract plasmids (TransGen Biotech, Beijing, China), which were used in the standard curve.

Real-time PCR for the quantification of F. oxysporum

A standard curve was established by plotting the log of known concentrations of *F. oxysporum* DNA against the associated Ct values, with an R^2 value higher than 0.99 (Fig. 4b). Fig. 4a shows a melting curve of the amplification products, which indicated the specificity of the amplification product. For the rhizosphere soil samples, total soil DNA was extracted (Fig. 3d) and used as a template in qPCR assays. The Ct values observed with unknown soil samples were interpolated with the standard curve, and the calculated *F. oxysporum* DNA concentration from the soil samples was record in Table 4. Compared with the control group, the XH-9 strain significantly inhibited the growth and reproduction of *F. oxysporum* in the wheat rhizosphere at definable intervals. In the control group, the amount of *F. oxysporum* drastically decreased to $3.74 (\pm 0.26) \times 10^4$ CFUs/g of soils during the first 20 d, and thereafter, it gradually increased and reached $46.42 (\pm 1.96) \times 10^4$ CFUs/g of soil after 50 d. Following treatment with the XH-9 strain, the amount of *F. oxysporum* decreased drastically,

being gradually reduced during the first 20 d, and then gradually declining and stabilizing until the final concentration ($1-2 \times 10^4$ CFUs/g of soil) was relatively stable.

Discussion

In the absence of targeted plant-disease control approaches, soil-borne pathogens continue lead to significant crop losses worldwide in both conventional and organic field production systems (11). Development of biological control measures using antagonistic bacteria is one of the most attractive alternatives to chemical fungicides, although it has many limitations including environmental concerns, food quality issues, and tolerance to antifungal agents after long-term applications (14). A trend exists in terms of utilizing microorganisms isolated from rhizosphere soil as biological agents, which may be better adapted to colonize the rhizosphere of crop plants and show better disease control than organisms isolated from the other sources, such as seawater, compost, and silt (15, 16).

Several species of nonpathogenic rhizobacteria, mainly belonging to the *Pseudomonas*, *Bacillus*, and *Paenibacillus* genera, have been identified from soils and used as potential biocontrol agents against soil-borne pathogens. For example, *B. amyloliquefaciens* FZB45 has been found to be an effective biocontrol agent with P-mobilizing capabilities (17). Treating seeds with *Pseudomonas fluorescens* enhanced resistance to tomato damping off (18). *Paenibacillus polymyxa* E681, with its plant-growth promotion and root-colonization ability, showed promise as a biocontrol agent for cucumber and barley (19). However, *Bacillus* species have attracted more attention than other species because their long-term viability facilitates the development of commercial products (20). Previous data suggest that *Bacillus* species exert antagonistic activities against pathogens by synthesizing antimicrobial peptides, secreting lytic enzymes, competing for nutrients and space, and inducing systemic resistance (21).

At present, *B. amyloliquefaciens*, which is known promote plant growth and to produce diverse secondary metabolites (22), was studied widely and applied as a biofertilizer and biocontrol agent in agriculture. Compared with other representatives of endospore-forming *B. amyloliquefaciens*, *B. amyloliquefaciens* subsp. *plantarum* was distinguished by the ability to efficiently colonize plant rhizospheres and suppress competing phytopathogenic bacteria and fungi (20). For example, *B.amyloliquefaciens plantarum* GR53 mitigated biotic stress by regulating the levels of plant stress hormones and antioxidant activity (21). In addition, the FZB42 strain of *B. amyloliquefaciens* subsp. *plantarum* was used to enhancing the yield of tomatoes, based on positive results for producing indole compounds and siderophores, and for their biocontrol functions against plant pathogens (23, 24). The S499 strain of *B. amyloliquefaciens* subsp. *Plantarum* was applied as a plant-beneficial rhizobacterium with good antagonistic potential against phytopathogens through the release of active secondary metabolites and the induction of systemic resistance in plants by producing considerable amounts of surfactins (25).

It should be noted that the plant growth-promoting effects of these PGPR are influenced by environmental factors. Previous findings have shown that soil moisture, temperature, and the availability of an energy source (carbon) determine the activity of these organisms and soil microflora, which are important for the incidence of disease and suppression thereof (26). These data explained why only a limited number of strains have been successfully applied in agriculture and developed commercially as biofertilizers or biocontrol agents, although several *Bacillus* strains have been developed as biocontrol agents in for plant protection and improving plant growth (27). Thus, it is a substantial advantage that indigenous microorganisms have been developed as biocontrol agents and applied in local arable land.

In conclusion, we successfully selected an effective antagonistic bacterial strain (XH-9) from wheat rhizosphere and identified it as *B. amyloliquefaciens subsp. Plantarum*, based on

phylogenetic analysis of the 16S rRNA sequence. Data from pot experiments indicated that the XH-9 strain increased in the growth of wheat and reduced the number of *F. oxysporum* in the treatment group. Antagonistic mechanisms and PGP characteristics were revealed in terms of nitrogen fixation, phosphate and potassium solubilization, and production of growth hormones, ACC deaminase, diffusible and volatile antibiotics, siderophores, cellulase, glucanase, protease, and chitinase. The key point is that the XH-9 strain can colonize the root system of different plant crops, which has been considered a relevant trait for biocontrol (28). Additionally, Chowdhury et al. (24) reported that plant growth-promoting activity is linked with the ability to suppress soil-borne plant pathogens. To our knowledge, bacteria exhibiting all known plant growth-promoting features simultaneously are rare (29). Fortunately, the XH-9 strain possesses all plant growth-promoting characteristics. Although further experiments are needed to determine the effectiveness of the XH-9 strain under field conditions, we have confidence that the XH-9 strain can be developed as a biocontrol agent for improving crop productivity in traditional and organic production systems.

Experimental Procedures

Soil sampling

Soil samples from healthy wheat plants were considered good sources of PGPR and were collected randomly from a wheat field located in Taian City (Shandong Province, China). One whole wheat plant was carefully uprooted during the grain-filling stage in 2015, the aerial parts were chopped off, and the residual root portion was placed in a polythene bag, labeled and tied, and further placed in a box containing ice. The ice box was immediately transported to a laboratory and stored in a refrigerator at 4°C for further studies.

Isolation of bacterial strains

A mixed rhizosphere soil sample was collected from the roots as described previously (6). Ten grams of soil were suspended in 90 ml of sterile distilled water and mixed on a table concentrator for 30 min. The soil samples were serially diluted (up to 10^6 -fold), plated on PDA plates, and incubated at $28^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for 2–3 d. Bacterial colonies growing on the plates were isolated according to their different visual characteristics and maintained on PDA slants at 4°C for further studies (6).

Screening and identification of antagonistic bacteria

All isolates were screened for antagonistic activity against soil-borne plant pathogenic fungi by performing dual culture assays. The isolates included member of the *Fusarium pseudograminearum*, *F. oxysporum*, *Alternaria* (Nees), *Colletotrichum gloeosporioides*, *Bipolaris sorokiniana*, *Botryosphaeria ribis*, and *Rhizoctonia solani* taxa, which are common phytopathogens and often cause soil diseases in various plants (30-33). The centers of PDA plates were inoculated with pure cultures of pathogens (5-mm discs) and incubated for 24 h. next; the bacterial isolates were incubated on both sides into 9-cm diameter Petri dishes containing PDA. Three replicate plates were maintained for each isolate. All plates were incubated at 30°C in a regulated incubator for 72 h. The percent growth inhibition was calculated using the following formula: inhibition (%) = $1 - (\text{fungal growth}/\text{control growth}) \times 100\%$ (34).

Ribosomal sequencing and phylogeny of isolates

The single most effective antagonistic bacterial strain (XH-9) showing a wide broad-spectrum against phytopathogens was selected for further characterization, based on sequence analysis of 16S rDNA gene (11). The bacterium was grown in Luria-Bertani(LB)broth culture at $28^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for 24 h and was then centrifuged at $3480 \times g$ for 2

min. DNA was extracted using the EasyPure Bacteria Genomic DNA Kit (TransGen Biotech, Beijing, China) according to the manufacturer's instructions. The 16S rDNA gene sequences from bacterial genomic DNA samples were amplified using universal primers 27F and 1492R (35). PCR amplification was performed in 50- μ L reactions containing 25 μ L of 2 $\times$ Taq PCR Master Mix (TransGen Biotech, Beijing, China), 2 μ L of each primer (10 μ M), 2 μ L of template DNA (10 ng/ μ L), and 19 μ L of ddH₂O. PCR amplification was performed under the following conditions: initial denaturation at 94°C for 3 min; 35 cycles of 94°C for 1 min, 54°C for 1 min, and 72°C for 1 min; and a final extension at 72°C for 10 min. The obtained amplicons were sequenced by a commercial sequencing company (Sangon Biotech, Shanghai, China). The obtained gene sequences were analyzed and BLAST searched against with the GenBank database (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>), and a phylogenetic tree was constructed using MEGA 5.1 software (36).

Antagonistic mechanisms

Bipolaris sorokiniana, *F. oxysporum*, and *Rhizoctonia solani* were used to test the diffusible and volatile antibiotics, as described previously (37). Chitinase activity was determined using colloidal chitin as the substrate (38); Glucanase activity was determined using yeast glucans as the substrate (39); Cellulase activity was determined using sodium carboxymethyl cellulose as the substrate (40). The amount of increased reducing sugars in the reaction mixture was determined using the dinitrosalicylic acid solution (41). Protease activity was determined by measuring tyrosine released from casein in the reaction mixture (42). Siderophore production was detected by the universal chrome azurol S) assay. An orange halo around the colonies grown on CAS media was considered as positive for the production of siderophores (43). Each experiment was performed in triplicate, and the results are expressed as mean values and standard deviations.

300

301 ***Growth-promoting characteristics of the isolates***

302 All selected antagonistic isolates were characterized for differences in their traits.
 303 Qualitative determination of phosphate and potassium solubilization by strains that induced a
 304 clear zone around colonies on NBRIP growth medium and silicate solid medium, which were
 305 considered as positive results for phosphate and potassium solubilization (44). Quantitative
 306 analysis of the phosphate- and potassium-solubilization efficiencies was conducted as
 307 described previously (45, 46). Nitrogenase activity was measured by performing a previously
 308 described (47) Salkowski assay. Salkowski assays were used to analyze IAA production in liquid cultures
 309 supplemented with L-tryptophan (0.5 mg ml⁻¹) for 48 h (48). Qualitative detection of ACC
 310 deaminase produced by the isolates was performed using DF-ACC salts minimal medium
 311 (49)

312 ***Root-colonization assay***

313 To examine the root-colonizing capacity of selected strains in vitro, the original XH-9
 314 strain was replaced by a rifampicin- and spectinomycin-resistant mutant of the XH-9 strain,
 315 which was designated XH-9M. The process of introducing mutations was performed as
 316 described previously (50). The wheat seed surface was sterilized with 75% ethanol for 1 min,
 317 exposed to 1% sodium hypochlorite for 30 min, and rinsed thoroughly with sterile distilled
 318 water. Surface-sterilized wheat seeds were sown in 10-L plastic pots containing soil
 319 collected from which the XH-9 strain was isolated. The experiment involved a completely
 320 randomized block design with three replicates for each treatment, with 5 pots per replicate;
 321 each pot has 5 wheat seeds. The XH-9M strain was grown overnight in LB broth at 30°C and
 322 200 rpm, after which the bacterial suspension was centrifuged at $1073 \times g$ for 10 min and
 323 resuspended in phosphate-buffered saline (1×10^8 CFUs/mL) prior to use. Twenty milliliters
 324 of bacterial suspension was poured into pots containing wheat seedlings that were

germinated a week before (51). Bacteria were recovered from the rhizosphere by collecting wheat rhizosphere soil every 10 days after bacterial inoculation. Five-gram soil samples were placed into 250-mL conical flasks containing 50 mL of sterile water and mixed on a table concentrator for 30 min. The suspensions were serially diluted and plated on PDA agar containing rifampicin (300 $\mu\text{g mL}^{-1}$) and spectinomycin (300 $\mu\text{g mL}^{-1}$). The plates were incubated for 2–3 d at 30°C, the number of colonies was counted, and the total population was expressed as the CFU g^{-1} of dry soil. Furthermore, the colonization ability of the selected strains was also tested on corn and chili crops in the same manner.

Pot experiment design and qPCR

In this study, qPCR was used to quantitatively measure the *F. oxysporum* biomass in wheat rhizosphere soil. Wheat seeds were sterilized and sown in clay tile pots, each of which had been inoculated with a 20-mL suspension of *F. oxysporum* conidia (10^7 mL^{-1}). After one week, the roots of plantlets were inoculated with 20 mL of bacterial suspension; the roots of the control plantlets were inoculated with the same volume of sterile water. After inoculation, both the treated and control rhizosphere soil were collected every 10 d and stored at -80°C to detect the amount of *F. oxysporum* by qPCR. The pot experiments followed a completely randomized block design with three replicates for each treatment, with 5 pots per replicate. Each pot had 5 wheat seeds. After 50 d, the wheat seedlings were uprooted separately to determine the biomass indices of the plants.

Fungal and soil DNA extraction

Mycelia of fung collected from cultures grown on PDA medium were freeze-dried liquid nitrogen and ground to a fine powder using an autoclaved mortar and pestle (52). Next, DNA

was extracted from 100–200 mg of the powdered mycelia using the HP Fungal DNA Kit (TransGen Biotech, Beijing, China).

Primer selection and specificity in vitro

qPCR assays of *F. oxysporum* DNA were conducted using the following specific primer pair: JB-F (5'-CATAACCACTTGTGTCTCGGC-3') and JB-R (5'-GAACGCGAATTAACGCGAGTC-3'), which was provided by Professor Zhiquan Mao of the College of Horticulture at Shandong Agricultural University. To evaluate the specificity of the primer pairs, genomic DNA samples were extracted from *F. pseudograminearum*, *F. graminearum*, *B. sorokiniana*, and *R. solani* and used as templates for PCR amplification, with sterile water used as a negative control. Each reaction mixture contained 25 µL of 2×Easy Taq PCR Super Mix (TransGen Biotech, Beijing, China), 2.0 µL of each primer, 2.0 µL of DNA template (10 ng/µL), and 19 µL of sterile distilled water. The thermal cycling conditions were as follows: initial denaturation at 95°C for 5 min, followed by 55 cycles of 95°C for 1 min and 54°C for 1 min, and a final extension at 72°C for 10 min. The PCR products were verified by running them on 2% agarose gels and visualizing them under UV light.

Plasmid construction and extraction

Specific target fragment of *F. oxysporum* were amplified using the specific PCR primer pair, JB-F/JB-F. The PCR amplification system and thermo cycling program are described above. The amplicons were extracted from 1% agarose gels using the EasyPure Quick Gel Extraction Kit (TransGen Biotech, Beijing, China) and spectrophotometrically quantified in a BioPhotometer to determine the amplicon concentrations. Furthermore, these amplicons were used to build carriers by ligation into the pEASY-T1 cloning vector at a 7: 1 molar ratio

and then transformed into phage-resistant, chemically competent Trans1-T1 cells. Moreover, these transformants were incubated for 1 h in LB medium at 37°C 200 rpm and plated on PDA medium containing 8 µL IPTG and 40 µL X-gal. The plates were incubated for 24 h at 30°C; several white colonies were selected and sequenced in a commercial sequencing company (Sangon Biotech, Shanghai, China).

Real-time qPCR experiments

qPCR was performed with a BioRad CFX Manager real-time PCR machine using TIAN GEN SuperReal PreMix plus SYBR Green. Plasmids were serially diluted (up to 10⁻⁶-fold) and used as DNA standards, with each standard measured in triplicate independent experiments. Each amplification was carried out in a 15-µL volume containing 7.5 µL of 2× TransStart Tip Green qPCR SuperMix (TransGen Biotech, Beijing, China), 0.6 µL of each primer (10 µM), 1 µL of template DNA (10 ng/µL), and ddH₂O added to 15 µL. A two-step amplification program was performed as follows: initial denaturation at 94°C for 30 s, followed by 45 cycles of 94°C for 5 s, 60°C for 15 s, and 72°C for 10 s. Melting curve was made at 0.6°C temperature increments every 10 s within a temperature range of 60–100°C and carried out at 95°C for 10 s, 65°C for 5 s, and 95°C for 5 s. Ct values were determined as the cycle number at which the PCR product was first fluorescently detected and was inversely related to log of the initial concentration, such that lower Ct values corresponded to higher initial DNA concentrations. A standard curve was developed by plotting the logarithm of known concentrations against the Ct values. To quantify the amount of *F. oxysporum* in the wheat rhizosphere soil, total soil DNA was extracted from wheat rhizosphere soil (treatment and control groups) every 10 d after bacterial inoculation and used as a DNA template for real-time qPCR. For each sample, 500 mg of rhizosphere soil was used to extract total soil DNA with a Soil DNA Kit (O Bio-Tek, Guangzhou, China).

The PCR amplification system and thermocycling program were the same as described above, with each DNA sample being measured in triplicate independent experiments.

Acknowledgements

This work was supported by the Special Fund for Agro-scientific Research in the Public Interest (no. 201503112) and the Forestry Industry Research special funds for public welfare projects (no. 201304212). The work was also supported by the Funds of from the Shandong forestry science and technology innovation project and Shandong “Double Tops” Program.

Compliance with Ethical Standards

This article does not contain any studies with human participants or animals performed by any of the authors.

Conflict of Interest

The authors have no conflict of interest to declare.

424 References

- 425 1. **Nourozian J, Etebarian HR, Khodakaramian G.** 2006. Biological control of *Fusarium*
426 *graminearum* on wheat by antagonistic bacteria. *SJST* **28**:29-38.
- 427 2. **Li E, Ling J, Wang G, Xiao J, Yang Y, Mao Z, Wang X, Xie B.** 2015. Comparative
428 Proteomics Analyses of Two Races of *Fusarium oxysporum* f. sp. *conglutinans* that Differ in
429 Pathogenicity. *SCI REP-UK* **5**:13663.
- 430 3. **Shanmugam V, Kanoujia N.** 2011. Biological management of vascular wilt of tomato caused by
431 *Fusarium oxysporum* f.sp. *lycopersici* by plant growth-promoting rhizobacterial mixture. *Biol*
432 *Control* **57**:85-93.
- 433 4. **Dufour MC, Fontaine S, Montarry J, Corio-Costet MF.** 2011. Assessment of fungicide
434 resistance and pathogen diversity in *Erysiphe necator* using quantitative real-time PCR assays.
435 *Pest Manag Sci* **67**:60–69.
- 436 5. **Jetiyanon K, Kloepper JW.** 2002. Mixtures of plant growth-promoting rhizobacteria for
437 induction of systemic resistance against multiple plant diseases. *Biol Control* **24**:285-291.
- 438 6. **Gopalakrishnan S, Humayun P, Kiran BK, Kannan IG, Vidya MS, Deepthi K, Rupela O.**
439 2011. Evaluation of bacteria isolated from rice rhizosphere for biological control of charcoal rot
440 of sorghum caused by *Macrophomina phaseolina* (Tassi) Goid. *World J Microb Biot* **27**:1313-
441 1321.
- 442 7. **Kloepper JW, Beauchamp CJ.** 1992. A review of issues related to measuring colonization of
443 plant roots by bacteria. *Can J Microbio* **38**:1219-1232.
- 444 8. **Ajit NS, Verma R, Shanmugam V.** 2006. Extracellular chitinases of *fluorescent pseudomonads*
445 antifungal to *Fusarium oxysporum* f. sp. *dianthi* causing carnation wilt. *Curr Microbiol* **52**:310-
446 316.
- 447 9. **Kamilova F, Leveau JH, Lugtenberg B.** 2007. *Collimonas fungivorans*, an unpredicted in vitro
448 but efficient in vivo biocontrol agent for the suppression of tomato foot and root rot. *Environ*
449 *Microbiol* **9**:1597-1603.
- 450 10. **Schippers B, Bakker AW, Bakker PAHM.** 1987. Interactions of Deleterious and Beneficial
451 Rhizosphere Microorganisms and the Effect of Cropping Practices. *Annu Rev Phytopathol*
452 **25**:339-358.
- 453 11. **Khabbaz SE, Abbasi PA.** 2014. Isolation, characterization, and formulation of antagonistic
454 bacteria for the management of seedlings damping-off and root rot disease of cucumber. *Can J*
455 *Microbio* **60**:25-33.
- 456 12. **Someya N, Nakajima M, Watanabe K, Hibi T, Akutsu K.** 2003. Influence of bacteria isolated
457 from rice plants and rhizospheres on antibiotic production by the antagonistic bacterium *Serratia*
458 *marcescens* strain B2. *J Plant Pathol* **69**:342-347.
- 459 13. **Minerdi D, Bossi S, Gullino ML, Garibaldi A.** 2009. Volatile organic compounds: a potential
460 direct long-distance mechanism for antagonistic action of *Fusarium oxysporum* strain MSA 35.
461 *Environ Microbiol* **11**:844-854.
- 462 14. **Zhan-Bin, Wu, Lei-Fei, Liang, Chen, Feng, Yong-Jun, Yuan, Xing-Fang.** 2013. Isolation,
463 screening and identification of antagonistic downy mildew/endophytic bacteria from cucumber.
464 *Eur J Plant Pathol* **137**:847-857.
- 465 15. **Cook RJ.** 2003. Making Greater Use of Introduced Microorganisms for Biological Control of
466 Plant Pathogens. *Annu Rev Phytopathol* **31**:53.
- 467 16. **Kamilova F, Leveau JH, Lugtenberg B.** 2010. *Collimonas fungivorans*, an unpredicted in vitro
468 but efficient in vivo biocontrol agent for the suppression of tomato foot and root rot. *Environ*
469 *Microbiol* **9**:1597-1603.
- 470 17. **Camiloa R, Josephw K.** 2010. Plant growth promotion by *Bacillus amyloliquefaciens* FZB45
471 depends on inoculum rate and P-related soil properties. *Biol Fert Soils* **46**:835-844.
- 472 18. **Harris D, Breese WA, Rao JVDKK.** 2005. The improvement of crop yield in marginal
473 environments using ‘on-farm’ seed priming: nodulation, nitrogen fixation, and disease resistance.
474 *Aust J Agr Res* **56**:1211-1218.
- 475 19. **Choi O, Kim J, Ryu CM, Chang SP.** 2004. Colonization and Population Changes of a
476 Biocontrol Agent, *Paenibacillus polymyxa* E681, in Seeds and Roots. *Plant Pathology J* **20**:712-
477 714 Vol. 712.

- 478 20. **Qiao JQ, Wu HJ, Huo R, Gao XW, Borriss R.** 2014. Stimulation of plant growth and
479 biocontrol by *Bacillus amyloliquefaciens* subsp. *plantarum* FZB42 engineered for improved
480 action. Chemical & Biological Technologies in Agriculture **1**:12.
- 481 21. **Kang SM, Radhakrishnan R, Lee IJ.** 2015. *Bacillus amyloliquefaciens* subsp. *plantarum* GR53,
482 a potent biocontrol agent resists Rhizoctonia disease on Chinese cabbage through hormonal and
483 antioxidants regulation. World J Microb Biot **31**:1517-1527.
- 484 22. **Borriss R.** 2011. Use of Plant-Associated Bacillus Strains as Biofertilizers and Biocontrol Agents
485 in Agriculture. Bacteria in Agrobiolgy Plant Growth Responses:41-76.
- 486 23. **Szilagyi-Zecchin VJ, Mógor ÁF, Ruaro L, Röder C.** 2015. Tomato seedlings growth (*Solanum*
487 *lycopersicum*) promoted by bacteria *Bacillus amyloliquefaciens* subsp. *plantarum* FZB42 in
488 organic system. Revista De Ciencias Agrarias **38**:26.
- 489 24. **Chowdhury SP, Hartmann A, Gao X, Borriss R.** 2015. Biocontrol mechanism by root-
490 associated *Bacillus amyloliquefaciens* FZB42 - a review. Front Microbiol **6**:780.
- 491 25. **Molinatto G, Puopolo G, Sonogo P, Moretto M, Engelen K, Viti C, Ongena M, Pertot I.** 2016.
492 Complete genome sequence of *Bacillus amyloliquefaciens* subsp. *plantarum* S499, a
493 rhizobacterium that triggers plant defences and inhibits fungal phytopathogens. J. Biotechnol.
494 **238**:56-59.
- 495 26. **Gupta VVSR, Rovira AD, Roget DK.** 2011. Principles and Management of Soil Biological
496 Factors for Sustainable Rainfed Farming Systems. Rainfed Farming Systems:149-184.
- 497 27. **Solanki MK, Kumar S, Pandey AK, Srivastava S, Singh RK, Kashyap PL, Srivastava AK,**
498 **Arora DK.** 2012. Diversity and antagonistic potential of *Bacillus* spp. associated to the
499 rhizosphere of tomato for the management of *Rhizoctonia solani*. Biocontrol Sci Techn **22**:203-
500 217.
- 501 28. **Kamilova F, Leveau JH, Lugtenberg B.** 2007. Collimonas fungivorans, an unpredicted in vitro
502 but efficient in vivo biocontrol agent for the suppression of tomato foot and root rot. Environ
503 Microbiol **9**:1597–1603.
- 504 29. **Muthukumarasamy R, Revathi G, Seshadri S, Lakshminarasimhan C.** 2002. G.
505 diazotrophicus (syn. Acetobacter diazotrophicus), a promising diazotrophic endophyte in tropics.
506 Curr Sci India **83**:137-145.
- 507 30. **Wildermuth GB, Mcnamara RB.** 1991. Effect of cropping history on soil populations of
508 *Bipolaris sorokiniana* and common root rot of wheat. Aust J Agr Res **42**:779-790.
- 509 31. **Vakalounakis DJ.** 1996. Root and stem rot of cucumber caused by *Fusarium oxysporum* f.sp.
510 *radicis-cucumerinum* f.sp. nov. Plant Dis **80**:313.
- 511 32. **Obanor F, Neate S, Simpfendorfer S, Sabburg R, Wilson P, Chakraborty S.** 2013. *Fusarium*
512 *graminearum* and *Fusarium pseudograminearum* caused the 2010 head blight epidemics in
513 Australia. Plant Pathol **62**:79–91.
- 514 33. **Wang QH, Niu SG, Liu XH, Liu YS.** 2010. Identification of species bf-02 and its antifungal
515 activity against several plant pathogens. Journal of Shandong Agricultural University **41**:513-516.
- 516 34. **Hameeda B, Harini G, Rupela OP, Kumar Rao JV, Reddy G.** 2010. Biological Control of
517 Chickpea Collar Rot by Co-inoculation of Antagonistic Bacteria and Compatible *Rhizobia*. Indian
518 J. Microbiol. **50**:419.
- 519 35. **Watanabe K, Kodama Y, Harayama S.** 2001. Design and evaluation of PCR primers to amplify
520 bacterial 16S ribosomal DNA fragments used for community fingerprinting. J Microbiol Meth
521 **44**:253-262.
- 522 36. **Mendez J.** 2014. Characterization of phosphate-solubilizing bacteria isolated from the arid soils
523 of a semi-desert region of north-east Mexico. Biol Agric Hortic **30**:211-217.
- 524 37. **Asha BB, Nayaka SC, Shankar ACU, Srinivas C, Niranjana SR.** 2011. Selection of effective
525 bio -antagonistic bacteria for biological control of tomato wilt caused by *fusarium oxysporum* f.
526 sp. *lycopersici*. Bioscan An International Quarterly Journal of Life Sciences.
- 527 38. **Jankiewicz U, Brzezinska MS, Saks E.** 2012. Identification and characterization of a chitinase
528 of *Stenotrophomonas maltophilia*, a bacterium that is antagonistic towards fungal phytopathogens.
529 J Biosci Bioeng **113**:30-35.
- 530 39. **Pérez LM, Besoain X, Reyes M, Pardo G, Montealegre J.** 2002. The expression of
531 extracellular fungal cell wall hydrolytic enzymes in different *Trichoderma harzianum* isolates
532 correlates with their ability to control Pyrenochaeta lycopersici. Biol Res. **35**:401.
- 533 40. **Miller GL.** 1959. Use of Dinitrosalicylic Acid Reagent for Determination of Reducing Sugar.
534 Anal Biochem **31**:426-428.

41. **Horikoshi K.** 1973. Comparative studies on beta-1,3-glucanases of microorganisms. Yeast Mould An Plant Protoplasts Proceedings.
42. **Bian HH, HUANG, WANG.** 2009. Protease activity and chyme protein tracing by GFP in the digestive tract of flounder *Paralichthys olivaceus*. Progress in Fishery Sciences.
43. **Schwyn B, Neilands JB.** 1987. Universal chemical assay for the detection and determination of siderophores. Anal Biochem **160**:47-56.
44. **Mehta S, Nautiyal CS.** 2001. An efficient method for qualitative screening of phosphate-solubilizing bacteria. Curr Microbiol **43**:51-56.
45. **Bray RH, Kurtz LT.** 1945. Determination of Total, Organic, and Available Forms of Phosphorus in Soils. Soil Sci **59**:39-46.
46. **LüChengQun, Huang BL.** 2010. Isolation and characterization of *azotobacteria* from pine rhizosphere. Afr J Microbiol Res **4**:1299-1306.
47. **Marra LM, Oliveira SMD, Ferreira PAA, Soares BL, Carvalho RDF, Lima JMD, Moreira FMDS.** 2012. Biological nitrogen fixation and phosphate solubilization by bacteria isolated from tropical soils. Plant Soil **357**:289-307.
48. **Abiala MA, Odebode AC, Hsu SF, Blackwood CB.** 2015. Phytobeneficial Properties of Bacteria Isolated from the Rhizosphere of Maize in Southwestern Nigerian Soils. Appl Environ Microb **81**:4736–4743.
49. **Husen E, Wahyudi AT, Suwanto A, Saraswati R.** 2009. Soybean seedling root growth promotion by 1-aminocyclopropane-1-carboxylate deaminase-producing *pseudomonads*. Indonesian Journal of Agricultural Science **10**:19-25.
50. **Ji X, Lu G, Gai Y, Gao H, Lu B, Kong L, Mu Z.** 2010. Colonization of *Morus alba* L. by the plant-growth-promoting and antagonistic bacterium *Burkholderia cepacia* strain Lu10-1. BMC Microbiol,10,1(2010-09-20) **10**:1-12.
51. **Someya N, Nakajima M, Akutsu K.** 2010. Potential of an Antagonistic Bacterium *Serratia marcescens* Strain B2 for the Biological Control of Cucumber Damping-Off Disease. Biocontrol Sci **10**:101-104.
52. **Perezartes E, Garciapedrajas MD, Bejaranoalcazar J, Jimenezdiaz RM.** 2000. Differentiation of cotton-defoliating and nondefoliating pathotypes of *Verticillium dahliae* by RAPD and specific PCR analyses. Eur J Plant Pathol **106**:507-517.

577 Tables

578 Table1 Diffusible and volatile antibiotics secreted by XH-9 strain against phytopathogens

Treatment	Diameter of the colonies (cm)		
	<i>Fusarium oxysporum</i>	<i>Rhizoctonia solani</i>	<i>Bipolaris sorokiniana</i>
CK	4.1±0.00a	7.0±0.15a	3.5±0.00a
Diffusible antibiotics	3.83±0.05b	6.0±0.06b	0.50±0.00b
Volatile antibiotics	2.17±0.05c	3.20±0.10c	0.50±0.00b

579 Note: Values are mean values ± standard deviations (n=3), Means sharing a common letter within the same column
 580 are not significantly different at $P < 0.05$.

581

582 Table 2 Population density of XH-9 strain on the different crops rhizosphere (10^5 cfu.g⁻¹ dry weight of soil)

Species	Days after application (d)				
	10d	20d	30d	40d	50d
wheat	5.70±0.5a	4.08±0.32b	0.61±0.06c	0.44±0.06c	0.63±0.03c
corn	3.22±0.22ab	3.53±0.37a	2.89±0.51b	0.80±0.10c	1.02±0.14c
chili	4.63±0.37a	0.29±0.03b	0.30±0.09b	0.27±0.07b	0.32±0.04b

583 Note: Values are mean values ± standard deviations (n=15), Means sharing a common letter within the same column
 584 are not significantly different at $P < 0.05$.

585

586

587 Table 3 Effect of the XH-9 strain on growth parameters of wheat seedling

Treatment	Shoot height (cm)	Root length (cm)	Fresh weight (g)	Dry weight (g)	Root/shoot ratio
CK	16.2±0.94b	9.83±0.60b	3.01±0.15b	0.503±0.06 b	0.352±0.01b
XH-9	18.5±0.68a	13.0±1.69a	3.50±0.34a	0.563±0.63a	0.392±0.011a

588 Notes: Values are means ± SD (n = 15). Means sharing a common letter within the same column are not significantly
 589 different at $P < 0.05$. “XH-9” denotes wheat seedlings treated with a suspension of strainXH-9 cells. “CK” denotes wheat
 590 seedlings treated with an equal volume of sterile water.

591

592 Table 4 Contents of *Fusarium oxysporum* in wheat rhizosphere ($\times 10^4$ cfu.g⁻¹ dry weight of soil)

Treatment	Days after inoculation XH-9 strain (d)				
	10d	20d	30d	40d	50d
CK	74.81±4.67a	3.74±0.26a	38.46±3.15ba	39.12±2.085a	46.42±1.96a
XH-9	37.01±1.93c	1.21±0.61c	0.9±0.06c	1.08±0.13 b	1.51±0.58b

593 Notes: Values are means $\pm$ SD (n = 15). Means sharing a common letter within the same column are not significantly
594 different at $P < 0.05$. “XH-9” denotes wheat seedlings treated with a suspension of strain XH-9 cells. “CK” denotes wheat
595 seedlings treated with an equal volume of sterile water.

596

597

598

599

600

601

602

603

Figure legends

Fig. 1. In vitro antagonistic activity of *Bacillus amyloliquefaciens* subsp. *plantarum* (XH-9 strain) against selected pathogen in dual culture assays on PDA medium, 4 d after incubation.

Fig.2. Fungal cell wall-degrading enzymes produced by XH-9 strain. (a) Protease activity analyzed using skim milk agar plates. (b) Cellulase activity analyzed using carboxyl methyl cellulose agar plates. (c) Glucanase activity analyzed using Pachyman solid medium supplemented with 6% aniline blue.

Fig. 3. Real-time PCR for quantification of *F. oxysporum*. (a) Fungal genomic DNA was extracted and observed under UV light by running it on a 2% agarose gel. Marker: Trans2K TM Plus II DNA. Lanes 1–5 represent *F. oxysporum*, *F. pseudograminearum*, *F. graminearum*, *B. sorokiniana*, and *R. solani*, respectively. (b) A specific fragment of *F. oxysporum* was amplified using the JB-F/R primer pair, which generated a 323-bp DNA product. Marker: Trans2K TM Plus II DNA. Lane 1: sterile water (negative control). Lanes 2–6: *F. oxysporum*, *F.pseudograminearum*, *F. graminearum*, *B. sorokiniana* and *R. solani* respectively. (c) Traditional blue/white screening technique used to the rapidly and conveniently detects recombinant plasmids. d) Total soil DNA of after control and XH-9 strain treatment. Lanes 1- 3 represent triplicate control group samples and lanes 4–6 represent triplicate XH-9 strain-treated samples. Marker: Trans2K TM Plus II DNA.


Fig. 4. Real-time PCR quantification of *F. oxysporum* in the wheat rhizosphere following XH-9 treatment and control treatment. (a) Melting curve analysis of PCR products from *F. oxysporum*. (b) qPCR standard curve




B. sorokiniana



F. oxysporum



F. pseudograminearum



Alternaria (Nees)



C. gloeosporioides



B. ribis






Fig.4a




Fig.4c




Fig.4b




Fig.4d

