

Neseliler et al. Neural and behavioral endophenotypes of obesity

Neural and behavioral endophenotypes of obesity.

Selin Neseliler¹, Uku Vainik^{1,2}, Mahsa Dadar¹, Yvonne H.C. Yau¹, Isabel Garcia-Garcia¹,
Stephanie G. Scala¹, Yashar Zeighami¹, D. Louis Collins¹ and Alain Dagher¹

^{1.} Montreal Neurological Institute, McGill University, Montréal, Canada

^{2.} Institute of Psychology, University of Tartu, Estonia

Word count of abstract: 249

Word count of main text: 3860

Figures and Tables: 4 Figures, 1 Table

Supplemental data: 3 Figures, 3 Tables

Short/running title: Endophenotypes of obesity

Keywords: Obesity, MRI, impulsivity, weight gain, uncontrolled eating

Abstract

Background

Impulsivity is a risk factor for obesity. It has different underlying facets that can be assessed using questionnaires. Impulsivity can be further refined by the use of food-specific questionnaires, which measure a tendency to uncontrolled eating. We examined how these impulsivity measures relate to each other, to obesity, and to brain anatomy.

Methods

We assessed students in their first year of university - a risky period for weight gain- at the beginning (N = 2214) and at the end of the school year (N = 1145) using questionnaire measures of impulsivity, personality, stress reactivity and eating-specific traits. A subset of participants (N = 72) underwent T1-weighted MRI to investigate the brain correlates of impulsivity.

Results

Using factor analysis, we show that impulsivity can be stratified into three domains, which we label stress reactivity, reward sensitivity and self-control, while eating questionnaires resolve into a single latent factor - uncontrolled eating. A watershed model shows that uncontrolled eating mediates the effect of impulsivity traits on BMI. Self-control and stress reactivity scores are associated with a thinner lateral orbitofrontal cortex. In addition, stress reactivity correlates positively with amygdala and negatively with hippocampal volume. Longitudinally, lack of self-control, not uncontrolled eating, correlates with weight gain, while stress reactivity correlates with weight loss in male students.

Conclusions

The brain-impulsivity-obesity relationship is hierarchical. Structural brain differences relate to differences in impulsivity domains which affect BMI via uncontrolled eating. However,

Neseliler et al. Neural and behavioral endophenotypes of obesity

46 longitudinally, low self-control, not uncontrolled eating, is a predictor of weight gain in this
47 sample.

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

Introduction

Obesity is a neurobehavioral disorder that stems from a vulnerable brain in a disease-promoting environment (1). Its risk factors can be studied as endophenotypes - intermediate phenotypes that link biological processes to observable outcomes (2, 3). Broadly, the neurobehavioral endophenotypes associated with obesity can be categorized as domain-general and eating-specific (4). The main domain-general endophenotype identified to date is impulsivity, defined as a tendency to act without full consideration of the consequences. Impulsivity is a multifaceted trait that has been associated with pathological gambling, substance abuse, and obesity (5, 6). Meta-analysis of impulsivity questionnaires revealed that impulsivity can further be subdivided into three domains that align with personality factors (5): 1) low *conscientiousness*, reflecting self-control ; 2) *neuroticism*, a reflection of an individual's sensitivity to stress and aversive events, and 3) *extraversion*, reflecting an individual's sensitivity to rewards (7–9). The magnitude and directionality of the association between impulsivity and obesity have been inconsistent (4, 10, 11). This inconsistency could be explained by different impulsivity measures and domains used across studies (4, 5, 11), and to weak effect sizes relating general impulsivity to BMI (12). Therefore, eating specific impulsivity questionnaires may better capture the vulnerability endophenotype.

Eating-specific impulsivity constructs include emotional eating (13), disinhibited eating (14) and power of food (15). Scores on all these questionnaires consistently and strongly correlate with BMI (4, 16) and with each other (17). Different eating behavior questionnaires may be depicting a common underlying latent factor for eating specific impulsivity (16, 18). This can be labelled uncontrolled eating (UE). While both general impulsivity traits and UE have been linked to

increased BMI, the relationships between different impulsivity domains, UE and obesity remain to be tested (i.e. some impulsivity traits might be associated with eating-specific impulsive behavior which in turn will be linked to BMI (11, 12, 19)). Statistically, if true, this hypothesis implies a complete mediation of general impulsivity traits on BMI via eating-specific impulsivity. However, so far, comprehensive models have not been built for all impulsivity domains and eating questionnaires, and have not been applied to the prediction of weight gain.

There is considerable overlap between the brain regions associated with obesity and the brain regions associated with impulsivity (10, 20, 21). Scores from both general impulsivity and conscientiousness scales are inversely correlated with the thickness of the lateral prefrontal cortex (PFC) (8, 22–25) - a region that has been related to cognitive control. Neuroticism is positively correlated with amygdala volume and negatively correlated with hippocampal volume and prefrontal cortex thickness (26–29). Extraversion and sensation seeking correlate positively with medial orbitofrontal cortex (mOFC) and ventromedial PFC (vmPFC) volume (8, 30, 31), a region associated with the computation of value (32). These results support a model where the impulsivity endophenotype might be mapped to the brain as reduced cortical thickness in the prefrontal cortex and increased gray matter volume in brain structures (e.g. amygdala, OFC, mPFC) associated with reward and value computation (33). Such a brain endophenotype has also been associated with obesity (4, 10, 21). Differences in brain morphometry and function might underlie differences in impulsivity and might thereby be associated with obesity.

If impulsivity (either general or eating-specific) is a risk factor for obesity, people with these risk endophenotypes will have higher BMI and will gain more weight compared to their peers with

non-risk endophenotypes. So far, most analyses have been conducted on cross-sectional datasets making it difficult to say if impulsivity is a risk factor or a consequence of obesity (4). The freshman year of university creates an ideal time period to test these hypotheses both cross-sectionally and longitudinally. During this time, students transition into a new environment with access to similar food and exercise options, allowing underlying vulnerability to express itself. Weight gain often happens in this short period of time and affects approximately 50-60% of students (34–40). Of the risk factors for weight gain, eating specific behaviors that revolve around the concept of impulsivity have been most commonly studied, but the results have been contradictory (38, 41–44).

We used structural brain imaging measures, general and eating-specific impulsivity questionnaires, and measures of BMI and weight gain in freshman students to test the hypothesis that the relationship between brain, impulsivity and obesity is hierarchical. We utilized a hierarchical statistical framework referred to as a 'watershed model' to study the relationship between impulsivity endophenotypes. This methodological framework has previously proven useful to study the relationships between age, brain structure and intelligence (2, 45, 46). From this perspective, brain structure will influence impulsivity measures which could affect eating-specific impulsivity (namely, UE) that is associated with over-eating. We predicted that UE would mediate the relationship between general impulsivity and BMI domains, and act as a risk factor for weight gain during the freshman year of university.

Methods and Materials

Participants

Participants were first year McGill university students, at least 18 years of age, recruited via an advertisement sent to the incoming class electronic mailing list. Participants provided their consent online and data were collected using the online survey tool LimeSurvey (<https://www.limesurvey.org>) over three consecutive years. The study was approved by the Montreal Neurological Institute Research Ethics Board. Participants for the brain imaging experiment gave additional consent before participating.

Questionnaires

Participants filled out the following online questionnaires: "Big Five" personality dimensions (Openness/ Imagination, Conscientiousness, Agreeableness, Extraversion, and Neuroticism) from the International Personality Item Pool (IPIP) (47), two subscales (lack of perseverance and sensation seeking) of the UPPS Impulsive Behavior Scale (48), Cohen's Perceived Stress Scale (PSS) (49), and Rosenberg Self Esteem Scale (50) as well as eating specific questionnaires such as the disinhibition subscale of Three-Factor Eating Questionnaire (TFEQ)(14), emotional eating subscale of Dutch Eating Behavior Questionnaires (DEBQ) (13), and all subscales of Power of Food Scale (PFS) (15) (Table 1). A subset (N=1145) of our initial population completed a follow-up questionnaire in the spring for longitudinal assessment. Our questionnaire included two "catch" questions and three "catch-match" questions as a measure of the level of participants' attention in completing the survey. Participants with total catch scores above three were excluded. An additional exclusion criterion was BMI under 15 or above 45, resulting in a sample of 2213 participants (Males N= 752; Females N= 1461). Only after completion of the

questionnaire, participants were asked to visit our laboratory to have their BMI measured in both the fall and spring. BMI was measured from 333 participants in the fall, and from 209 participants in the spring semester (N =115 overlap with the fall group) using a medical scale and a stadiometer. Reported BMI was highly correlated with measured BMI in the fall ($r = 0.91$, $p < 10^{-5}$) and in the spring ($r = 0.92$, $p < 10^{-5}$). We replaced the reported BMI with measured BMI in the analyses for higher accuracy. BMI was further residualized for age, sex and a covariate to account for whether it was reported or measured. Brain imaging was conducted on a subset of participants (N=72) selected from this sample in the fall.

Magnetic Resonance Imaging parameters and preprocessing

High-resolution T1-weighted anatomical images with voxel size = 1x1x1 mm were obtained (TR = 2.3 s; TE = 2.98 ms; FOV phase=93.8°; FOV = 256mm) with a Siemens Magnetom Trio 3T MRI scanner at the Montreal Neurological Institute (MNI).

Pre-processing of T1-weighted MRIs included denoising using optimized non-local means filtering (51), correction for intensity inhomogeneity (52) and linear intensity scaling using histogram matching to the ICBM-MNI152 template. The images were linearly registered to ICBM-MNI152 template (9 parameter registration) (53). A mask of the brain was generated using BEaST, a nonlocal segmentation method applied to the linearly registered images in stereotaxic space (54).

Deformation Based Morphometry

Deformation Based Morphometry is utilized to calculate the local difference in tissue volume between subjects (55). Each MRI was non-linearly transformed to the stereotaxic ICBM-MNI152 template. The local deformation, a measure of tissue expansion or shrinkage, was then computed at each voxel. We utilized Deformation Based Morphometry as it is suggested to be the most sensitive morphometric method for studying subcortical anatomy (as explained in (56)).

Cortical Thickness

All T1-weighted MRI images were processed using the CIVET pipeline (version 2.1; <http://www.bic.mni.mcgill.ca/ServicesSoftware/CIVET>). Native T1-weighted MRI scans were first corrected using the N3 algorithm, underwent brain masking, and registration to ICBM-MNI152 template. Images were then segmented into gray matter (GM), white matter (WM), cerebrospinal fluid (CSF) and background using a neural net classifier. The white matter (inner) and gray matter (outer) cortical surfaces were extracted using the Constrained Laplacian-based Automated Segmentation with Proximities algorithm (CLASP). The resulting surfaces were resampled to a stereotaxic surface template to provide vertex-based measures of cortical thickness. All resulting images went through a stringent quality control by two inspectors in which 69 of 72 images were accepted for further analysis. Cortical thickness was then measured in MNI space using the linked distance between the two surfaces across vertices and a 20mm surface smoothing kernel was applied to the data. The Desikan–Killiany–Tourville (DKT) atlas was used to parcellate the surface into 64 cortical regions (57). Cortical thickness was averaged over all vertices (left and right) in each region of interest (ROI) for each subject. The subject mean cortical thickness was regressed out from the ROIs before further analysis.

Factor Analysis

Factor analysis was carried out in order to (i) understand the shared variance of the measured variables, which can be attributed to latent factor(s) and (ii) reduce the collinearity between the questionnaire scores. The correlation matrix of the impulsivity traits is shown in Figure S2. Questionnaire scores were z-scored for this group and entered into the factor analysis. Exploratory Factor Analysis was carried out first in order to identify the underlying factor structure of our set of observed variables. All of the statistical analyses are carried out in R version 3.3 (58). To account for missing values, we used the R “Amelia” library to impute data (59). We conducted our exploratory factor analysis on the split half of our population of 2213 participants (N=1107) to create our model. Factor selection was conducted using parallel analysis via the fa.parallel function from the Psych package in R (60). The generated pattern matrix of standardized loadings suggested the existence of a 3-factor latent structure (Table S1).

Afterwards, confirmatory factor analysis (CFA) was utilized to test the relationship between our observed variables and their underlying latent factors in both split-halves of the sample (61). All structural equation models were fit using package Lavaan 0.5 (62). BMI was log transformed to increase normality. Models were fit using full information maximum likelihood (FIML) using robust standard errors. Model fit was assessed with the chi-square test (X^2), the Comparative Fit index (CFI), Root Mean Square Error of Approximation (RMSEA) and the standardized root mean squared residuals (SRMR). The following guidelines were utilized for judging good fit: CFI>0.95; RMSEA<0.08 and SRMR <0.10 (63). The hierarchical relationship between impulsivity, UE and BMI was tested using the watershed framework (see supplementary

material, also explained in (46)). Only variables that survived multiple comparisons with False Discovery Rate (FDR) are displayed as significant. The p values reported are FDR corrected.

Results

Eating-specific impulsivity factor uncontrolled eating is correlated with BMI

Participant characteristics are listed in Table 1. We first tested if UE emerged as a latent factor that captures the covariance between the eating questionnaires, namely Disinhibition (TFEQ), Power of Food, and Emotional Eating (DEBQ) (Figure 1-A). We created the model in the first split-half sample, and verified it in the second split-half. Both models resulted in good fits and returned similar loadings (Table S2). Following the best practices in structural equation modeling (64), here we report our model in the full sample. A single factor confirmatory factor analysis for UE provided a good fit for our data. Figure 1-B shows that UE correlated with BMI ($\beta = 0.37, p < 10^{-5}$) and explained 9.2% of the variance in BMI ($F(1, 2206) = 223.7, p < 10^{-5}$).

General impulsivity traits can be explained with a three-factor latent structure and are correlated with BMI

We next tested the relationship between general impulsivity measures and UE and BMI. A data driven exploratory factor analysis in a split-half of the data suggested a three-factor latent factor structure for impulsivity traits (Table S1). The first factor had a negative loading from the Rosenberg self-esteem questionnaire and positive loadings from Neuroticism (IPIP) and the Perceived Stress Scale (PSS). We labelled this factor ‘Stress Reactivity (STRESS)’. The second factor had a positive loading from Lack of Perseverance (UPPS) and a negative loading from Conscientiousness (IPIP). We labelled this latent factor ‘Lack of Self Control’. The third factor

was named ‘Reward Sensitivity’ (REWARD) and it was characterized by positive loadings from the Extraversion (IPIP) and Sensation Seeking (UPPS) subscales (Table 1). We next tested if these general impulsivity domains were associated with BMI. Using confirmatory factor analysis, we showed that there is a small but significant correlation between STRESS and BMI ($\beta = 0.08$, $p=0.028$), and REWARD and BMI ($\beta=0.09$, $p = 0.028$) (Figure 1-C).

The relationship between general impulsivity, uncontrolled eating, and BMI

General impulsivity traits might exert their effects on BMI via UE, eating-specific impulsivity. To test this, we fit the full watershed model integrating the general impulsivity latent factors, the UE latent factor and BMI (Figure 1-D). Our model satisfied the predictions of the watershed model (46, see supplementary materials). These results showed that the relationship between general impulsivity, UE and BMI is hierarchical. Furthermore, our watershed model was selective for eating behaviors. For example, when alcohol intake (i.e. how many alcoholic beverages do you consume a week?) was utilized as the outcome variable in the watershed model, general impulsivity factors, but not UE, explained 25.6% of the variance in alcohol intake (Figure S3).

UE mediates the relationship of stress reactivity and reward sensitivity to BMI

We additionally utilized mediation analysis to test if UE mediates the relationships between STRESS and BMI and REWARD and BMI. Using the structural equation model framework, we simultaneously modelled two pathways: UE mediates (i) the effect of REWARD on BMI and (ii) the effect of STRESS on BMI. The analysis confirmed that the effects of STRESS and REWARD on BMI were mediated by UE (76% mediation; Figure 2; Table S3).

Prefrontal and subcortical brain structures are associated with impulsivity traits

We hypothesized that structural brain measures might be related to the latent impulsivity factors. To test this hypothesis, we included brain regions that have been linked to impulsivity traits and obesity in the full model. Cortical regions were measured using cortical thickness (CT) and anatomical differences in subcortical regions were measured using deformation-based morphometry. We found that STRESS was positively associated with amygdala volume ($\beta = 0.29$, $p = 0.035$) and inversely with hippocampal volume ($\beta = -0.33$, $p = 0.021$) and lateral OFC thickness ($\beta = -0.31$, $p = 0.035$). Lack of self-control inversely correlated with lateral OFC cortical thickness ($\beta = -0.33$, $p = 0.041$). Independent of these relationships, superior frontal gyrus thickness inversely correlated with BMI ($\beta = -0.38$, $p = 0.014$).

Longitudinal Predictors of Weight Gain

We next looked for predictors of change in body weight in the first year of university (Figure 4-A). Our results revealed that students gained on average 0.51 kg ($SD \pm = 3.22$) in the first year of university (August/September to April/May), which is a small but significant effect ($t = 5.4$, $df = 1133$, $p\text{-value} = p < 10^{-5}$, $95\% \text{ CI} = 0.32\text{-}0.68$). We utilized percent BMI change as the outcome variable to account for the initial BMI. Our model explained 1.5 % of the variance in percent BMI change (Figure 4-A). According to our model, UE did not significantly predict BMI change in this sample ($\beta = 0.03$, $p = 0.57$). Among the impulsivity factors, Lack of Self Control predicted weight change ($\beta = 0.10$, $p = 0.028$), while stress reactivity predicted weight loss ($\beta = -0.13$, $p = 0.028$). A post-hoc regression analysis of the relationship between stress reactivity and weight change revealed that this negative correlation was driven by males. In male students,

STRESS explained 2% of the variance in weight reduction ($\beta = -0.17$, $p = 0.0053$, $N = 360$, $R^2 = 0.02$). No significant correlation emerged for female students ($\beta = -0.061$, $p = 0.15$, $N = 788$, $R^2 = 0.001$) (Figure 4-B).

Discussion

The purpose of the current study was twofold (i) to provide a comprehensive link between brain structure, impulsivity and obesity, and (ii) to characterize the temporal relationship between impulsivity and weight changes. We utilized a large sample size ($N > 2000$) of first-year students and accounted for the multidimensional nature of impulsivity to study these relationships. The models were developed in the first split-half of the sample and replicated in the second.

We first stratified impulsivity into two domains: general and eating-specific, and then tested the relationship between these two domains. We showed that the variance of different eating-specific questionnaires can be captured by a general underlying latent factor (UE). UE had a strong relationship to BMI. This result was consistent with previous studies suggesting that different food questionnaires are measuring different severity levels of a common latent factor: eating-specific impulsivity (16, 18). Next, we demonstrated that the commonly used general impulsivity questionnaires can be explained in a three-factor structure that align with the general personality domains of neuroticism (stress sensitivity), conscientiousness (self-control), and extraversion (reward sensitivity) (Figure 1-C). This result is in agreement with the previously published meta-analysis of impulsivity factor structure (5), and highlights the multifaceted nature of impulsivity.

We hypothesized that people with general impulsivity traits might also exhibit eating-specific impulsivity, and that the relationship between general personality domains, UE and BMI would be hierarchical. Using the hierarchical watershed model, we first showed that each general impulsivity domain had an independent and positive correlation with UE (Figure 1-D). While stress reactivity and reward sensitivity weakly correlated with BMI, we showed that this was mediated by UE (Figure 2). This mediation was specific for BMI. For example, increased scores in each general impulsivity domain, but not UE, were linked to increased alcohol intake (Figure S3). This suggests eating-specific impulsivity (UE) captures the relationship between impulsivity and obesity.

Differences in brain structure might underlie differences in impulsivity domains (20). We next tested if brain morphometry related to impulsivity or if it had an independent correlation with BMI. Our results showed that cortical thickness in lateral OFC is inversely related to the self-control and stress reactivity latent factor scores (Figure 3). This is in line of the role of lateral OFC in maintaining goal-oriented focus by suppressing irrelevant information (65–67). Furthermore, lateral OFC and its neighboring ventrolateral PFC have also been implicated in emotional regulation, particularly in the reduction of negative affect (66, 68, 69). Lateral OFC thickness has been shown to be lower in adults with obesity (70). Our results add to this literature and suggest that reduced lateral OFC thickness might affect impulsivity and thereby appetite control and obesity.

Both amygdala and hippocampus have been implicated in the control of food intake as well as in stress reactivity (71–74). In this study we showed that a personality dimension related to stress

reactivity was associated with greater amygdala volume and lower hippocampal volume, consistent with previous reports (26, 27, 75). Our results suggest that differences in amygdala and hippocampal volumes might relate to increased stress reactivity which can lead to UE and obesity. Independent of these relationships, we showed that superior frontal gyrus thickness was inversely related to BMI consistent with the literature (70, 76, 77). This suggests that some brain regions might be important in overeating and obesity independent of impulsivity.

Finally, we tested the hypothesis UE underlies vulnerability for weight gain. We utilized the transition period into university that is associated with high risk of weight gain to study this hypothesis. In our sample, the average weight gain was small, but significant, and was similar to previously reported studies (40). However, in this sample we failed to find that UE was a predictor of weight gain. This could be due to limited duration of the study. Furthermore, the directionality of the UE-BMI relationship could contribute to this result: changes in body weight might lead to longitudinal variations in eating-specific impulsivity. For instance, scores in different uncontrolled eating scales decrease after bariatric surgery (78–81) or voluntary weight loss (82, 83). In this study, weight changes correlated with changes in disinhibition score from TFEQ (has the highest loading UE) between spring and fall semesters ($\beta = 0.088$, $p = 0.02$, $N = 660$), supporting this view.

Our results support a model in which weight gain results from lack of self-control. Upon transitioning into a new environment, difficulties in self-control could compromise the maintenance of behaviors that promote healthy weight. High scores on conscientiousness have been consistently linked to a lower risk of having and developing obesity across independent

samples (84, 85). These results are generally derived from sample populations with a higher mean age than ours, suggesting that self-control deficits may be more associated with BMI as individuals age and start making their own food decision. Our findings thus point to self-control deficits as risk factors for future weight gain in young populations.

Higher stress reactivity correlated with longitudinal weight loss in our study. This result is surprising, and yet, it is consistent with the existence of bidirectional effects of stress on eating behaviors. Stress is associated both with weight-gain and weight-loss longitudinally (86, 87). This bidirectional relationship might be partly mediated by sex differences, since there seem to be biological and psychological variations in stress responses among sexes (88). Male participants that report distress at the transition from high school to university tend to lose weight (89) and decrease their food intake compared to female participants (90). In a similar vein, our post hoc analysis revealed that stress reactivity was associated with weight loss only in male students. Conversely, the stress reactive endophenotype was associated with higher BMI, possibly due to differences in the effects of chronic versus acute stress on appetite and weight gain (91).

The results of our study should be considered with regards to its limitations. Although we had a large sample group, whereby we were able to detect and replicate results from meta-analyses, most of our measures were self-reported. In addition, brain imaging was conducted with a smaller subset from the sample. Therefore, the relationship between impulsivity and brain morphometry were only tested in regions of interest based on the literature. This limits the generalizability of the brain imaging results.

386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410

Conclusions

The present study provides a comprehensive characterization of the relationships between different facets of impulsivity and obesity. We show that structural brain differences relate to differences in impulsivity domains which affect BMI via eating-specific impulsivity (UE). However, predictors of weight gain, at least in the short-term, are different than the risk factors associated with increased BMI. Deficits in self-control, not UE, predict longitudinal weight gain, while higher stress reactivity predicts weight loss in the first year of university (in males). These findings highlight the role of self-control deficits in longitudinal weight gain. Our results might inform future clinical strategies aimed at fostering self-control abilities to prevent and/or treat unhealthy weight gain.

Acknowledgments

This research was supported by a Canadian Institutes of Health Research Grant to AD. SN was supported by a Frederick Banting and Charles Best Canada Graduate Scholarship. UV is supported by Personal Post-doctoral Research Funding project PUTJD654 and by Fonds de recherche du Québec – Santé (FRQS) foreign post-doctoral training award.

Disclosures

SN, UV, MD, YY, IGG, SGS, YZ, DLC, AD reported no financial interests or potential conflicts of interest.

References

1. O’Rahilly S, Farooqi IS (2008): Human obesity: a heritable neurobehavioral disorder that is highly sensitive to environmental conditions. *Diabetes*. 57: 2905–10.
2. Cannon TD, Keller MC (2006): Endophenotypes in the Genetic Analyses of Mental Disorders. *Annu Rev Clin Psychol*. 2: 267–290.
3. Dagher A, Neseliler S, Han JE (2017): Appetite as motivated choice: Hormonal and environmental influences. *Decis Neurosci An Integr Perspect*. pp 397–409.
4. Vainik U, Dagher A, Dubé L, Fellows LK (2013): Neurobehavioural correlates of body mass index and eating behaviours in adults: a systematic review. *Neurosci Biobehav Rev*. 37: 279–99.
5. Sharma L, Markon KE, Clark LA (2014): Toward a theory of distinct types of “impulsive” behaviors: A meta-analysis of self-report and behavioral measures. *Psychol Bull*. 140: 374–408.
6. Dalley JW, Robbins TW (2017): Fractionating impulsivity: neuropsychiatric implications. *Nat Rev Neurosci*. 18.
7. Costa PT, McCrae RR (1992): Professional manual: revised NEO personality inventory (NEO-PI-R) and NEO five-factor inventory (NEO-FFI). *Odessa FL Psychol Assess Resour*.
8. DeYoung CG, Hirsh JB, Shane MS, Papademetris X, Rajeevan N, Gray JR (2010): Testing predictions from personality neuroscience. Brain structure and the big five. *Psychol Sci a J Am Psychol Soc / APS*. 21: 820–828.
9. Gray JA (1987): The Psychology of Fear and Stress. *Probl Behav Sci*. New York, NY: Cambridge University Press., p 422.
10. Michaud A, Vainik U, Garcia-Garcia I, Dagher A (2017): Overlapping Neural

- Endophenotypes in Addiction and Obesity. *Front Endocrinol (Lausanne)*. 8: 127.
11. Emery RL, Levine MD (2017): Questionnaire and behavioral task measures of impulsivity are differentially associated with body mass index: A comprehensive meta-analysis. *Psychol Bull.* 143: 868–902.
12. Meule A (2017): Commentary: Questionnaire and behavioral task measures of impulsivity are differentially associated with body mass index: a comprehensive meta-analysis. *Front Psychol.* 8: 1222.
13. van Strien T, Frijters JER, Bergers GPA, Defares PB (1986): The Dutch Eating Behavior Questionnaire (DEBQ) for assessment of restrained, emotional, and external eating behavior. *Int J Eat Disord.* 5: 295–315.
14. Stunkard AJ, Messick S (1985): The three-factor eating questionnaire to measure dietary restraint, disinhibition and hunger. *J Psychosom Res.* 29: 71–83.
15. Lowe MR, Butryn ML, Didie ER, Annunziato RA, Thomas JG, Crerand CE, *et al.* (2009): The Power of Food Scale. A new measure of the psychological influence of the food environment. *Appetite.* 53: 114–118.
16. Price M, Higgs S, Lee M (2015): Self-reported eating traits: Underlying components of food responsiveness and dietary restriction are positively related to BMI. *Appetite.* 95: 203–10.
17. Mason AE, Vainik U, Acree M, Tomiyama AJ, Dagher A, Epel ES, Hecht FM (2017): Improving Assessment of the Spectrum of Reward-Related Eating: The RED-13. *Front Psychol.* 8: 795.
18. Vainik U, Neseliler S, Konstabel K, Fellows LK, Dagher A (2015): Eating traits questionnaires as a continuum of a single concept. Uncontrolled eating. *Appetite.* 90: 229–239.

- 481 19. Meule A, Blechert J (2017): Indirect effects of trait impulsivity on body mass. *Eat Behav.* 26:
482 66–69.
- 483 20. Allen TA, DeYoung CG (2016): *Personality Neuroscience and the Five Factor Model*. (Vol.
484 1).
- 485 21. Willette AA, Kapogiannis D (2015): Does the brain shrink as the waist expands? *Ageing Res*
486 *Rev.* 20: 86–97.
- 487 22. Matsuo K, Nicoletti M, Nemoto K, Hatch JP, Peluso MAM, Nery FG, Soares JC (2009): A
488 Voxel-Based Morphometry Study of Frontal Gray Matter Correlates of Impulsivity. *Hum*
489 *Brain Mapp.* 30: 1188–1195.
- 490 23. Schilling C, Kühn S, Romanowski A, Schubert F, Kathmann N, Gallinat J (2012): Cortical
491 thickness correlates with impulsiveness in healthy adults. *Neuroimage.* 59: 824–830.
- 492 24. Korponay C, Pujara M, Deming P, Philippi C, Decety J, Kosson DS, *et al.* (2017): Impulsive-
493 antisocial psychopathic traits linked to increased volume and functional connectivity within
494 prefrontal cortex. *Soc Cogn Affect Neurosci.* 12: 1169–1178.
- 495 25. Schilling C, Kühn S, Paus T, Romanowski A, Banaschewski T, Barbot A, *et al.* (2013):
496 Cortical thickness of superior frontal cortex predicts impulsiveness and perceptual
497 reasoning in adolescence. *Mol Psychiatry.* 18: 624–630.
- 498 26. Pruessner M, Bechard-Evans L, Pira S, Joober R, Collins DL, Pruessner JC, Malla AK
499 (2017): Interplay of hippocampal volume and hypothalamus-pituitary-adrenal axis function
500 as markers of stress vulnerability in men at ultra-high risk for psychosis. *Psychol Med.* 47:
501 471–483.
- 502 27. Holmes AJ, Lee PH, Hollinshead MO, Bakst L, Roffman JL, Smoller JW, Buckner RL
503 (2012): Individual differences in amygdala-medial prefrontal anatomy link negative affect,

- 504 impaired social functioning, and polygenic depression risk. *J Neurosci.* 32: 18087–100.
- 505 28. Kapogiannis D, Sutin A, Davatzikos C, Costa P, Resnick S, Resnick S (2013): The five
- 506 factors of personality and regional cortical variability in the Baltimore longitudinal study of
- 507 aging. *Hum Brain Mapp.* 34: 2829–40.
- 508 29. Bjørnebekk A, Fjell AM, Walhovd KB, Grydeland H, Torgersen S, Westlye LT (2013):
- 509 Neuronal correlates of the five factor model (FFM) of human personality: Multimodal
- 510 imaging in a large healthy sample. *Neuroimage.* 65: 194–208.
- 511 30. Cremers HR, Demenescu LR, Aleman A, Renken R, van Tol M-J, van der Wee NJA, *et al.*
- 512 (2010): Neuroticism modulates amygdala-prefrontal connectivity in response to negative
- 513 emotional facial expressions. *Neuroimage.* 49: 963–70.
- 514 31. Omura K, Todd Constable R, Canli T (2005): Amygdala gray matter concentration is
- 515 associated with extraversion and neuroticism. *Neuroreport.* 16: 1905–8.
- 516 32. Bartra O, McGuire JT, Kable JW (2013): The valuation system: a coordinate-based meta-
- 517 analysis of BOLD fMRI experiments examining neural correlates of subjective value.
- 518 *Neuroimage.* 76: 412–27.
- 519 33. Hawes SW, Chahal R, Hallquist MN, Paulsen DJ, Geier CF, Luna B (2017): Modulation of
- 520 reward-related neural activation on sensation seeking across development. *Neuroimage.*
- 521 147: 763–771.
- 522 34. Mokdad AH, Serdula MK, Dietz WH, Bowman BA, Marks JS, Koplan JP (1999): The
- 523 spread of the obesity epidemic in the United States, 1991-1998. *JAMA.* 282: 1519–1522.
- 524 35. Nelson MC, Story M, Larson NI, Neumark-Sztainer D, Lytle LA (2008): Emerging
- 525 adulthood and college-aged youth: an overlooked age for weight-related behavior change.
- 526 *Obesity (Silver Spring).* 16: 2205–2211.

- 527 36. Levitsky DA, Youn T (2004): The More Food Young Adults Are Served, the More They
528 Overeat. *J Nutr.* 134: 2546–2549.
- 529 37. Lloyd-Richardson EE, Bailey S, Fava JL, Wing R, Tobacco Etiology Research Network
530 (TERN) (2009): A prospective study of weight gain during the college freshman and
531 sophomore years. *Prev Med (Baltim).* 48: 256–261.
- 532 38. Pliner P, Saunders T (2008): Vulnerability to freshman weight gain as a function of dietary
533 restraint and residence. *Physiol Behav.* 93: 76–82.
- 534 39. Provencher V, Polivy J, Wintre MG, Pratt MW, Pancer SM, Birnie-Lefcovitch S, Adams GR
535 (2009): Who gains or who loses weight? Psychosocial factors among first-year university
536 students. *Physiol Behav.* 96: 135–141.
- 537 40. Vadeboncoeur C, Townsend N, Foster C (2015): A meta-analysis of weight gain in first year
538 university students: is freshman 15 a myth? *BMC Obes.* 2: 22.
- 539 41. Lowe MR, Annunziato RA, Markowitz JT, Didie E, Bellace DL, Riddell L, *et al.* (2006):
540 Multiple types of dieting prospectively predict weight gain during the freshman year of
541 college. *Appetite.* 47: 83–90.
- 542 42. Vella-Zarb RA, Elgar FJ (2009): The ‘Freshman 5’: A Meta-Analysis of Weight Gain in the
543 Freshman Year of College. *J Am Coll Heal.* 58: 161–166.
- 544 43. Finlayson G, Cecil J, Higgs S, Hill A, Hetherington M (2012): Susceptibility to weight gain.
545 Eating behaviour traits and physical activity as predictors of weight gain during the first
546 year of university. *Appetite.* 58: 1091–1098.
- 547 44. Meule A, Platte P (2018): An Examination of the “Freshman-15” in Germany. *Eur J Heal*
548 *Psychol.* 25: 2–8.
- 549 45. Kievit RA, Davis SW, Mitchell DJ, Taylor JR, Duncan J, Tyler LK, *et al.* (2014): Distinct

- aspects of frontal lobe structure mediate age-related differences in fluid intelligence and multitasking. *Nat Commun.* 5: 5658.
46. Kievit RA, Davis SW, Griffiths J, Correia MM, Cam-Can, Henson RN (2016): A watershed model of individual differences in fluid intelligence. *Neuropsychologia.* 91: 186–198.
47. Donnellan MB, Oswald FL, Baird BM, Lucas RE (2006): The Mini-IPIP Scales: Tiny-yet-effective measures of the Big Five Factors of Personality. *Psychol Assess.* 18: 192–203.
48. Whiteside SP, Lynam DR (2001): The Five Factor Model and impulsivity: using a structural model of personality to understand impulsivity. *Pers Individ Dif.* 30: 669–689.
49. Cohen S, Kamarck T, Mermelstein R (1983): A global measure of perceived stress. *J Health Soc Behav.* 24: 385–396.
50. Rosenberg M (1965): *Society and the adolescent self-image.* .
51. Coupe P, Yger P, Prima S, Hellier P, Kervrann C, Barillot C (2008): An Optimized Blockwise Nonlocal Means Denoising Filter for 3-D Magnetic Resonance Images. *IEEE Trans Med Imaging.* 27: 425–441.
52. Sled JG, Zijdenbos AP, Evans AC (1998): A nonparametric method for automatic correction of intensity nonuniformity in MRI data. *IEEE Trans Med Imaging.* 17: 87–97.
53. Collins DL, Neelin P, Peters TM, Evans AC (1994): Automatic 3D intersubject registration of MR volumetric data in standardized Talairach space. *J Comput Assist Tomogr.* 18: 192–205.
54. Eskildsen SF, Coupé P, Fonov V, Manjón J V., Leung KK, Guizard N, *et al.* (2012): BEaST: Brain extraction based on nonlocal segmentation technique. *Neuroimage.* 59: 2362–2373.
55. Borghammer P, Østergaard K, Cumming P, Gjedde A, Rodell A, Hall N, Chakravarty MM (2010): A deformation-based morphometry study of patients with early-stage Parkinson’s

573 disease. *Eur J Neurol*. 17: 314–320.

574 56. Zeighami Y, Ulla M, Iturria-Medina Y, Dadar M, Zhang Y, Larcher KM-H, *et al.* (2015):

575 Network structure of brain atrophy in de novo Parkinson's disease. *Elife*. 4: 1472–1475.

576 57. Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, *et al.* (2006): An

577 automated labeling system for subdividing the human cerebral cortex on MRI scans into

578 gyral based regions of interest. . doi: 10.1016/j.neuroimage.2006.01.021.

579 58. Team R (2013): R Development Core Team. *R A Lang Environ Stat Comput*. .

580 59. James Honaker A, King G, Blackwell M, Matthew Blackwell M (2018): A Program for

581 Missing Data. . Retrieved from <https://cran.r-project.org/web/packages/Amelia/Amelia.pdf>.

582 60. Revelle W (2016): How To: Use the psych package for Factor Analysis and data reduction. .

583 61. Suhr DD (2006): Exploratory or Confirmatory Factor Analysis? *SAS Users Gr Int Conf*. (pp.

584 1-17).

585 62. Rosseel Y (2012): **lavaan** : An R Package for Structural Equation Modeling. *J Stat Softw*. 48:

586 1–36.

587 63. Kenny DA, Kaniskan B, McCoach DB (2015): The Performance of RMSEA in Models With

588 Small Degrees of Freedom. *Sociol Methods Res*. 44: 486–507.

589 64. Schreiber JB, Nora A, Stage FK, Barlow EA, King J (2006): Reporting Structural Equation

590 Modeling and Confirmatory Factor Analysis Results: A Review. *J Educ Res*. 99: 323–338.

591 65. Vuilleumier P, Armony JL, Driver J, Dolan RJ (2001): Effects of attention and emotion on

592 face processing in the human brain: an event-related fMRI study. *Neuron*. 30: 829–41.

593 66. Hooker CI, Knight RT (2006): Role of the Orbitofrontal Cortex in the Inhibition of Emotion.

594 *The Orbitofrontal Cortex*. 307–324.

595 67. Hong S-B, Kim J-W, Choi E-J, Kim H-H, Suh J-E, Kim C-D, *et al.* (2013): Reduced

- 596 orbitofrontal cortical thickness in male adolescents with internet addiction. *Behav Brain*
597 *Funct.* 9: 11.
- 598 68. Phan KL, Fitzgerald DA, Nathan PJ, Moore GJ, Uhde TW, Tancer ME (2005): Neural
599 substrates for voluntary suppression of negative affect: A functional magnetic resonance
600 imaging study. *Biol Psychiatry.* 57: 210–219.
- 601 69. Wager TD, Davidson ML, Hughes BL, Lindquist MA, Ochsner KN (2008): Prefrontal-
602 Subcortical Pathways Mediating Successful Emotion Regulation. *Neuron.* 59: 1037–1050.
- 603 70. Ottino-González J, Jurado MA, García-García I, Segura B, Marqués-Iturria I, Sender-
604 Palacios MJ, *et al.* (2017): Allostatic Load Is Linked to Cortical Thickness Changes
605 Depending on Body-Weight Status. *Front Hum Neurosci.* 11: 639.
- 606 71. Kanoski SE, Grill HJ (2017): Hippocampus Contributions to Food Intake Control:
607 Mnemonic, Neuroanatomical, and Endocrine Mechanisms. *Biol Psychiatry.* 81: 748–756.
- 608 72. Neseliler S, Han J, Dagher A (2017): The Use of Functional Magnetic Resonance Imaging in
609 the Study of Appetite and Obesity. *Appet Food Intake.* (Vol. 2), pp 117–134.
- 610 73. Stevenson RJ, Francis HM (2017): The hippocampus and the regulation of human food
611 intake. *Psychol Bull.* 143: 1011–1032.
- 612 74. Arnsten AFT (2015): Stress weakens prefrontal networks: molecular insults to higher
613 cognition. *Nat Neurosci.* 18: 1376–1385.
- 614 75. McEwen BS, Bowles NP, Gray JD, Hill MN, Hunter RG, Karatsoreos IN, Nasca C (2015):
615 Mechanisms of stress in the brain. *Nat Neurosci.* 18: 1353–1363.
- 616 76. Marqués-Iturria I, Pueyo R, Garolera M, Segura B, Junqué C, García-García I, *et al.* (2013):
617 Frontal cortical thinning and subcortical volume reductions in early adulthood obesity.
618 *Psychiatry Res Neuroimaging.* 214: 109–115.

- 619 77. Lavagnino L, Mwangi B, Bauer IE, Cao B, Selvaraj S, Prossin A, Soares JC (2016): Reduced
620 Inhibitory Control Mediates the Relationship Between Cortical Thickness in the Right
621 Superior Frontal Gyrus and Body Mass Index. *Neuropsychopharmacology*. 41: 2275–2282.
- 622 78. Ivezaj V, Wiedemann AA, Grilo CM (2017): Food addiction and bariatric surgery: a
623 systematic review of the literature. *Obes Rev*. 18: 1386–1397.
- 624 79. Koball AM, Clark MM, Collazo-Clavell M, Kellogg T, Ames G, Ebbert J, Grothe KB
625 (2016): The relationship among food addiction, negative mood, and eating-disordered
626 behaviors in patients seeking to have bariatric surgery. *Surg Obes Relat Dis*. 12: 165–170.
- 627 80. Sevinçer GM, Konuk N, Bozkurt S, Coşkun H (2016): Food addiction and the outcome of
628 bariatric surgery at 1-year: Prospective observational study. *Psychiatry Res*. 244: 159–164.
- 629 81. Holsen LM, Davidson P, Cerit H, Hye T, Moondra P, Haimovici F, *et al.* (2017): Neural
630 predictors of 12-month weight loss outcomes following bariatric surgery. *Int J Obes*. 1–9.
- 631 82. Martin CK, O’Neil PM, Pawlow L (2006): Changes in Food Cravings during Low-Calorie
632 and Very-Low-Calorie Diets*. *Obesity*. 14: 115–121.
- 633 83. Batra P, Das SK, Salinardi T, Robinson L, Saltzman E, Scott T, *et al.* (2013): Relationship of
634 cravings with weight loss and hunger. Results from a 6month worksite weight loss
635 intervention. *Appetite*. 69: 1–7.
- 636 84. Jokela M, Hintsanen M, Hakulinen C, Batty GD, Nabi H, Singh-Manoux A, Kivimäki M
637 (2013): Association of personality with the development and persistence of obesity: a meta-
638 analysis based on individual-participant data. *Obes Rev*. 14: 315–23.
- 639 85. Sutin AR, Terracciano A (2017): Personality and Body Weight: Mechanisms, Longitudinal
640 Associations and Context. *Pasonariti kenkyu*. 26: 1–11.
- 641 86. Serlachius A, Hamer M, Wardle J (2007): Stress and weight change in university students in

- the United Kingdom. *Physiol Behav.* 92: 548–553.
87. Kivimäki M, Head J, Ferrie JE, Shipley MJ, Brunner E, Vahtera J, Marmot MG (2006):
Work stress, weight gain and weight loss: evidence for bidirectional effects of job strain on
body mass index in the Whitehall II study. *Int J Obes.* 30: 982–987.
88. Verma R, Balhara YPS, Gupta CS (2011): Gender differences in stress response: Role of
developmental and biological determinants. *Ind Psychiatry J.* 20: 4–10.
89. Economos CD, Hildebrandt ML, Hyatt RR (2008): College freshman stress and weight
change: differences by gender. *Am J Health Behav.* 32: 16–25.
90. Zellner DA, Loaiza S, Gonzalez Z, Pita J, Morales J, Pecora D, Wolf A (2006): Food
selection changes under stress. *Physiol Behav.* 87: 789–793.
91. Sominsky L, Spencer SJ (2014): Eating behavior and stress: a pathway to obesity. *Front
Psychol.* 5: 434.

Table 1. Descriptive statistics of the study variables.

2214 Students (34% Males) filled the questionnaires in the fall semester. Out of the 2214 students, 1145 answered the questionnaires in the spring. There were no significant differences in the variables of interest between participants who completed only fall and who completed both fall and spring questionnaires (Figure S1). Paired t-tests revealed that BMI in the spring (mean difference 0.15 kg/m²; t = 4.40, p-value = 1.5*10⁻⁵, 95% CI: 0.08 - 0.22) and weight in the spring (mean difference 0.51 kg; t = 5.32, p< 10⁻⁵, 95% CI: 0.32 - 0.70) were significantly different from the fall values. 95% CI refer to 95 percent confidence intervals. UE: Uncontrolled Eating.

Variables	Mean	SD (±)	min	max
Age	19.8	3.4	18.0	52.0
BMI (kg/m ²)	22.2	3.4	15.5	42.8
Spring BMI (kg/m ²)	22.3	3.3	14.7	44.3
Disinhibition (Three-Factor Eating Questionnaire (14))	6.4	3.4	0.0	16.0
Emotional Eating (Dutch Eating Behavior Questionnaire (13))	2.5	0.7	1.0	4.8
Power of Food (PFS) (15)	12.4	3.7	5.0	25.0
Lack of Perseverance (the UPPS Impulsive Behavior Scale (48))	1.9	0.4	1.0	3.7
Sensation Seeking (the UPPS Impulsive Behavior Scale (48))	2.8	0.6	1.0	4.0
Rosenberg Self Esteem (50)	29.9	5.0	13.0	40.0
Cohen's Perceived Stress Scale (49)	24.5	7.2	2.0	52.0

Neseliler et al. Neural and behavioral endophenotypes of obesity

Neuroticism (International Personality Item Pool (47))	11.6	2.9	4.0	20.0
Agreeableness (International Personality Item Pool(47))	15.6	2.8	4.0	20.0
Imagination (International Personality Item Pool(47))	15.2	2.7	4.0	20.0
Conscientiousness (International Personality Item Pool (47))	14.0	2.9	4.0	20.0
Extraversion (International Personality Item Pool(47))	12.1	3.5	4.0	20.0
Change in weight in first year (kg)	0.51	3.22	-13.0	15.0

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689






Figure 1. Eating-specific impulsivity UE correlates with general impulsivity traits and with BMI. A) A single-factor confirmatory factor analysis for uncontrolled eating fits the data well ($X^2=30.28$, $df=4$, $p<.001$, $CFI=.992$, $RMSEA=.054$, $SRMR=.015$). B) Linear fit of UE with log BMI. C) Confirmatory factor analysis analysis of impulsivity traits with three latent factors fit the data well ($X^2=147.277$, $df=14$, $p<.001$, $CFI=.963$, $RMSEA=.066$, $SRMR=.03$). After multiple comparison correction, STRESS and REWARD are weakly correlated with BMI. D) The full watershed model that links general impulsivity traits to UE and UE to BMI fits the data

well. ($X^2=328.03$, $df=34$, $p<.001$, $CFI=.957$, $RMSEA=.063$, $SRMR=.035$) REWARD: Reward Sensitivity; STRESS: Stress Reactivity; UE: Uncontrolled Eating. B-D represent normalized scores that have been residualized for age and sex. DEBQ: Emotional eating from DEBQ scale; DIS: Disinhibition from TFEQ; PFS: Power of Food Scale; PSS: Perceived Stress Scale; Self-esteem: Rosenberg Self Esteem Scale. Significant relationships after FDR correction for multiple comparisons are shown in blue or yellow based on the directionality of the relationship.

A



B



$$\text{Mediation Effect Size} = \frac{(a1 \times b) + (a2 \times b)}{c1 + c2 + (a1 \times b) + (a2 \times b)} \times 100 = 76 \%$$

Figure 2. Mediation model for BMI. The mediation model tests the hypothesis that the effect of general personality traits- STRESS and REWARD- on BMI is mediated by UE. In mediation models, there are three paths, a, b and c, between the variables. Full mediation will result in the direct relationship between the independent variable and the dependent variable (path c) to become insignificant when taking the mediator into account (path a). The mediation models were

fit together in structural equation model framework but displayed on separate figures as A and B. REWARD: Reward Sensitivity; STRESS: Stress Reactivity; UE: Uncontrolled Eating. Gray numbers represent the regression coefficients between general impulsivity traits and BMI without UE as shown in Figure 1-C. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.




Figure 3. The full watershed model with structural brain parameters. This model fits the data well ($X^2 = 440.485$, $df = 120$, $p < .001$, $CFI = .954$, $RMSEA = .035$, $SRMR = .07$). All of the numbers shown reflect parameters that are standardized and residualized for age and sex. The correlations within the latent factors and within the brain parameters are allowed but not shown for simplicity. Significant relationships after FDR correction for multiple comparisons are shown in blue or yellow based on the directionality of the relationship. Dashed lines represent tested nonsignificant regressions. REWARD: Reward Sensitivity; STRESS: Stress Reactivity; UE: Uncontrolled Eating.







Figure 4. Predictors of weight gain in first year university students. A) This model represents the whole watershed model with percent BMI change as the outcome variable in (N = 1134). The data fits this model well ($X^2=124.855$, $df=34$, $p<.001$, $CFI=.972$, $RMSEA=.048$, $SRMR=.03$). All of the numbers shown reflect parameters that are standardized and residualized for age and sex. Significant relationships after FDR correction for multiple comparisons are shown in blue or yellow based on the directionality of the relationship. B) Regression analysis of STRESS endophenotype with weight change is only significant in male students. Raw data is plotted in panel B. UE: Uncontrolled Eating; STRESS: Stress Reactivity; REWARD: Reward Sensitivity.

Neseliler et al. Supplementary Information




Supplementary Figure 1. Descriptive statistics of participants who completed only the first session in the fall and who completed both sessions. There were no significant differences in the variables of interest between participants between these groups after FDR correction.

Neseliler et al. Supplementary Information



Supplementary Figure 2. Correlation plot of the impulsivity traits used in the exploratory factor analysis. Blue colors refer to positive correlations, red refer to negative correlations. The size of the circle is modulated by the correlation coefficient value. The correlation is shown only for the split half of the data.

Neseliler et al. Supplementary Information



Supplementary Figure 3. Uncontrolled eating does not correlate with alcohol intake.

The full watershed model that links general impulsivity traits to UE and alcohol fits the data well ($X^2=242.058$, $df=43$, $p<.001$, $CFI=.971$, $RMSEA=.046$, $SRMR=.032$). This model explains 25.6 % of the variance in reported alcoholic beverages consumption per week. All of the numbers shown reflect parameters that are fully standardized and residualized for age and sex. Significant relationships after FDR correction for multiple comparisons are shown in blue or yellow based on t

Neseliler et al. Supplementary Information

he directionality of the relationship. UE: Uncontrolled Eating; STRESS: Stress Reactivity; REW
ARD: Reward Sensitivity.

Supplementary Table 1. Results of the exploratory factor analysis.

Parallel analysis is a technique that compares the scree plot of the observed data with that of a random data matrix of the same size as the original (Revelle). Sharp breaks in the scree plot, which plots successive eigenvalues, provide an estimation of the appropriate number of factors to extract. After analyzing the generated pattern matrix, we decided to exclude Imagination, Restraint, Agreeableness from our factor structure. These variables had poor communalities, meaning our factor structure explained a relatively low fraction of the variance for these variables. Re-running our parallel analysis suggested that the optimal number of factors = 3.

Parallel analysis suggests that the number of factors = 3

```
Factor Analysis using method = minres
Call: fa(r = r, nfactors = 3, n.obs = nrow(means))
Standardized loadings (pattern matrix) based upon correlation matrix
```

	item	MR1	MR2	MR3	h2	u2	com
Perceived Stress Scale	7	0.85			0.73	0.27	1.0
Neuroticism	6	0.70			0.46	0.54	1.1
Rosenberg Self Esteem	3	-0.56			0.60	0.40	1.6
Lack of Perseverance	1		-0.84		0.70	0.30	1.0
Conscientiousness	5		0.60		0.43	0.57	1.3
Sensation seeking	2			0.62	0.39	0.61	1.0
Extraversion	4			0.50	0.27	0.73	1.0

Neseliler et al. Supplementary Information

48				
49		MR1	MR2	MR3
50	SS loadings	1.62	1.17	0.78
51	Proportion Var	0.23	0.17	0.11
52	Cumulative Var	0.23	0.40	0.51
53	Proportion Explained	0.45	0.33	0.22
54	Cumulative Proportion	0.45	0.78	1.00
55				
56	With factor correlations of			
57		MR1	MR2	MR3
58	MR1	1.00	-0.39	-0.36
59	MR2	-0.39	1.00	0.11
60	MR3	-0.36	0.11	1.00
61				
62	Tucker Lewis Index of factoring reliability = 0.889			
63	RMSEA index = 0.094 and the 90 % confidence intervals are 0.066 0.124			
64	BIC = 10.88			
65	Fit based upon off diagonal values = 1			
66	Measures of factor score adequacy			
67				

Neseliler et al. Supplementary Information

Supplementary Table 2. Values from the CFA analysis of uncontrolled eating (UE) in the first and second split half of the sample.

	X²	df	P value	CFI	RMSEA	SRMR
UE1	11.83	4	0.019	0.995	0.042	0.014
UE2	19.32	4	0.001	0.990	0.059	0.017

Supplementary Table 3. Mediation Analysis Results

Neseliler et al. Supplementary Information

lavaan (0.5-23.1097) converged normally after 43 iterations

	Used	Total
Number of observations	2208	2213
Number of missing patterns	2	
Estimator	ML	
Minimum Function Test Statistic	328.032	
Degrees of freedom	34	
P-value (Chi-square)	0.000	

Parameter Estimates:

Information	Observed
Standard Errors	Standard

Latent Variables:

	Estimate	Std.Err	z-value	P(> z)	Std.lv	Std.all
STRESS =~						
PSS	1.000				0.691	0.702
Neuroticism	0.736	0.029	25.789	0.000	0.509	0.533
Rosenberg	-1.248	0.054	-23.236	0.000	-0.863	-0.883
Lack of Self Control =~						

Neseliler et al. Supplementary Information

112	Lack of Persever	1.000				0.830	0.846
113	Conscientiousness	-0.726	0.049	-14.704	0.000	-0.603	-0.609
114	REWARD ~						
115	Extraversion	1.000				0.675	0.677
116	SensationSeeking	0.701	0.064	10.987	0.000	0.473	0.492
117	UE ~						
118	Disinhibition	1.000				0.845	0.886
119	PFST	0.820	0.024	33.882	0.000	0.693	0.721
120	DEBQ	0.864	0.024	36.314	0.000	0.730	0.780
121							
122	Regressions:						
123		Estimate	Std.Err	z-value	P(> z)	Std.lv	Std.all
124	UE ~						
125	STRESS (a1)	0.526	0.051	10.286	0.000	0.430	0.430
126	LackofSC	0.140	0.035	4.034	0.000	0.137	0.137
127	REWARD (a2)	0.385	0.058	6.627	0.000	0.308	0.308
128	logBMI ~						
129	STRESS (c1)	-0.085	0.055	-1.555	0.120	-0.059	-0.061
130	LackofSC	-0.025	0.034	-0.724	0.469	-0.021	-0.021
131	REWARD (c2)	-0.014	0.055	-0.257	0.797	-0.010	-0.010
132	UE (b)	0.352	0.031	11.289	0.000	0.297	0.309
133							
134							
135	Mediation Parameters:						
136		Estimate	Std.Err	z-value	P(> z)	Std.lv	Std.all

Neseliler et al. Supplementary Information

137	STRESS to BMI (a1)	-0.085	0.055	-1.555	0.120	-0.059	-0.061
138	REWARD to BMI (a2)	-0.014	0.055	-0.257	0.797	-0.010	-0.010
139	STRESS via UE (c1)	0.185	0.025	7.449	0.000	0.128	0.133
140	REWARD via UE (c2)	0.136	0.024	5.599	0.000	0.091	0.095
141	total effect	0.221	0.093	2.387	0.017	0.151	0.157
142	mediation effect	0.763	0.026	29.443	0.000	0.762	0.762

143

144

145

146

147

148

149

150

151

152

153

154 **Watershed model**

155 In order to study the endophenotypes that related to BMI in a watershed model, we needed to
 156 statistically satisfy three major criteria (as explained in 1, 2). 1) Higher statistical dimensionality

Neseliler et al. Supplementary Information

between upstream variables compared with downstream variables. The statistical dimensionality of the covariance pattern between all variables will increase as we move upstream in the model. The result will be that the upstream effects (i.e. brain or personality traits) cannot be fully captured by a single summary statistic. We tried to capture our cortical thickness variables as a single latent factor (not shown), but this model had a poor fit suggesting that upstream brain variables cannot be reduced to a single dimension. In addition, general impulsivity traits were only captured using three factors as opposed to one latent factor. Furthermore, the upstream effects were partially independent: all of the endophenotypes (i.e. STRESS, Lack of Self Control, REWARD) did better than each individual endophenotype (e.g. STRESS) in predicting the downstream outcome (i.e. UE). 2) Multiple realisability. According to the watershed model, there are multiple precursory causes for a complex endophenotype. In other words, a single behavioural dimension such as UE is likely to have multiple neural determinants. In an SEM framework, this means the variability in each endophenotype (i.e. STRESS, Lack of Self Control, REWARD) will make partially independent contributions to variability in UE endophenotype. 3) Hierarchical dependence. The watershed model should exhibit a hierarchical dependence. That means statistically there should be no residual, or direct, relationships between levels. In the present context this implies that the influence of general personality variables on BMI will be indirect, namely through UE. In the SEM formalization of this hypothesis, any direct paths between general impulsivity domains and BMI will be a source of model misfit, and mediation analysis will reveal either partial or no mediation.

1. Kievit RA, Davis SW, Griffiths J, Correia MM, Cam-Can, Henson RN (2016): A watershed model of individual differences in fluid intelligence. *Neuropsychologia*. 91: 186–198.

2. Kievit RA, Davis SW, Mitchell DJ, Taylor JR, Duncan J, Tyler LK, *et al.* (2014): Distinct

Neseliler et al. Supplementary Information

180 aspects of frontal lobe structure mediate age-related differences in fluid intelligence and
181 multitasking. *Nat Commun.* 5: 5658.

182

183