

Neural dynamics of associative learning during human sleep

Andrés F. Canales-Johnson^{1,2,3}, Emiliano Merlo^{1,4,5}, Tristan A. Bekinschtein¹ and Anat Arzi^{1*}

¹Department of Psychology, University of Cambridge, Cambridge, CB2 3EB, UK.

²Center for Social and Cognitive Neuroscience (CSCN), Universidad Adolfo Ibáñez, Santiago, Chile. ³The Neuropsychology and Cognitive Neurosciences Research Center (CINPSI Neurocog), Universidad Católica del Maule, Talca, Chile.

10 ⁴IFIBIO-Houssay, Facultad de Medicina, Universidad de Buenos Aires - CONICET,
11 Buenos Aires, Argentina. ⁵School of Psychology, University of Sussex, Brighton BN1
12 9RH, UK.

13

14

15

16

17

18

19 Anat Arzi

20 Department of Psychology

21 University of Cambridge

22 Cambridge, CB2 3EB, UK

22 Tel: (+) 44 1223765280

2015 © All rights reserved.

1

27

1 **Abstract**

2 Recent evidence indicate that humans can learn entirely new information during
3 sleep. To elucidate the neural dynamics underlying sleep-learning we investigated
4 brain activity during auditory-olfactory discriminatory associative learning in human
5 sleep. We found that learning-related delta and sigma neural changes are involved in
6 early acquisition stages, when new associations are being formed. In contrast,
7 learning-related theta activity emerged in later stages of the learning process, after
8 tone-odour associations were already established. These findings suggest that
9 learning new associations during sleep is signalled by a dynamic interplay between
10 slow-waves, sigma and theta activity.

11

12 **Introduction**

13 The possibility to learn during sleep has intrigued humanity for over a century. In his
14 1911 science fiction novel “Ralph 124C 41+”, Hugo Gernsback described the
15 *Hypnobioscope*, a device that transmits words directly to the sleeping brain such that
16 they would be fully remembered in the next morning. However, decades of scientific
17 efforts to teach sleeping humans new verbal information have been largely
18 unsuccessful (for review (Peigneux et al. 2001)). Recently, the question of learning
19 during sleep was revisited, and by applying simple forms of learning, such as
20 associative and perceptual learning, it has been found that humans (Arzi et al. 2012,
21 2014; Ruch et al. 2014; Andrillon and Kouider 2016; Andrillon et al. 2017; Züst et al.
22 2019) and animals (De Lavilléon et al. 2015) can learn entirely new information
23 during sleep. Yet, the brain mechanisms enabling learning of novel information

1 during sleep are still unknown. Here, we aimed to identify brain processes supporting
2 discriminatory associative learning during sleep.

3

4 We hypothesized that main brain sleep signals associated with consolidation and
5 reactivation of information learned in awake state (Diekelmann and Born 2010;
6 Oudiette and Paller 2013): slow-waves (delta: 0.5–4 Hz) (Marshall et al. 2006; Rasch
7 et al. 2007; Antony et al. 2012), theta (4-7Hz) (Schreiner and Rasch 2015; Schreiner
8 et al. 2015, 2018) and spindles (sigma: 11-16Hz) (Schabus et al. 2004; Tamminen et
9 al. 2011; Laventure et al. 2016; Cairney et al. 2018), will be associated with learning
10 novel associations during sleep. To test this hypothesis, we analysed
11 electroencephalograph (EEG) activity recorded during auditory-olfactory partial-
12 reinforcement conditioning in non-rapid eye movement (NREM) sleep (Figure 1a-d).
13 On reinforced trials, each tone (400Hz or 1200Hz) was paired with either a pleasant
14 or an unpleasant odour. Tone-odour pairings were counter-balanced across
15 participants. On non-reinforced trials, either tone was presented without an ensuing
16 odour, enabling the measure of learning-related neural correlates without the
17 interference of odour. Stimuli were presented in blocks, each consisted of four
18 reinforced trials and two non-reinforced trials (for detailed experimental design see
19 methods, Figure S1 and Table S1). The conditioned response was the tone-induced
20 sniff response, a behavioural change in nasal airflow in response to tone-odour
21 pairings. Invariably, unpleasant odours drove smaller sniffs than pleasant odours
22 (Arzi et al. 2012). Sleeping participants that learned these tone-odour associations
23 subsequently showed modulated sniffs in response to tones alone, in accordance
24 with the odour valence associated with the tone during sleep [behavioural data

1 published in (Arzi et al. 2012). We found that learning-related delta and sigma
2 activity are involved at early acquisition stages of the learning procedure, while
3 learning-related theta activity emerges only after the discrimination is established,
4 suggesting that timely modulation of slow-waves, sigma and theta rhythms during
5 learning in sleep may prompt the encoding and stabilisation of new associative
6 memories.

7

8 **Materials and Methods**

9

10 The data used here was collected as part of a study that examined whether humans
11 can learn new associations during sleep and was published independently (Arzi et al.
12 2012). Thus, detailed information about participants, experimental design, data
13 acquisition and behavioural results can be found in the original article.

14

15 **Participants**

16 Fourth-three healthy participants (mean age = 25.2 ± 3.2 years, 17 females) gave
17 informed consent to procedures approved by the Weizmann Institute Ethics
18 Committee. Participant exclusion criteria included use of medication, history of sleep
19 disorders and nasal insults, or insufficient sleeping time. Out of these, 28 participants
20 were presented with the auditory-olfactory conditioning during both NREM and REM
21 sleep, and 15 participants during NREM sleep-only. Participants were unaware of
22 specific experimental aims and conditions.

23

24

1 **Stimuli**

2 Pleasant (shampoo or deodorant) and unpleasant (rotten fish or carrion) odorants
3 were presented in a nasal mask by computer-controlled air-dilution olfactometer from
4 an adjacent room (stimulus duration = 3 s, constant flow = 6 litres per minute). Tones
5 (400, 800 and 1,200 Hz, duration = 1 s, at a non-arousing 40 dB) were presented by
6 a loudspeaker ~2 m from participants' heads.

7

8 **Procedures**

9 Participants arrived at the sleep laboratory at a self-selected time, based on their
10 usual sleep pattern, typically at 11:00 pm. After fitting of the polysomnography
11 devices (Iber et al. 2007), subjects were left alone in the darkened room to be
12 observed from the neighbouring control room via infrared video camera. The
13 experimenters observed the real-time polysomnography reading and, after they
14 determined that the subject had entered the desirable sleep stage, they initiated the
15 experimental protocol. The conditioned and unconditioned stimuli were partially
16 reinforced at a ratio of 2:1; on reinforced trials (two-thirds of trials), each 1-s auditory
17 conditioned stimulus (either 1,200 Hz or 400 Hz) was triggered by inhalation and
18 paired with a 3-s olfactory unconditioned stimulus (either pleasant or unpleasant). On
19 non-reinforced trials (one-third of trials), a tone triggered by inhalation was generated
20 without an odorant (tone alone). Stimuli were generated in blocks of six trials (two
21 reinforced trials with pleasant odour, two with unpleasant odour and two non-
22 reinforced trials, one of each tone, randomized between blocks, Inter-trial interval 25-
23 40 seconds). Tone-odour contingencies were counter-balanced across participants.
24 The conditioned response was measured by the sniff response magnitude elicited

1 during tone alone trials. During wakefulness the sniff response can be conditioned to
2 a tone such that different tones can drive different sniffs (Resnik et al. 2011).
3 Therefore, the sniff response was chosen to be the conditioned response in this
4 experiment. In the NREM and REM group (28 participants), in a night without
5 arousals/wakes within a window of 30 s from tone onset, five blocks were presented
6 in NREM sleep, then the procedure was halted up to stable REM sleep, at which
7 point an additional five blocks were presented. In NREM-only, the procedure was
8 triggered during NREM sleep only (15 participants). If an arousal/wake was detected
9 in the ongoing polysomnographic recording, the experiment was immediately
10 stopped until stable sleep was resumed and continued up to a maximum of 18
11 blocks. Thus, the distribution between NREM and REM depended on each
12 participant's sleep structure. Since the experiment was halted following arousal or
13 wake, different participants had different numbers of trials (Mean = 62.9 ± 19.3 trials)
14 and varying training intervals between blocks imposed by individual sleep structure.
15 About half an hour after spontaneous morning wake, conditioned response was
16 tested in a retention procedure: three auditory stimuli, 1,200 Hz and 400 Hz that
17 were presented during the night, and a new 800-Hz tone (eight repetitions each),
18 were sequentially presented while nasal respiration was recorded. Retention
19 procedure data from two subjects was lost due to technical problems.
20

21 **Polysomnography**

22 Sleep was recorded by standard polysomnography (Iber et al. 2007). EEG (obtained
23 from C3 and C4, referenced to opposite mastoid), electro-oculogram (placed 1 cm
24 above or below and laterally of each eye, referenced to opposite mastoid),

1 electromyogram (located bilaterally adjacent to the *submental* muscles), and
2 respiration were simultaneously recorded (Power-Lab 16SP and Octal Bio Amp
3 ML138, ADInstruments) at 1 kHz. Nasal respiration was measured using a
4 spirometer (ML141, ADInstruments) and high-sensitivity pneumotachometer (#4719,
5 Hans Rudolph) in line with the vent ports of the nasal mask.

6

7 **Nasal airflow analysis**

8 Nasal inhalation volume in the retention session was normalised by dividing the sniff
9 volume for each tone by the baseline nasal inhalation volume (averaged volume of
10 15 nasal inhalations preceding retention procedure onset). Participants' normalized
11 sniff volume differing by 3 SD were excluded (one participant).

12

13 **EEG analysis**

14 EEG activity was recorded from C3 and C4 electrodes. Trials with EEG artefacts
15 within a 5.5-second window before tone onset or 5-second after tone onset were
16 excluded. EEG spectral analysis in the 0.5-40 Hz frequency range for all non-
17 reinforced trials that met study criteria was conducted using Hilbert transform on a
18 10.5-second window using customized MATLAB scripts, with time-frequency
19 resolution of 0.5Hz bin per 1 msec. The data was filtered using Hamming window
20 sinc finite impulse response (FIR) filter implemented in EEGLAB. The filter
21 order/transition band width is 25% of lower bandpass edge but not lower than 2Hz
22 where possible. Specifically, Hilbert transform was conducted on a longer time
23 window to avoid edge artefact and the time points before and after the 10.5-second
24 window of interest were trimmed. The power in each trial was then z-scored. Non-

1 reinforced trials from each participant were averaged across the two electrodes, and
2 then averaged across trials to create a single time-frequency representation per
3 participant, and per condition for the early-training phase (averaged across the non-
4 reinforced trials in the first five blocks) and late-training phase (averaged across the
5 non-reinforced trials in the sixth to the last blocks) during NREM sleep.

6

7 **Inclusion/exclusion criteria**

8 An independent experienced sleep technician, blind to experimental conditions and
9 to stimulus onset/offset times, scored the data off-line according to American
10 Academy of Sleep Medicine criteria (Iber et al. 2007). We then used these blindly
11 obtained scorings to include participants and/or trials. We included only EEG
12 artefact-free trials without wake or arousal within 30 s of tone onset presented during
13 NREM sleep. Importantly, trials presented in REM sleep in the 'NREM and REM'
14 group were not included in the analysis. To avoid a bias of the results by individual
15 trials we included in the analysis only participants with a minimum of 10 EEG-clean
16 and arousals-free non-reinforced trials. Out of the 43 participants, five participants
17 had less than 10 EEG-clean and arousals-free non-reinforced trials. Excluding these
18 five participants, data from 38 participants remained for the EEG power analysis. In
19 addition, two participants lacked the retention paradigm (morning testing) due to
20 technical error and one participant's sniff response was an outlier (> 3 SD).
21 Excluding these three participants, data from 35 participants remained for the
22 regression analyses presented in the supplementary materials. Total number of
23 included non-reinforced trials in NREM sleep per learning phase and condition were
24 153 trials for tone alone previously paired with an unpleasant odour (CSu) and 170

1 trials for tone alone previously paired with a pleasant odour (CSp) in early-training,
2 and 146 trials for CSu and 133 for CSp for late-training. The average number of
3 included trials per participant was 4.03 ± 0.94 for CSu and 4.47 ± 0.98 for CSp in
4 early-training, and 3.8 ± 2.9 for CSu and 3.5 ± 2.7 for CSp in late-training. The
5 average number of trials was similar between early- and late-training phases
6 however the variance was larger in the late-training due to individual differences in
7 sleep architecture between participants.

8

9 **Statistical analysis**

10 A permutation-based statistical test of time-frequency data was applied using
11 FieldTrip (Oostenveld et al. 2011) and customized scripts. Time-frequency
12 representation of each condition (CSu and CSp) and learning phase (early and late)
13 in NREM sleep, were submitted to a cluster-based non-parametric permutation test
14 (Maris and Oostenveld 2007), to determine in which time-frequency points in a 5-
15 second window from tone onset there was a significant ($\alpha = 0.05$) change from
16 baseline (-5500 ms to -500 ms pre-tone onset; Figure 2a-d). In addition, the power
17 envelope in each frequency band of interest [delta (0.5-4 Hz), theta (4-7 Hz) and
18 sigma (11-16Hz)] was calculated for each condition separately (CSu and CSp), and
19 submitted to cluster-based non-parametric permutation test to determine in which
20 time points in a 5-second window from tone onset there was a significant increase
21 from baseline per condition per learning phase. To determine whether there was a
22 significant difference in power between CSu and CSp, non-reinforced trials were
23 averaged across conditions (CSu and CSp) and learning phases (early and late).
24 Then the averaged signal was submitted to cluster-based non-parametric

1 permutation test to identify the time points where increase in power following the
2 conditioned stimuli was significantly greater than baseline. A significant cluster was
3 found in each one of the three frequency bands: delta (13 - 2184 ms, $p_{\text{cluster}} < 0.001$),
4 theta (1 - 850 ms, $p_{\text{cluster}} < 0.001$) and sigma (727 - 1553 ms, $p_{\text{cluster}} < 0.001$). The
5 cluster-based non-parametric permutation test between CSu and CSp in each
6 learning phase was applied on the time window defined by the above-mentioned
7 clusters in each frequency band. Multiple regression and correlation analyses were
8 performed using Matlab and open-source statistical software JASP (JASP Team
9 (2017), version 0.8.3.1). Nonparametric effect size was calculated by the following
10 formula $r = z/\sqrt{n}$ (Rosenthal et al. 1994).

11


12 **Results**

13 First, we verified that the behavioural results observed in the current examined
14 NREM sleep dataset (see Methods) were similar to those reported before (The data
15 was collected as part of a previous study published independently; Arzi et al. 2012).
16 In order to investigate learning dynamics during discriminatory associative learning in
17 NREM sleep we first examined the behavioural sniff response dynamics. We
18 extracted the sniff volume during NREM sleep for a tone alone previously paired
19 during sleep with an unpleasant odour (CSu) and for a tone alone previously paired
20 during sleep with a pleasant odour (CSp) up to the fifth non-reinforced trial of each
21 condition or first arousal, whichever came first (see Methods). One can see a
22 gradual increase in CSp-CSu sniff volume difference across trials (Figure 1e).
23 Moreover, the CSp-CSu sniff volume difference was significantly larger in the 4th -
24 5th block in comparison to the 1st - 2nd block ($p < 0.05$), similarly to what was


1 previously shown in a subset of the dataset (Arzi et al. 2012). These findings suggest
2 that the new tone-odour discriminatory associations were learned within the first five
3 blocks. Therefore, we analysed the effects of training during sleep in early- (first five
4 blocks; where discriminatory paired associations were acquired) and late-training
5 (sixth to the last block ($M = 5.5 \pm 3.2$ SD); where paired associations were well
6 established). The following morning, during a test session, we observed that sniff
7 response was larger for CSu than for CSp ($t_{35} = 2.86$, $p = 0.015$; excluding outlier: t_{34}
8 = 2.75, $p = 0.017$; Figure S3c), suggesting the new associations learned during sleep
9 were stored as memories readily retrievable upon awake (Figure 1f).

Differential associative learning experimental design

a Auditory-Olfactory conditioning in sleep



b Reinforced trial:




c Non-Reinforced trial:




d Retention in the morning



e Behavioral sniff-response during sleep



f Behavioral sniff-response in the morning



3 **Figure 1: Auditory-olfactory discriminatory learning paradigm**

4 **(a) Experimental design:** stimuli were generated in blocks of six trials: two
5 reinforced trials with pleasant odour (pink), two reinforced trials with unpleasant
6 odour (grey) and two non-reinforced trials (tone alone), one of each tone (see

1 methods). T, tone; O, Odour. **(b)** On reinforced trials, each auditory stimulus (1,200
2 Hz or 400 Hz) was paired with either a pleasant (shampoo or deodorant) or
3 unpleasant (rotten fish or carrion) odour. **(c)** On non-reinforced trials, either tone was
4 presented alone. **(d)** Structure of the retention session performed during the
5 subsequent morning where three auditory stimuli [1,200 Hz, 400 Hz and a novel 800-
6 Hz tone (green), eight repetitions each] were presented without odours. **(e)**
7 Normalized sniff response across continuous repetitions of a tone alone previously
8 paired during sleep with a pleasant odour (CSp, pink) and continuous repetitions of a
9 tone alone previously paired during sleep with unpleasant odour (CSu, grey) during
10 the first five non-reinforced presentation of each CS during sleep, or first arousal,
11 whichever came first. **(f)** Normalized sniff response during the retention session for
12 CSp (pink bar), novel tone (800Hz, green bar), and for CSu (grey bar). nvu:
13 normalized sniff volume units [sniff volume divided by the baseline nasal inhalation
14 (see Methods)]. The data used here was collected as part of a study that examined
15 whether humans can learn new associations during sleep and was published
16 independently (Arzi et al. 2012).

17

18 Then, to elucidate discriminatory associative learning-related brain dynamics during
19 sleep, we analysed brain activity during non-reinforced CSu and CSp trials during
20 NREM sleep for early- and late-training phases separately. Analysis of delta
21 frequency band showed that during both early- and late-training, CSu elicited higher
22 delta power in comparison to CSp (early-training: cluster1 396-1096 msec, p =
23 0.013, effect size r = 0.60; cluster2 1551-2320 msec, p = 0.011, effect size r = 0.43;
24 Figure 2a,c,e; late-training: cluster 888-1418 msec, p = 0.028, effect size r = 0.42;
25 Figure 2b,d,f), suggesting a learning-related delta modulation. Interestingly, the
26 discriminatory neural response was different between the two training phases. While
27 in the early-training phase two clusters were revealed, in the late-training phase only
28 one cluster was found. While there was overlapping between the first cluster in early-
29 training and the cluster in late-training, the second cluster from early-training
30 uncovered a prolonged learning-related differential response that was not observed
31 in the late-training phase. Furthermore, the differential response between CSu and

1 CSp in early and late-training was similar in the first 1500 msec bout (no cluster) but
2 was larger in early-training compared to late-training in the second 1500 msec bout
3 from tone onset (1959-2273 msec, $p_{\text{cluster}} = 0.05$, effect size $r = 0.31$). These findings
4 indicate a modulation of delta activity between early- and late-training, suggesting
5 this neural correlate could signal the acquisition of new associative memory traces.

6

7 Similar analysis of theta frequency band revealed an opposite response pattern to
8 the one observed in delta. No cluster was found in theta power when comparing CSu
9 and CSp in early-training (Figure 2a, c, g), while late-training CSu elicited higher
10 theta power (cluster 94-311 msec, $p = 0.026$, effect size $r = 0.36$; Figure 2b, d, h)
11 compared to CSp. Moreover, the differential neural response between CSu and CSp
12 was larger in late-training when compared with early-training (cluster = 1-156, msec,
13 $p = 0.05$, effect size $r = 0.37$). These findings suggest that learning-related theta
14 modulations emerge when new associations are well-trained or already established.

15

16 Analysis of sigma frequency band showed that CSu trials induced higher power than
17 CSp homologous events in early-training (cluster 1080-1321 msec, $p = 0.032$, effect
18 size $r = 0.32$; Figure 2a, c, i), but not in late-training (no cluster) (Figure 2b, d, j).
19 However, we did not find reliable differences between early- and late-training in CSu
20 and CSp differential response (no cluster). Thus, modulation in sigma power may
21 underlie acquisition of associative memories in early learning stages, but may not
22 have a distinct contribution to early- versus late-training stages.




Figure 2: Learning-related electrophysiological activity in non-reinforced trials during NREM sleep.

(a-d) Time-frequency decomposition of the EEG signal averaged across C3 and C4 electrodes in non-reinforced trials during NREM sleep time-locked to: **(a-b)** tone previously paired during sleep with an unpleasant odour (CSu) during **(a)** early-training phase, or **(b)** late-training phase; **(c-d)** tone previously paired during sleep

1 with a pleasant odour (CSp) during **(c)** early-training phase, or **(d)** late-training
2 phase. Areas inside black contours indicate significant deviations from zero
3 compared to baseline (cluster permutation *t*-test, $P_{\text{cluster}} < 0.05$), each dotted vertical
4 line represents tone onset, tone duration was 1 sec. EEG delta power during **(e)**
5 early-training and **(f)** late-training for CSu (dark red) and CSp (light red). EEG theta
6 power during **(g)** early-training and **(h)** late-training for CSu (dark green) and CSp
7 (light green). EEG sigma power during **(i)** early-training and **(j)** late-training for CSu
8 (dark blue) and CSp (light blue). Horizontal lines in the colour of the curve represent
9 significant deviation from zero compared to baseline. Horizontal black lines represent
10 significant difference between CSu and CSp (cluster permutation test, $P_{\text{cluster}} < 0.05$).
11
12
13

14 **Discussion**

15 Here, we aimed to elucidate the brain activity supporting discriminatory associative
16 learning in sleep. Using EEG recordings during auditory-olfactory conditioning we
17 uncovered learning-related delta, sigma and theta power modulation in NREM sleep.
18 Moreover, in the course of discriminatory associative learning, learning-related delta
19 and sigma activity are modulated at early acquisition stages while theta activity
20 modulation emerges only after stimuli discrimination is well established. These
21 effects were evident despite the variability in training history introduced by individual
22 differences in sleep architecture (Table S1 and Figure S1).
23

24 During the discriminatory associative learning procedure sleeping participants
25 learned to implicitly discriminate between tones predicting odours with different
26 valence. This process involves learning two independent contingencies between
27 specific tones and odours, and results in the ability to discriminate between the
28 expected value of each tone. During the early-training phase, spanning the first five
29 training blocks, the new tone-odour associations were readily acquired as indicated
30 by the behavioural sniff response (Figure 1e). However, as contingency learning and

1 discrimination occur during the same time frame, and discrimination is an integral
2 part of the learning, the observed learning-related EEG power may reflect either or
3 both of these processes. If a specific brain activity correlate is involved in
4 contingency encoding it should be more apparent when prediction error is high (i.e.
5 during early-training phase). On the contrary, if the same brain activity correlate
6 indicates stimuli discrimination it should be increasingly recruited or stay constant as
7 training progress from early to late trials. Thus, learning-related delta power
8 modulation observed in cluster1 in both early- and late-training suggests an
9 involvement of slow waves in the discrimination process occurring all along training.
10 Modulation of delta power in cluster2 and sigma power observed specifically in early-
11 training implies a role of these brain correlates in acquisition of new associations
12 during sleep. Notably, the contribution of sleep depth to learning cannot be fully
13 dissociated from training phase. In late-training, trials distribution was even between
14 N2 and N3, however in early-training the vast majority of trials were presented in N3
15 (supplementary materials and Table S2), implying that sleep depth may interact with
16 different stages of the learning process. Altogether, these findings suggest that slow-
17 waves and spindles activity is part of the required conditions for encoding of novel
18 associations in sleep.

19
20 To date, only a few studies investigate the neural activity underlying sleep-learning
21 (Peigneux et al. 2001; Ruch et al. 2014; Andrillon and Kouider 2016; Andrillon et al.
22 2017; Farthouat et al. 2018; Züst et al. 2019). The observed learning-related delta
23 power modulation in this study is in line with recent findings showing that successful
24 verbal associative learning during NREM sleep is bound to slow-wave peaks (Ruch

1 et al. 2014; Züst et al. 2019), and that unsuccessful retrieval of auditory perceptual
2 learning during NREM sleep is associated with decreased delta power (Andrillon et
3 al. 2017). In addition, the absence of theta modulation by learning during early-
4 training stage is in agreement with findings of low theta activity during vocabulary
5 encoding in NREM sleep (Züst et al. 2019), and could imply that theta activity does
6 not signal encoding of new associations in NREM sleep. That said, perceptual
7 learning during REM sleep was found to increase theta power (Andrillon et al. 2017).
8 On the other hand, learning-related theta activity in later phases of the training
9 procedure, after the new associations were formed, may be involved in
10 strengthening already-established memories. For sigma, a more complex picture
11 emerges. We observed learning-related modulation in early-training, however not in
12 late-training and found no interaction between learning phases. Other studies, found
13 no relationship between sigma activity and associative learning retrieval (Züst et al.
14 2019), and marginal modulation during perceptual learning (Andrillon et al. 2017).
15 Taken together, these findings suggest that the same brain rhythm may have
16 different roles during different learning processes and sleep stages.

17

18 Delta, sigma and theta activity are key players in memory reactivation and
19 consolidation of information learned before sleep (Diekelmann and Born 2010;
20 Oudiette and Paller 2013). Slow-waves have been associated with and causally
21 related to memory consolidation (Marshall et al. 2006; Rasch et al. 2007; Antony et
22 al. 2012); spindles have a role in integrating new memories and existing knowledge,
23 and in memory consolidation (Diekelmann and Born 2010; Tamminen et al. 2011;
24 Antony et al. 2012, 2019; Oudiette and Paller 2013; Cairney et al. 2018); theta is

1 involved in memory consolidation and reactivation of information learned while
2 awake (Schreiner and Rasch 2015; Schreiner et al. 2015, 2018). Unlike during
3 sleep-learning, the involvement of these brain oscillations in memory formation,
4 consolidation and retrieval is well-established (Diekelmann and Born 2010;
5 Hanslmayr et al. 2016; Antony et al. 2019). However, whether consolidation,
6 targeted memory reactivation and sleep-learning share similar brain mechanism is
7 still an open question.

8

9 Understanding the brain process underlying acquisition, consolidation and retrieval
10 of new information presented during sleep constitutes an important step in the
11 course of identifying what and how is possible to learn during sleep (Simon and
12 Emmons 1956; Wood et al. 1992; Arzi et al. 2012, 2014; Ruch et al. 2014; Andrlillon
13 and Kouider 2016; Andrlillon et al. 2017; Farthouat et al. 2018; Züst et al. 2019).
14 Here, we start to elucidate part of these processes showing that encoding of a
15 discriminatory associative memory during sleep is associated with a dynamic
16 interplay between learning-related slow-waves, sigma and theta activity. Timely
17 modulation of these brain rhythms occurring during learning in sleep may determine
18 the acquisition and storage of new associative memories.

1 **Acknowledgements**

2 This work was supported by the Blavatnik family Foundation [to AA], Royal Society –
3 Kohn International fellowship [NF150851 to AA], European Molecular Biology
4 Organization (EMBO) fellowship [ALTF 33-2016 to AA], Fondo Nacional de
5 Desarrollo Científico y Tecnológico [FONDECYT N 1171200 to ACJ], and The
6 Wellcome Trust [WT093811MA to TAB]. Collection of the data at the Weizmann
7 Institute of Science in Israel was supported by the Rob and Cheryl McEwen Fund for
8 Brain Research. We are indebted to Noam Sobel for his valuable contribution to this
9 work. We are grateful to Sridhar R. Jagannathan, Daniel Bor, Roni Kahana-Zweig,
10 and Lavi Secundo for their comments and for fruitful discussions.

11

12 **Author Contributions**

13 Conceptualization: AFC, EM, TAB, AA; Methodology: AFC, EM, TAB, AA
14 Software: AFC, AA; Validation: AA; Formal analysis: AFC, AA; Investigation: AA
15 Resources: TAB, AA; Data curation: AA; Writing - original draft: AA; Writing – Review
16 & editing: AFC, EM, TAB, AA; Visualization: AFC, EM, TAB, AA; Supervision: AA,
17 TAB; Project Administration: AA; Funding Acquisition: AFC, TAB, AA.

18

19 **Declaration of Interests**

20 The authors declare no competing interests.

21

22 **Data availability**

23 The raw data of this manuscript are available for download at:
24 <https://www.weizmann.ac.il/neurobiology/worg/materials.html>.

25

1 **References**

2 Andrillon T, Kouider S. 2016. Implicit memory for words heard during sleep. *Neurosci*

3 *Conscious.* 2016:niw014.

4 Andrillon T, Pressnitzer D, Léger D, Kouider S. 2017. Formation and suppression of

5 acoustic memories during human sleep. *Nat Commun.* 8.

6 Antony JW, Gobel EW, O'Hare JK, Reber PJ, Paller KA. 2012. Cued memory

7 reactivation during sleep influences skill learning. *Nat Neurosci.* 15:1114–1116.

8 Antony JW, Schönauer M, Staresina BP, Cairney SA. 2019. Sleep Spindles and

9 Memory Reprocessing. *Trends Neurosci.* 42:1–3.

10 Arzi A, Holtzman Y, Samnon P, Eshel N, Harel E, Sobel N. 2014. Olfactory aversive

11 conditioning during sleep reduces cigarette-smoking behavior. *J Neurosci.*

12 34:5382–15393.

13 Arzi A, Shedlesky L, Ben-Shaul M, Nasser K, Oksenberg A, Hairston IS, Sobel N.

14 2012. Humans can learn new information during sleep. *Nat Neurosci.* 15:1460–

15 1465.

16 Cairney SA, Guttesen A á. V, El Marj N, Staresina BP. 2018. Memory Consolidation

17 Is Linked to Spindle-Mediated Information Processing during Sleep. *Curr Biol.*

18 28:948–954.e4.

19 De Lavillénat G, Lacroix MM, Rondi-Reig L, Benchenane K. 2015. Explicit memory

20 creation during sleep demonstrates a causal role of place cells in navigation.

21 *Nat Neurosci.* 18:493–495.

22 Diekelmann S, Born J. 2010. The memory function of sleep. *Nat Rev Neurosci.* 114–

23 126.

24 Farthouat J, Atas A, Wens V, Tiege X De, Peigneux P. 2018. Lack of frequency-

1 tagged magnetic responses suggests statistical regularities remain undetected
2 during NREM sleep. *Sci Rep.* 1–16.

3 Hanslmayr S, Staresina BP, Bowman H. 2016. Oscillations and Episodic Memory:
4 Addressing the Synchronization/Desynchronization Conundrum. *Trends
5 Neurosci.* 39:16–25.

6 Iber C, Ancoli-Israel S, A C. 2007. The AASM manual for the scoring of sleep and
7 associated events: Rules, terminology and technical specifications, American
8 Academy of Sleep Medicine.

9 Laventure S, Fogel S, Lungu O, Albouy G, Sévigny-Dupont P, Vien C, Sayour C,
10 Carrier J, Benali H, Doyon J. 2016. NREM2 and Sleep Spindles Are
11 Instrumental to the Consolidation of Motor Sequence Memories. *PLoS Biol.*
12 14:1–27.

13 Maris E, Oostenveld R. 2007. Nonparametric statistical testing of EEG- and MEG-
14 data. *J Neurosci Methods.* 164:177–190.

15 Marshall L, Helgadóttir H, Mölle M, Born J. 2006. Boosting slow oscillations during
16 sleep potentiates memory. *Nature.* 444:610–613.

17 Oostenveld R, Fries P, Maris E, Schoffelen JM. 2011. FieldTrip: Open source
18 software for advanced analysis of MEG, EEG, and invasive electrophysiological
19 data. *Comput Intell Neurosci.* 2011.

20 Oudiette D, Paller KA. 2013. Upgrading the sleeping brain with targeted memory
21 reactivation. *Trends Cogn Sci.* 17:142–149.

22 Peigneux P, Laureys CAS, Delbeuck X, Maquet P. 2001. Sleeping brain , learning
23 brain . The role of sleep for memory systems. 12:111–124.

24 Rasch B, Büchel C, Gais S, Born J. 2007. Odor cues during slow-wave sleep prompt

1 declarative memory consolidation. *Science* (80-). 315:1426–1429.

2 Resnik J, Sobel N, Paz R. 2011. Auditory aversive learning increases discrimination

3 thresholds. *Nat Neurosci*. 14:791–796.

4 Rosenthal R, Cooper H, Hedges L. 1994. Parametric measures of effect size. *Handb*

5 *Res Synth*. 621:231–244.

6 Ruch S, Koenig T, Mathis J, Roth C, Henke K. 2014. Word encoding during sleep is

7 suggested by correlations between word-evoked up-states and post-sleep

8 semantic priming. *Front Psychol*. 5:1–14.

9 Schabus M, Gruber G, Parapatics S, Sauter C, Klösch G, Anderer P, Klimesch W.

10 2004. Sleep Spindles and Their Significance for Declarative Memory

11 Consolidation. *Sleep*. 27:1479–1485.

12 Schreiner T, Doeller CF, Jensen O, Rasch B, Staudigl T. 2018. Theta Phase-

13 Coordinated Memory Reactivation Reoccurs in a Slow-Oscillatory Rhythm

14 during NREM Sleep. *Cell Rep*. 296–301.

15 Schreiner T, Lehmann M, Rasch B. 2015. Auditory feedback blocks memory benefits

16 of cueing during sleep. *Nat Commun*. 6.

17 Schreiner T, Rasch B. 2015. Boosting vocabulary learning by verbal cueing during

18 sleep. *Cereb Cortex*. 25:4169–4179.

19 Simon CW, Emmons WH. 1956. Responses to material presented during various

20 levels of sleep. *J Exp Psychol*. 51:89–97.

21 Tamminen J, Payne JD, Stickgold R, Wamsley EJ, Gareth M. 2011. Sleep spindle

22 activity is associated with the integration of new memories and existing

23 knowledge. *J Neurosci*. 30:14356–14360.

24 Wood JM, Bootzin RR, Kihlstrom JF, Schacter DL. 1992. Implicit and Explicit

- 1 Memory for Verbal Information Presented during Sleep. *Psychol Sci*. 3:236–240.
- 2 Züst MA, Ruch S, Wiest R, Henke K. 2019. Implicit Vocabulary Learning during
- 3 Sleep Is Bound to Slow-Wave Peaks. *Curr Biol*. 1–13.
- 4