

1 Chikungunya virus outbreak in the Amazon region: replacement of 2 the Asian genotype by an ECSA lineage?

3

4 Naveca, F. G. ¹, Claro, I. ², Giovanetti, M. ^{3,4}, Jesus, J. G. ^{3,5}, Xavier, J. ^{3,5}, Iani, F. C.
5 M. ^{4,6}, do Nascimento V. A. ¹, de Souza, V. C. ¹, Silveira, P. P. ⁷, Lourenço, J. ⁸,
6 Santillana, M. ^{9,10}, Kraemer, M. U. G. ^{8,11}, Quick, J. ¹², Hill, S. C. ⁸, Thézé, J. ⁸,
7 Carvalho, R. D. O. ⁴, Azevedo, V. ⁴, Salles, F. C. S. ², Nunes, M. R. T. ¹³, Lemos, P. S.
8 ¹³, Candido, D. S. ⁸, Pereira, G. C. ⁴, Oliveira, M. A. A. ⁴, Meneses, C. A. R. ¹⁴, Maito,
9 R. M. ¹⁴, Cunha, C. R. S. B. ¹⁴, Campos, D. P. S. ¹⁵, Castilho, M. C. ¹⁶, Siqueira, T. C.
10 S. ¹⁵, Terra, T. M. ¹⁷, de Albuquerque, C. F. C. ¹⁸, da Cruz, L. N. ¹⁹, Abreu, A. L. ¹⁹,
11 Martins, D. V. ¹⁹, Simoes, D. S. M. V. ¹⁹, Aguiar, R. S. ⁷, Luz S. L. B. ¹, Loman, N. ¹²,
12 Pybus, O. G. ⁸, Sabino, E. C. ², Okumoto, O. ¹⁹, Alcantara, L. C. J. ^{3,4*}, Faria, N. R. ^{8*}

13

14 1. Laboratório de Ecologia de Doenças Transmissíveis na Amazônia, Instituto
15 Leônidas e Maria Deane, FIOCRUZ, Manaus, Brazil.

16 2. Instituto de Medicina Tropical e Faculdade de Medicina da Universidade de São
17 Paulo, São Paulo, Brazil.

18 3. Laboratório de Flavivírus, Instituto Oswaldo Cruz, FIOCRUZ, Rio de Janeiro,
19 Brazil.

20 4. Laboratório de Genética Celular e Molecular, Instituto de Ciências Biológicas,
21 Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais, Brazil.

22 5. Laboratório de Patologia Experimental, Instituto Gonçalo Moniz, Fundação
23 Oswaldo Cruz, Salvador, Bahia, Brazil.

24 6. Laboratório Central de Saúde Pública, Instituto Octávio Magalhães, FUNED, Belo
25 Horizonte, Minas Gerais, Brazil.

26 7. Laboratório de Virologia Molecular, Departamento de Genética, Instituto de
27 Biologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil.

28 8. Department of Zoology, University of Oxford, South Parks Road, Oxford, United
29 Kingdom.

30 9. Harvard Medical School, Department of Pediatrics, Boston, MA, USA.

31 10. Computational Health Informatics Program, Boston Children's Hospital, Boston,
32 MA, USA.

11. Computational Epidemiology Lab, Boston Children's Hospital, Boston, MA, USA.
 12. Institute of Microbiology and Infection, University of Birmingham, Birmingham B15 2TT, United Kingdom.
 13. Evandro Chagas Institute, Brazilian Ministry of Health, Ananindeua, Brazil.
 14. Laboratório Central de Saúde Pública, Boa Vista, Roraima, Brazil.
 15. Superintendência de Vigilância em Saúde, Secretaria Municipal de Saúde de Boa Vista, Roraima, Brazil.
 16. Departamento de Virologia, Fundação de Medicina Tropical Doutor Heitor Vieira Dourado, Manaus, Amazonas, Brazil.
 17. Laboratório Central de Saúde Pública do Amazonas, Manaus, Amazonas, Brazil.
 18. Organização Pan - Americana da Saúde/Organização Mundial da Saúde - (OPAS/OMS), Brasília-DF, Brazil.
 19. Secretaria de Vigilância em Saúde, Ministério da Saúde (SVS/MS), Brasília-DF, Brazil.
- Corresponding authors:** luiz.alcantara@ioc.fiocruz.br and nuno.faria@zoo.ox.ac.uk

Abstract

Background

Since its first detection in the Caribbean in late 2013, chikungunya virus (CHIKV) has affected 51 countries in the Americas. The CHIKV epidemic in the Americas was caused by the CHIKV-Asian genotype. In August 2014, local transmission of the CHIKV-Asian genotype was detected in the Brazilian Amazon region. However, a distinct lineage, the CHIKV-East-Central-South-America (ECSA)-genotype, was detected nearly simultaneously in Feira de Santana, Bahia state, northeast Brazil. The genomic diversity and the dynamics of CHIKV in the Brazilian Amazon region remains poorly understood despite its importance to better understand the epidemiological spread and public health impact of CHIKV in the country.

Methodology/Principal Findings

We report a large CHIKV outbreak (5,928 notified cases between August 2014 and August 2018) in Boa vista municipality, capital city of Roraima's state, located in the Brazilian Amazon region. In just 48 hours, we generated 20 novel CHIKV-ECSA genomes from the Brazilian Amazon region using MinION portable genome sequencing. Phylogenetic analyses revealed that despite an early introduction of the Asian genotype in 2015 in Roraima, the large CHIKV outbreak in 2017 in Boa Vista was caused by an ECSA-lineage most likely introduced from northeastern Brazil. Epidemiological analyses suggest a basic reproductive number of R_0 of 1.66, which translates in an estimated 39 (95% CI: 36 to 45) % of Roraima's population infected with CHIKV-ECSA. Finally, we find a strong association between Google search activity and the local laboratory-confirmed CHIKV cases in Roraima.

Conclusions/Significance

This study highlights the potential of combining traditional surveillance with portable genome sequencing technologies and digital epidemiology to inform public health surveillance in the Amazon region. Our data reveal a large CHIKV-ECSA outbreak in Boa Vista, limited potential for future CHIKV outbreaks, and indicate a replacement of the Asian genotype by the ECSA genotype in the Amazon region.

Author Summary

Until the end of 2017, Brazil notified the highest number of infections caused by chikungunya virus (CHIKV) in the Americas. We investigated a large CHIKV outbreak in Boa vista municipality in the Brazilian Amazon region. Rapid portable genome sequencing of 20 novel isolates and subsequent genetic analysis revealed that ECSA lineage was introduced from northeastern Brazil to Roraima around July 2016. Epidemiological analyses suggest a basic reproductive number of R_0 of 1.66, which suggests that approximately 39% of Roraima's population was infected with CHIKV-ECSA. Given the dominance of the CHIKV-Asian genotype in the Americas, our data highlights the rapid spread of a less understood and poorly characterized CHIKV-ECSA genotype in Brazil. Investigations on potential associations between public health impact of CHIKV and genetic diversity of circulating strains are warranted to better evaluate its impact in Brazil and beyond.

Keywords

Chikungunya, East-Central-South-African, surveillance, Amazon region, MinION genome sequencing, traditional epidemiology, genomic epidemiology, digital epidemiology, phylodynamics.

Introduction

In August 2014, local transmission of chikungunya virus (CHIKV) was detected in Brazil for the first time, with cases being reported nearly simultaneously in Oiapoque (Amapá state, north Brazil) and Feira de Santana (Bahia state, northeast Brazil), two municipalities separated by >2000 km distance. Genetic analysis confirmed the co-circulation of distinct virus lineages in Brazil: the Asian genotype (CHIKV-Asian) was introduced to Oiapoque possibly from neighbouring French Guiana, while the East-Central-South-African genotype (CHIKV-ECSA) was introduced to Feira de Santana from a traveller returning from Angola [1].

Since 2014 and until the end of September 2018, a total of 697,564 CHIKV cases have been notified in Brazil (including 94,672 laboratory-confirmed cases). This is the largest number recorded in any of the 51 countries or territories reporting local CHIKV transmission in the Americas [2]. The virus has been circulating in the Americas since 2013 where approximately 260 million people live in areas at-risk of transmission [2-4]. Similar to the recent Zika virus epidemic [5], the rapid spread of CHIKV in the Americas, including in Brazil, results from several factors, including the establishment and abundance of competent *Aedes* spp. vectors, lack of population immunity, and increased mobility of vectors and humans between regions reporting current presence of the virus [6].

Chikungunya virus is an enveloped, non-segmented, single-stranded positive polarity RNA alphavirus that is a member of the *Togaviridae* family and is transmitted predominately by the *Aedes aegypti* and *Aedes albopictus* vectors, which are widespread in Brazil [7]. There are four main genotypes: (i) the West African genotype is maintained in an enzootic cycle in Africa, (ii) the Asian genotype, which is endemic in Asia, (iii) the East-Central-South-African genotype, endemic to Africa, and (iv) the Indian Ocean Lineage (IOL) genotype, an epidemic lineage that emerged from the ECSA genotype around 2004 and swept through the Indian Ocean region causing a series of explosive outbreaks [8].

The first symptoms of CHIKV infection are a rapid increase in temperature (>38.9°C), followed by severe, often debilitating polyarthralgia. Serological data from La Reunion, Philippines and the Indian Ocean island of Mayotte suggest that 75-97% of persons infected with CHIKV develop symptomatic infections [9]. Seroprevalence data from Brazil suggests that 45.7 to 57.1% Riachão do Jacuípe

and of Feira de Santana, both located in Bahia state, were exposed to CHIKV in 2015, with a total of 32.7% to 41.2% of the population reporting symptoms [10].

Throughout Asia and the Americas, chikungunya virus outbreaks have been associated with unique clinical features [11], including long-lasting symptoms [12], and high mortality resulting from complications associated with CHIKV infection [13, 14]. In Brazil, a striking proportion of 68.1 to 75% of the population with positive serological results reporting symptoms contracted a chronic form of the disease [12, 15]. However, the epidemiological features, genomic diversity, and transmission dynamics of recent CHIKV outbreaks in this country remain poorly understood. Inferences that are based only on clinical-epidemiological notifications are complicated by underreporting of cases by the national reporting system [16], mostly due to the co-circulation and co-infection with viruses that cause overlapping symptoms, such as Zika and dengue viruses [17-19]. Moreover, CHIKV serological tests may cross-react with other alphaviruses, such as Mayaro virus, that circulate in the north and centre-west regions of Brazil [20, 21]. In this context, it is challenging to use only clinical-epidemiological and serological data to evaluate the true extent of the disease. Moreover, accurate incidence data is critical to forecast and provide prediction of the course of epidemics [22].

Until the end of 2016, 83.3% of the cases in Brazil were reported in northeast region of the country [23]. However, in 2017, Roraima state, located in the Amazon basin in the north of Brazil, reported its first large CHIKV outbreak. Roraima is the northernmost state of Brazil, lies in the Amazon basin, borders Venezuela and French Guiana to the north, and Amazonas and Pará states to the south, and its equatorial climate favours year round transmission of mosquito-borne viruses [24]. Within Brazil's northern states, Roraima has been implicated as a stepping-stone to virus introductions from other Latin American regions, such as dengue [25], and yellow fever virus in the past [26]. Moreover, the Amazon region has recently been highlighted as a region with high transmission potential of vector-borne diseases [4] and, more generally, a region with high potential for virus zoonoses and emergence [27].

Due to its connectivity and potential impact on global epidemiology of vector-borne and zoonotic virus from the Amazon basin, it is important to improve genomic pathogen surveillance in Roraima. By August 2018, the public health laboratory of Boa Vista (capital city of Roraima state) had reported 5,928 CHIKV cases, 3,795 of

171 which were laboratory-confirmed. Here we use a combination of on-site portable
172 virus genome sequencing, and epidemiological analysis of case count and web search
173 data to describe the circulation, genetic diversity, epidemic potential and attack rates
174 of a large CHIKV outbreak in Boa Vista.

175

Methods

Connectivity in study area

Roraima is the northernmost of Brazil's 27 federal units (**Figure 1a**) and has an estimated population of 450,479, of whom 284,313 live in the capital city of Boa Vista (ibge.gov.br/). Despite being Brazil's least populated federal unit, Roraima is one of the best-connected Brazilian states in the Amazon basin [28]. Within Brazil, Roraima is connected to Amazonas state in the south via the road BR-174. This road also connects Roraima's capital city, Boa Vista, to the states of Bolivar and Amazonas in Venezuela in the north. Further, the road BR-401 links Boa Vista to Guyana in the east. There are four daily flights connecting Boa Vista with Brasília, capital of Brazil, as well as six weekly flights to Manaus, the capital city of Amazonas state and the biggest city in the north of the country, with connecting daily nonstop flights to all other Brazilian states/regions and international destinations, including important international airport hubs in Panamá City and Miami, USA. There are also less-commonly used seasonal fluvial networks that connect Boa Vista and Manaus via the Amazonas river.

[Figure 1 around here]

Chikungunya virus case count time series

The Roraima State Central Laboratory (LACEN-RR) is responsible for the differential diagnosis of suspected arbovirus cases presenting to Roraima's public health units. Between Jan 2014 and September 2018, LACEN-RR notified 5,928 CHIKV cases in Boa Vista alone, 3,795 of these laboratory-confirmed, to the National Reportable Disease Information System (SINAN). Case count time series are available from Github (<http://github.com/arbospread>). We follow the Brazilian Ministry of Health's guidelines and define a notified CHIKV case as a suspected case characterized by (i) acute onset of fever $>38.5^{\circ}\text{C}$, (ii) severe arthralgia and/or arthritis not explained by other medical conditions, and (iii) residing or having visited epidemic areas within 15 days before onset of symptoms. A laboratory-confirmed case is a suspected case confirmed by laboratory methods such as (i) virus isolation in cell culture, (ii) detection of viral RNA, (iii) detection of virus-specific IgM antibodies in a single serum sample collected in the acute or convalescent stage of infection; or (iv) a four-

fold rise of IgG titres in samples collected during the acute phase, in comparison with a sample collected in the convalescent period.

Nucleic acid isolation and RT-qPCR

Residual anonymized clinical diagnostic samples were sent to Instituto Leônidas e Maria Deane, FIOCRUZ Manaus, Amazonas, Brazil, for molecular diagnostics as part of the ZiBRA-2 project. The ZiBRA-2 project was approved by the Pan American Health Organization Ethics Review Committee (PAHOERC) nº PAHO-2016-08-0029. Total RNA extraction was performed with QIAmp Viral RNA Mini kit (Qiagen), following manufacturer's recommendations. Samples were first tested using a multiplexed qRT-PCR protocol against CHIKV, dengue virus (DENV1-4), yellow fever virus, Zika virus, Oropouche virus and Mayaro virus [29]. All qRT-PCR results were corroborated using a second protocol [30]; comparable Ct values were obtained with the two protocols. CHIKV positive samples tested negative for all other arboviruses tested. Samples were selected for sequencing based on Ct-value <30 (to maximize genome coverage of clinical samples by nanopore sequencing [31]), and based on the availability of epidemiological metadata, such as date of onset of symptoms, date of sample collection, gender, municipality of residence, and symptoms (**Table 1**). We included 13 samples from Roraima state plus 5 additional samples from patients visiting the LACEN-Amazonas in Manaus, under the auspices of the ZiBRA project (<http://www.zibraproject.org/>). All samples were processed in accordance with the terms of Resolution 510/2016 of CONEP (National Ethical Committee for Research, Brazilian Ministry of Health).

Complete genome MinION nanopore sequencing

Between the 1st and 7th June 2017, we attempted sequencing at Instituto Leônidas e Maria Deane, FIOCRUZ Manaus on all selected samples with Ct-value <30. We used an Oxford Nanopore MinION device with protocol chemistry R9.4, as previously described [32]. Sequencing statistics can be found in **Table S1 (Julien)**. In brief, we employed a protocol with cDNA synthesis using random primers followed by strain-specific multiplex PCR [32]. Extracted RNA was converted to cDNA using the Protoscript II First Strand cDNA synthesis Kit (New England Biolabs, Hitchin, UK) and random hexamer priming. CHIKV genome amplification by multiplex PCR was attempted using the CHIKAsianECSA primer scheme and 35 cycles of

PCR using Q5 High-Fidelity DNA polymerase (NEB) as described in [32]. PCR products were cleaned up using AmpureXP purification beads (Beckman Coulter, High Wycombe, UK) and quantified using fluorimetry with the Qubit dsDNA High Sensitivity assay on the Qubit 3.0 instrument (Life Technologies). PCR products for samples yielding sufficient material were barcoded and pooled in an equimolar fashion using the Native Barcoding Kit (Oxford Nanopore Technologies, Oxford, UK). Sequencing libraries were generated from the barcoded products using the Genomic DNA Sequencing Kit SQK-MAP007/SQK-LSK108 (Oxford Nanopore Technologies). Libraries were loaded onto a R9/R9.4 flow cell and sequencing data were collected for up to 48 hr. Consensus genome sequences were produced by alignment of two-direction reads to a CHIKV virus reference genome (GenBank Accession number: N11602) as previously described in [32]. Positions with $\geq 20\times$ genome coverage were used to produce consensus alleles, while regions with lower coverage, and those in primer-binding regions were masked with N characters. Validation of the sequencing protocol was previously performed in [32].

Collation of CHIKV-ECSA complete genome datasets

Genotyping was first conducted using the phylogenetic arbovirus subtyping tool available at <http://www.krisp.org.za/tools.php>. Complete and near complete sequences were retrieved from GenBank on June 2017 [33]. Two complete or near-complete CHIKV genome datasets were generated. Dataset 1 included ECSA-PreAm (ECSA sampled outside the Americas) and ECSA-Br (ECSA sequences sampled in the Americas) sequences. This dataset contained 36 complete genomes from the ECSA genotype, including 7 from East and Central Africa (HM045823 from Angola 1962; HM045784 from Central African Republic 1984; HM045812 from Uganda 1982; KY038947 from Central African Republic 1983; HM045793 from Central African Republic 1986; HM045822 from Central African Republic 1978; and KY038946 from Central African Republic 1975). Dataset 1 also included 29 sequences from Brazil, including the new 18 genomes reported here from the ECSA lineage and 3 genomes from the outbreak caused by the ECSA lineage in June 2016 in Maceió, Alagoas states, northeast Brazil (**Figure 1a**) [34]. Dataset 2 (ECSA-Br) included only the 29 Brazilian genome sequences. Using a robust nonparametric test [35], no evidence of recombination was found in both datasets.

Maximum likelihood analysis and temporal signal estimation

Maximum likelihood (ML) phylogenetic analyses were performed for each dataset using RAxML v8 [36]. We used a GTR nucleotide substitution model with 4 gamma categories (GTR+4 Γ). In order to investigate the evolutionary temporal signal in each dataset, we regressed root-to-tip genetic distances against sample collection dates using TempEst [37]. For both datasets we obtained a strong linear correlation (dataset 1: $r^2=0.93$; dataset 2: $r^2=0.84$) suggesting these alignments contain sufficient temporal information to justify a molecular clock approach. However, for dataset 1, the Angola/M2022/1962 strain was positioned substantially above the regression line. Previous investigations have suggested this strain may have been the result of contamination or high passage in cell culture [8], so this sequence was removed from subsequent analyses.

Molecular clock phylogenetic analysis

To estimate time-calibrated phylogenies we used the BEAST v.1.10.1 software package [38]. To infer historical trends in effective population size from the genealogy we used several different coalescent models. Because preliminary analysis indicated oscillations in epidemic size through time (as also expected from national case report data), we used three flexible, non-parametric models: a) the standard Bayesian skyline plot (BSP; 10 groups) [39], b) the Bayesian skyride plot [40], and c) the Bayesian skygrid model [41], with 45 grid points equally spaced between the estimated TMRCA of the CHIKV-ECSA genotype in Brazil and the date of the earliest available isolate, collected in 18 March 2017 [41]. For comparison, we also used a constant population size coalescent model. We tested two molecular clock models: a) the strict molecular clock model, which assumes a single rate across all phylogeny branches, and b) the more flexible uncorrelated relaxed molecular clock model with a lognormal rate distribution (UCLN) [42]. Because the marginal posterior distribution of the coefficient of variation of the UCLN model did not exclude zero (most likely due to the small alignment size), we used a strict molecular model in all analyses. For each coalescent model, Markov

Chain Monte Carlo analyses were run in duplicate for 10 million steps using a ML starting tree, and the GTR+4 Γ codon partition (CP)1+2,3 model [42].

Epidemiological analysis

The epidemic basic reproductive number (R_0) was estimated from monthly confirmed cases, as previously described [31, 43]. Because (i) the Asian genotype was circulating in the north region of Brazil since 2014 [1], and (ii) we observed a relatively small number of cases both in the notified and confirmed time series, we assume cases from June 2014 and December 2016 did not represent autochthonous transmission of CHIVK-ECSA. We assume a mean generation time of 14 days, as previously reported elsewhere [44]. We report R_0 estimates for different values of the generation time (g) parameter, along with corresponding estimates of the epidemic exponential growth rate, per month (r).

Web search query data

Available in near-real time, disease-related Internet search activity has been shown to track disease activity (a) in seasonal mosquito-borne disease outbreaks, such as those caused by dengue [42, 82], and (b) in unexpected and emerging mosquito-borne disease outbreaks such as the 2015-2016 Latin American Zika outbreak [45]. Here, we investigated whether we could find a meaningful relationship between Internet search activity and the local chikungunya outbreak in Roraima. Indeed, novel Internet-based data sources have the potential to complement traditional surveillance by capturing early increases in disease-related search activity that may signal an increase in the public's perception of a given public health threat and may additionally capture underlying increases in disease activity. Internet searches may be particularly important and indicative of changes in disease transmission early during an outbreak, when ongoing information on the virus transmission is obfuscated by a lack of medical surveillance. In addition, Internet search trends may also help track disease activity in populations that may not seek formal medical care. We used the Google Trends (GT) tool [45] to compile the monthly fraction of online searches for the term "Chikungunya", that originated from Boa Vista municipality (Roraima state), between January 2014 and July 2018. For comparison, GT search activity for the term "Chikungunya" was collected for the

same time period for Manaus municipality (Amazonas state). The synchronicity of GT time series and notified and confirmed case counts from Boa Vista and Manaus was assessed using the Spearman's rank correlation test in the R software [46].

Data availability

XML files and datasets analysed in this study are available in the GitHub repository (<http://github.com/arbospread>). New sequences have been deposited in GenBank under accession numbers MK121891-MK121908 (CHIKV-ECSA) and MK134712-MK134713 (CHIKV-Asian).

Results

Although most CHIKV notified cases in Brazil were reported in 2016 (**Figure 1**), in Roraima, the majority of notified and confirmed cases in Roraima state were reported in 2017 (5,027 notified cases and 3,720 laboratory-confirmed infections). The number of cases in Roraima started increasing exponentially in January 2017, and the outbreak peaked in July 2017.

Figure 1 around here

We attempted on-site portable nanopore sequencing of isolates collected during the early phase of the outbreak (February to March 2017). We selected 15 RT-qPCR+ virus isolates from autochthonous cases in Roraima state (11 from Boa Vista, 1 from Bonfim, and 1 from Iracema municipalities) (**Table 1**) with a cycle threshold (Ct) ≤ 30 (mean 20.3, range 13.7 – 27.41). We included two isolates from two infected travellers returning to Roraima in December 2014, and an additional five isolates from Amazonas state (all from Manaus municipality), sampled between July 2015 and March 2017. In less than 48 hours genome sequence data was obtained for all selected isolates and in less than 72 hours preliminary results were shared with local public health officials and the Brazilian Ministry of Health. A mean genome coverage of 86% (20x) per base pair was obtained for the sequenced data; mean coverage increased to 90% when focusing on samples with Ct < 26 (**Figure 2a**). Coverage of individual sequences and epidemiological information for each sequenced isolate can be found in **Table 1**.

Table 1 around here

Manual and automated phylogenetic analysis identified the ECSA genotype as the dominant genotype circulating in both Roraima and Manaus between 2015 and 2017. However, two cases from late 2014 returning from Venezuela to Roraima (AMA294 and AMA295) were classified as Asian genotype, the dominant lineage circulating in Latin America. Regression analysis of genetic divergence and sampling dates shows accumulation of temporal signal in the ECSA-Br dataset ($r^2 = 0.84$) (**Figure 2b**).

We estimated the evolutionary time-scale of the ECSA-Br lineage using several well-established molecular clock coalescent methods. Our substitution rate estimates indicate that the ECSA-Br lineage is evolving at 7.15×10^{-4} substitutions per site per year (s/s/y; 95% Bayesian credible interval: $5.04 - 9.55 \times 10^{-4}$). This estimated rate is higher than that estimated for endemic lineages, and is similar to the evolutionary rates estimated for the epidemic lineage circulating in the Indian Ocean region (**Figure 2c**). A closer inspection of amino acid mutations indicate that the ECSA-Br strains lack both the A226V (E1 protein) and the L210Q (E2 protein) mutations that has been reported to increase virus transmissibility and persistence in *Ae. albopictus* populations in the Indian Ocean [47].

ML and Bayesian phylogenetic analyses reveal that the ECSA sequences from Brazil (hereafter named ECSA-Br lineage) form a single well-supported clade (bootstrap support = 100) (**Figure 3**). This is consistent with the establishment of the ECSA genotype in Brazil following the introduction of a single strain to the Americas [1]. The two isolates collected in late 2014 in Roraima cluster together and fall as expected within the diversity of other Asian genotype sequences from the Americas. Our phylogenetic reconstruction suggests at least five separate introductions of the Asian genotype strain Brazil (**Figure S1**), in contrast to a single introduction of the ECSA genotype followed by onward transmission. Moreover, all 13 ECSA isolates sampled in Roraima (*node C*) cluster together with maximum phylogenetic support (bootstrap support = 100; posterior probability = 1.00) (**Figure 3**). We consistently estimate the date of the most recent common ancestor of ECSA-Br Roraima clade to be mid-July 2016 (95% BCI: late March to late October 2016) (**Figure 3**); similar dating estimates under different coalescent models (**Figure S2**). In contrast to the Roraima strains, sequences from Manaus were found to be interspersed with isolates from Bahia and Pernambuco (**Figure 3**), indicating separate introductions of the CHIKV-ECSA lineage, some in early 2015 (*node B*), possibly from the northeast region of Brazil. Interestingly, according to travel history reports, the first autochthonous transmission of CHIKV in Manaus was linked to an index patient who reported spending holidays in Feira de Santana (Bahia state) in early 2015, during a period when this city was experiencing a large CHIKV outbreak [5]. The date of *node A* was estimated to be around mid-July 2014 (95% BCI: early Jul – late Aug 2014), shortly after the arrival of the presumed index case in Feira de Santana, Bahia [5]. This is in line with a single introduction to Bahia (*node A*), followed by

subsequent waves of transmission across the northeast and southeast regions of Brazil [5, 48, 49]. Our demographic reconstructions indicate that the outbreak in Roraima 2017 probably represents the third epidemic wave spreading across Brazil (**Figure S3**).

Next, we used notified case counts to estimate the basic reproductive number, R_0 , of the epidemic. R_0 is the average number of secondary cases caused by an infected individual and can be estimated from epidemic growth rates during its early exponential phase [43]. We find that $R_0 \approx 1.66$ (95% CI: 1.51 – 1.83), in line with previous reports from other settings [50-52]. A sensitivity analysis considering different exponential growth phase periods resulted in a lower bound for R_0 of around 1.23 (**Figure S4**). To gain insights into the possible magnitude of the outbreak and local surveillance capacity we used the equilibrium end state of a simple susceptible-infected-recovered (SIR) model: $N = S + I + R$, $S \sim 1/R_0$, $I \sim 0$, with N being the total population size of Roraima. Using this simple mathematical approach, we obtain an attack rate (R) of 0.39 (95% CI: 0.36 – 0.45), slightly lower than elsewhere in Brazil [12, 15]. This corresponds to an estimated 110,882 (95% CI: 102,352 – 127,940) infected individuals, and a case detection rate of 5.34% (95% CI: 4.63 – 5.79). This implies that approximately 1 case was notified for every 19 infections. If we assume 32.7 – 41.2% of the estimated infections are symptomatic, as previously reported in Bahia and Sergipe [53], then we estimate that the local observation success of symptomatic cases was between 12.8 – 16.1%. However, if we assume that 75 – 97% of people infected with CHIKV will develop symptomatic infections, as reported for the Indian Ocean lineage [10, 54, 55], then the chances of reported a symptomatic CHIKV case decrease to 5 – 7% [9]. Case reports suggest that the beginning of the exponential phase of the outbreak was in December 2016 (**Figure S4**), while genetic data suggests that the outbreak clade emerged around July 2016. However, between August 2014 and June 2016, 612 CHIKV notified cases and 40 confirmed cases were reported by the LACEN-RR. It is therefore likely that prior to Jan 2017, low but non-neglectable transmission of the Asian genotype occurred in Roraima.

We investigated the public's awareness of the chikungunya outbreak by retrospectively monitoring Google searches of the search term “chikungunya” in Roraima state from January 2014 to July 2018 (**Figure 4**). As a comparison, we

performed a similar search focusing on the neighbouring state of Amazonas. We found that web search activity and CHIKV cases counts in Roraima are highly correlated (notified cases: $r = 0.89$; confirmed cases: $r = 0.92$, **Figure 4d – e**). Additionally, the timing of the peak of Google searches corresponds to that of notified and confirmed cases with a peak in July 2017 (**Figure 4a and c, Figure 4b and f**). It is important to note that web search activity was available weeks or months before the final number of confirmed (and suspected) cases were made publicly available. This fact highlights the potential utility of monitoring disease-related searches during the outbreak. Interestingly, we find some web-search activity in Roraima before June 2016, particularly in September 2014, March 2015 and March 2016 (**Figure 4f**). These patterns are distinct to those in the Amazonas neighbouring state (notified cases: $r = 0.65$; confirmed cases: $r = 0.15$), which shows an early peak in November 2014, soon after the estimated age of *node B* (**Figure 3b**), followed by a peak in February 2016 and another in March 2017 (**Figure 4c**). These multiple peaks in internet search queries are consistent with the timing of at least 3 introductions detected in our phylogenetic analyses (**Figure 3b**), each possibly resulting in small epidemic waves of CHIKV in Manaus and Amazonas states.

Discussion

We describe a genomic epidemiological study which used genetic, epidemiological, and digital search data to investigate an outbreak caused by CHIKV in Boa Vista city, Roraima state, northern Brazil, in 2017. Using a combination of genetic, laboratory-confirmed and -suspected, and digital search data from 2014 to 2018, we find evidence for the replacement of the Asian lineage by the ECSA lineage in the north of Brazil. Moreover, we find that ECSA lineage was introduced in Roraima around July 2016, six months before the beginning of the exponential increase in case numbers. Using simple epidemiological models, we find that on average 1 in 17 (95% CI: 14 – 20) symptomatic CHIKV cases, a fraction of the 110,882 (95% CI: 102,352 – 127,940) estimated number of infections, sought medical care during the outbreak of CHIKV ECSA in Roraima. Finally, we find that Google search activity data shows a strong association with CHIKV notified cases in Roraima. Moreover, although the nanopore-based sequencing protocol for CHIKV utilized in this study has been described and validated previously [32], this study represent to our knowledge the first effort to generate on-site complete CHIKV genome sequences. Our results provide evidence of lineage replacement in Brazil, and to the best of our knowledge, deliver a description of the largest outbreak ever reported in north Brazil, revealing the circulation of the ECSA lineage in the Amazon region.

We estimate that 39% (95% CI: 36 – 45)% of Roraima's population was infected with CHIKV-ECSA-Br during the outbreak in 2017. Our estimates are higher than the 20% seropositive observed in a rural community in Bahia [10], and slightly lower than the 45.7 – 57.1% observed in two serosurveys conducted in the same state [12], where the ECSA lineage also seems to predominate. The observed differences in terms of the proportion of the population exposed to CHIKV in Roraima compared to previous estimates from the northeast region could result from partial protection resulting from low-level transmission of the CHIKV-Asian genotype during 2014 – 2016 in the north region. Alternatively, some level of cross-protection could have been conferred by previous exposure to Mayaro virus (MAYV); Mayaro is an antigenically-related alphavirus that may provide some level of cross-reactivity [56, 57] and is associated with *Haemagogus* spp. vectors [58], but has also been identified in *Culex quinquefasciatus* and *Aedes aegypti* mosquitoes [59]. MAYV has been detected in the north [60-64] and centre-west

[21, 59, 65-68] regions of Brazil. Moderate to high prevalence of MYV IgM have been found in urban northern areas [60], which could explain the limited spread of CHIKV in Manaus compared to Roraima.

Different CHIKV circulating lineages may have remarkably different public health consequences. Lineage-specific clinical presentations have been recently highlighted by a recent index cluster study which showed that 82% of CHIKV infections caused by the ECSA lineage are symptomatic, in comparison to only 52% of symptomatic infections caused by the Asian genotype [54]. While the Asian lineage seems to have circulated cryptically for 9 months before its first detection in the Caribbean [3], the faster detection of the ECSA lineage in Brazil could at least in part be a consequence of a higher rate of symptomatic to asymptomatic infections of the ECSA lineage circulating in Brazil. The time lag between the phylogenetic estimate of the date of introduction of a virus lineage and the date of the first confirmed case in a given region, enables us to identify surveillance gaps between the arrival and discovery of a virus in that region [69].

We used genomic data collected over a 3-year period to estimate the genetic history of the CHIKV-ECSA-Br lineage. We estimate that the CHIKV-ECSA-Br lineage arrived in Roraima around July 2016, whilst the first confirmed CHIKV cases in Roraima occurred earlier, in August 2014. That the discovery date anticipates the estimated date of introduction can be explained by initial introduction(s) of the Asian lineage (from the north of Brazil or from other south American regions) resulting in only limited onwards transmission, followed by the replacement of the Asian lineages by an epidemiological successful ECSA lineage. Transmission of the Asian genotype during this period is in line with an increase in notified and confirmed cases, as well internet search query data between August 2014 and June 2016. Nationwide molecular and seroprevalence studies combined with epidemiological modelling [70] will help to determine the proportion of cases caused by the ECSA compared to the Asian lineage in different geographic settings, and to identify which populations are still at risk of infection in Brazil.

We estimated high rates of nucleotide substitution for this lineage, which equates to around 8 (95% BCI: 6 – 11) nucleotide substitutions per year across the virus genome. Such rates are similar to the evolutionary rates estimated for the IOL lineage; these are typical of urban and epidemic transmission cycles in locations with an abundance of suitable hosts and lack of herd immunity [8]. None of the mutations

associated previously with increased transmissibility of the IOL lineage in *Ae. albopictus* mosquitos in the Indian Ocean region were identified in this study. However, it is currently unclear whether we should expect the same mutations to be linked with increased transmission in *Aedes* spp. populations both from Brazil and from Southeast Asia. Further, it is possible that CHIKV in Brazil is vectored mainly by the *Ae. aegypti* vector that is abundant throughout Brazil [71]. In line with this, CHIKV-ECSA was recently detected in *Aedes aegypti* from Maranhão [72] and Rio de Janeiro states [73].

The past dengue serotype 4 genotype II outbreak in Brazil ignited in the north of the country, and is inferred to have been introduced from Venezuela to Roraima, before spreading to the northeast and southeast region of Brazil [74]. Our genetic analysis reveals at least four instances of ECSA-Br virus lineage migration in the opposite direction, i.e., from northeastern to northern Brazil. Such a pattern may not be surprising due to the year-round persistence of *Aedes aegypti* mosquitos in the northeast and the north areas [31]. Within-country transmission will be dictated by human mobility, climatic synchrony, and levels of population immunity. Moreover, international spread of the ECSA-Br lineage is expected to regions linked to Brazil. Previous analyses of dengue virus serotypes has identified a strong connectivity between north Brazil and Venezuela [25, 75], and northeast Brazil and Haiti [31, 76]. In addition, Angola and Brazil are linked by human mobility and synchronous climates that have facilitated the migration of CHIKV-ECSA [1] and Zika virus (<http://virological.org/t/circulation-of-the-asian-lineage-zika-virus-in-angola/248>).

Improving surveillance in the Amazon region may help anticipate transmission of vector-borne diseases and also spillover from wild mammals of zoonotic viruses of particular concern [27]. Genomic portable sequencing of vector-borne viral infections in the Amazon may be particularly important in the context of early identification of circulation of strains newly (re)-introduced from wildlife. For example, yellow fever strains collected in Roraima seem to be at the source of the 2016-2018 yellow fever virus outbreak in southeast Brazil, which has affected large urban centres in Minas Gerais, São Paulo and Rio de Janeiro [26]. In the near future, the increasing rapidity and decreasing cost of genome sequencing in poorly sampled areas, combined with emerging theoretical approaches [77], will facilitate the investigation of possible associations between arbovirus lineage diversity, mosquito vectors, reservoir species, and transmission potential.

Finally, the reported synchronicities between notified chikungunya case counts in Roraima and the chikungunya-related Internet searches originated in the region highlight the potential complementarity that Internet search activity may offer in future disease outbreaks. Specifically, given that disease-related search activity can be monitored in near-real time, early signals of increases in disease activity may be spotted weeks or months before lab-confirmed case counts may be available in an unfolding outbreak.

Acknowledgments

We are thankful to all personnel from SVS/MS, PAHO/Brazil, Roraima and Boa Vista Health Surveillance System that coordinated surveillance and helped with data collection and assembly. We thank Oxford Nanopore Technologies for the support to the ZIBRA-2 (Zika in Brazil Real time Analyses-second round) project with additional flowcells and corresponding reagents, and also thank QIAGEN for donation of consumables.

Figure Legends

Figure 1. Context of this study. A. Map showing municipalities of Roraima state, including Boa Vista, bordering countries (Venezuela and French Guiana) and bordering Brazilian federal states (Amazonas and Pará). B. Map of Brazilian states, showing the states from which CHIKV sequence data in this study was analysed (Bahia, Alagoas, Pernambuco, Paraíba, Amazonas and Roraima). C. Barplot showing the annual number of notified CHIKV cases in selected states of Brazil (data obtained from the Brazilian Ministry of Health). Map was made with Natural Earth. Free vector and raster map data at naturalearthdata.com.

Fig. 2. Sequencing statistics, temporal signal and evolutionary rates of the CHIKV-ECSA lineage. A. Genome coverage plotted against RT-qPCR CT-values for the newly generated sequence data. B. Genetic divergence regressed against dates of sample collection for dataset 2 (CHIKV-ECSA-Br lineage). C. Evolutionary rate estimates for the CHIKV-ECSA-Br lineage obtained by this study (circle number 1) compared to published evolutionary rates obtained for other lineages. Circles numbered 2 to 8 represent point estimates reported in [1, 8, 78]. Horizontal bars represent 95% highest posterior density credible intervals for associated evolutionary rates.

Figure 3. Genetic analysis of the CHIKV-ECSA genotype. A. Maximum likelihood phylogeny depicting the monophyletic clade containing all the Brazilian ECSA isolates (ECSA-Br lineage). B. Time-calibrated phylogeny of all available CHIKV-ECSA whole genome sequences from Brazil, including 18 novel genomes from Roraima and Amazonas states. Colours correspond to state of sample collection. Violin plots show 95% Bayesian credible intervals for associated node heights [38].

Figure 4. Digital surveillance of chikungunya disease in northern Brazil. A and B show respectively the number of notified CHIKV cases in LACEN-RR and LACEN-AM between Jan 2014 and Sep 2018. Panels C and F show Google Trends activity for the term “chikungunya” in Amazonas (C) and Roraima (F) from Jan 2016 and Sep 2018. Panels D and G show the correlation between Google Trends activity and confirmed cases in Amazonas (D) and Roraima (G), while panels E and H show the

correlation between Google Trends activity and notified cases in Amazonas (E) and Roraima (H).

Table 1. Epidemiological data for virus isolates from Roraima (RR) and Amazonas (AM). CT=cycle threshold, *d*=days from onset of symptoms to sample collection. Corresponding sequencing statistics are available in **Table S1**. Isolates were collected around 2.3 (range: 0 – 5) days after onset of symptoms. Acc. Number = GenBank accession number.

Isolate	State, Municipality	Acc. Number	Ct RT- qPCR	Coverage (%)	Age, Sex	Collection date	<i>d</i>
AMA290	AM, Manaus	MK121891	NA	90.2	76, F	15/07/2015	5
AMA291	AM, Manaus	MK121892	NA	80.7	48, F	15/07/2015	4
AMA292	AM, Manaus	MK121893	NA	90.2	50, M	15/07/2015	0
AMA293	AM, Manaus	MK121894	NA	84.4	42, M	31/01/2016	4
AMA294	RR, Boa Vista	MK134712	NA	90.2	45, F	01/12/2014	2
AMA295	RR, Unknown	MK134713	NA	90.2	9, F	11/11/2014	1
AMA74	AM, Manaus	MK121895	15	90.2	32, F	20/03/2017	2
AMA346	RR, Boa Vista	MK121896	13.7	90.2	30, F	03/03/2017	1
AMA350	RR, Bonfim	MK121897	27.15	54.7	32, F	20/02/2017	1
AMA352	RR, Boa Vista	MK121898	17.33	88.6	3, F	22/02/2017	1
AMA354	RR, Boa Vista	MK121899	23.36	86.9	19, F	17/03/2017	1
AMA362	RR, Iracema	MK121900	18.63	88.6	31, F	17/03/2017	1
AMA364	RR, Boa Vista	MK121901	25.93	83.3	19, F	17/03/2017	2
AMA366	RR, Boa Vista	MK121902	19.87	90.0	36, F	17/03/2017	2
AMA368	RR, Boa Vista	MK121903	25.91	93.1	26, F	15/03/2017	2
AMA369	RR, Boa Vista	MK121904	21.55	95.6	52, M	02/03/2017	3
AMA374	RR, Boa Vista	MK121905	27.41	71.4	64, F	02/03/2017	4
AMA379	RR, Boa Vista	MK121906	17.5	96.1	38, F	27/02/2017	4
AMA381	RR, Boa Vista	MK121907	16.66	97.7	31, F	27/02/2017	4
AMA382	RR, Boa Vista	MK121908	14.58	76.6	30, F	05/03/2017	1

Supplementary Figure Legends

Figure S1. Maximum likelihood phylogenetic tree of the CHIKV Asian genotype.

Includes isolates from Southeast Asia, Americas and Brazil. Isolates represented by blue tips were sampled in Roraima, while isolates shown in red represent other strains sampled in Brazil.

Figure S2. Dating estimates obtained under different coalescent models.

Estimates for node A (time of the most recent common ancestor, in dark red, see Figure 3b), node B (main Amazonas clade, in green), and node C (Roraima clade, in purple) are shown for different non-parametric models (Bayesian skygrid, skyride, skyline) and for a simple constant population size model.

Figure S3. Demographic dynamics of CHIKV ECSA-Brazilian lineage in Brazil.

Fluctuation of effective population size over time as inferred through a Bayesian skygrid coalescent model.

Figure S4. Exponential Period of the CHIKV epidemic in Boa Vista

municipality, Roraima state. Log number of notified cases per month are plotted against number of months since January 2015.

Supplementary Table

Table S1. Minion sequencing statistics

Isolate	Mapped reads	Average depth coverage	Bases covered >10x	Bases covered > 25x	Reference covered (%)
AMA290	13624	767	10276	10258	90.2
AMA291	60261	2047	9489	9280	80.7
AMA292	68090	2746	10402	10223	90.2
AMA293	64953	2096	9745	9701	84.4
AMA294	21361	701	10252	10022	90.2
AMA295	16370	531	10188	10077	90.2
AMA74	42276	1951	10396	10195	90.2
AMA346	31210	1225	10243	10208	90.2
AMA350	63672	1673	7522	7168	54.7
AMA352	13530	536	10219	10184	88.6
AMA354	22214	752	10082	9985	86.9
AMA362	9938	398	10237	10128	88.6
AMA364	28494	1079	9813	9577	83.3
AMA366	38228	1441	10264	10224	90.0
AMA368	12968	503	11122	10825	93.1
AMA369	7280	311	11225	11149	95.6
AMA374	7030	305	10225	8805	71.4
AMA379	7970	348	11226	11092	96.1
AMA381	7522	327	11214	11208	97.7
AMA382	14040	411	9915	9424	76.6

References

1. Nunes MR, Faria NR, de Vasconcelos JM, Golding N, Kraemer MU, de Oliveira LF, et al. Emergence and potential for spread of Chikungunya virus in Brazil. *BMC medicine*. 2015;13:102. Epub 2015/05/16. doi: 10.1186/s12916-015-0348-x. PubMed PMID: 25976325; PubMed Central PMCID: PMC4433093.
2. Organization PAWH. Number of Reported Cases of Chikungunya Fever in the Americas, by Country or Territory. Washington: 2017.
3. Sahadeo NSD, Allicock OM, De Salazar PM, Auguste AJ, Widen S, Olowokure B, et al. Understanding the evolution and spread of chikungunya virus in the Americas using complete genome sequences. *Virus Evol*. 2017;3(1):vex010. Epub 2017/05/10. doi: 10.1093/ve/vex010. PubMed PMID: 28480053; PubMed Central PMCID: PMCPMC5413804.
4. Nsoesie EO, Kraemer MU, Golding N, Pigott DM, Brady OJ, Moyes CL, et al. Global distribution and environmental suitability for chikungunya virus, 1952 to 2015. *Euro surveillance : bulletin Europeen sur les maladies transmissibles = European communicable disease bulletin*. 2016;21(20). Epub 2016/05/31. doi: 10.2807/1560-7917.ES.2016.21.20.30234. PubMed PMID: 27239817; PubMed Central PMCID: PMCPMC4902126.
5. Faria NR, Lourenco, J., Cerqueira, E. M., Lima, M. M., Pybus, O. G., Alcantara, L. C. J. Epidemiology of Chikungunya Virus in Bahia, Brazil, 2014-2015. *PLoS Currents Outbreaks*. 2016;1. doi: 10.1371/currents.outbreaks.c97507e3e48efb946401755d468c28b2.
6. Johansson MA. Chikungunya on the move. *Trends in parasitology*. 2015;31(2):43-5. Epub 2015/02/05. doi: 10.1016/j.pt.2014.12.008. PubMed PMID: 25649340; PubMed Central PMCID: PMCPMC4583061.
7. Kraemer MUG, Sinka, M. E., Duda, K. A., Mylne, A., Shearer, F. M., Barker, C. M., Moore, C. G., Carvalho, R. G., Coelho, G. E., Bortel, W. V., Hendrickx, G., Schaffner, F., Elyazar, I. R. F., Teng, H. J., Brady, O. J., Messina, J. P., Pigott, D., M., Scott, T. W., Smith, D. L., Wint, G. R. W., Golding, N., Hay, S. I. The global distribution of the arbovirus vectors *Aedes aegypti* and *Ae. albopictus*. *eLife*. 2015;10.7554(eLife.08347).
8. Volk SM, Chen R, Tsetsarkin KA, Adams AP, Garcia TI, Sall AA, et al. Genome-scale phylogenetic analyses of chikungunya virus reveal independent

- 709 emergences of recent epidemics and various evolutionary rates. Journal of
710 virology. 2010;84(13):6497-504. Epub 2010/04/23. doi: 10.1128/JVI.01603-09.
711 PubMed PMID: 20410280; PubMed Central PMCID: PMC2903258.
- 712 9. Staples JE, Breiman RF, Powers AM. Chikungunya fever: an
713 epidemiological review of a re-emerging infectious disease. Clinical infectious
714 diseases : an official publication of the Infectious Diseases Society of America.
715 2009;49(6):942-8. Epub 2009/08/12. doi: 10.1086/605496. PubMed PMID:
716 19663604.
- 717 10. Cunha RV, Trinta KS, Montalbano CA, Sucupira MV, de Lima MM, Marques
718 E, et al. Seroprevalence of Chikungunya Virus in a Rural Community in Brazil.
719 PLoS Negl Trop Dis. 2017;11(1):e0005319. Epub 2017/01/21. doi:
720 10.1371/journal.pntd.0005319. PubMed PMID: 28107342; PubMed Central
721 PMCID: PMCPMC5287455.
- 722 11. Figueiredo LTM. Large outbreaks of Chikungunya virus in Brazil reveal
723 uncommon clinical features and fatalities. Revista da Sociedade Brasileira de
724 Medicina Tropical. 2017;50(5):583-4. Epub 2017/11/22. doi: 10.1590/0037-
725 8682-0397-2017. PubMed PMID: 29160502.
- 726 12. Dias JP, Costa M, Campos GS, Paixao ES, Natividade MS, Barreto FR, et al.
727 Seroprevalence of Chikungunya Virus after Its Emergence in Brazil. Emerging
728 infectious diseases. 2018;24(4):617-24. Epub 2018/03/20. doi:
729 10.3201/eid2404.171370. PubMed PMID: 29553317; PubMed Central PMCID:
730 PMCPMC5875253.
- 731 13. Brito CAA, Teixeira MG. Increased number of deaths during a
732 chikungunya epidemic in Pernambuco, Brazil. Memorias do Instituto Oswaldo
733 Cruz. 2017;112(9):650-1. Epub 2017/09/14. doi: 10.1590/0074-02760170124.
734 PubMed PMID: 28902292; PubMed Central PMCID: PMCPMC5572452.
- 735 14. Freitas ARR, Cavalcanti L, Von Zuben AP, Donalisio MR. Excess Mortality
736 Related to Chikungunya Epidemics in the Context of Co-circulation of Other
737 Arboviruses in Brazil. PLoS currents. 2017;9. Epub 2017/12/22. doi:
738 10.1371/currents.outbreaks.14608e586cd321d8d5088652d7a0d884. PubMed
739 PMID: 29263941; PubMed Central PMCID: PMCPMC5731794.
- 740 15. Cunha MDP, Santos CAD, Neto DFL, Schanoski AS, Pour SZ, Passos SD, et
741 al. Outbreak of chikungunya virus in a vulnerable population of Sergipe, Brazil-A

- 742 molecular and serological survey. Journal of clinical virology : the official
743 publication of the Pan American Society for Clinical Virology. 2017;97:44-9. Epub
744 2017/11/04. doi: 10.1016/j.jcv.2017.10.015. PubMed PMID: 29100064.
- 745 16. Silva MM, Rodrigues MS, Paploski IA, Kikuti M, Kasper AM, Cruz JS, et al.
746 Accuracy of Dengue Reporting by National Surveillance System, Brazil. Emerging
747 infectious diseases. 2016;22(2):336-9. Epub 2016/01/27. doi:
748 10.3201/eid2202.150495. PubMed PMID: 26812472; PubMed Central PMCID:
749 PMCPMC4734515.
- 750 17. Pessoa R, Patriota JV, Lourdes de Souza M, Felix AC, Mamede N, Sanabani
751 SS. Investigation Into an Outbreak of Dengue-like Illness in Pernambuco, Brazil,
752 Revealed a Cocirculation of Zika, Chikungunya, and Dengue Virus Type 1.
753 Medicine. 2016;95(12):e3201. Epub 2016/03/26. doi:
754 10.1097/MD.00000000000003201. PubMed PMID: 27015222; PubMed Central
755 PMCID: PMCPMC4998417.
- 756 18. Sardi SI, Somasekar S, Naccache SN, Bandeira AC, Tauro LB, Campos GS, et
757 al. Coinfections of Zika and Chikungunya Viruses in Bahia, Brazil, Identified by
758 Metagenomic Next-Generation Sequencing. Journal of clinical microbiology.
759 2016;54(9):2348-53. doi: 10.1128/JCM.00877-16. PubMed PMID: 27413190;
760 PubMed Central PMCID: PMCPMC5005514.
- 761 19. Cardoso CW, Paploski IA, Kikuti M, Rodrigues MS, Silva MM, Campos GS, et
762 al. Outbreak of Exanthematous Illness Associated with Zika, Chikungunya, and
763 Dengue Viruses, Salvador, Brazil. Emerging infectious diseases.
764 2015;21(12):2274-6. Epub 2015/11/20. doi: 10.3201/eid2112.151167. PubMed
765 PMID: 26584464; PubMed Central PMCID: PMCPMC4672408.
- 766 20. Mavian C, Rife BD, Dollar JJ, Cella E, Ciccozzi M, Prosperi MCF, et al.
767 Emergence of recombinant Mayaro virus strains from the Amazon basin.
768 Scientific reports. 2017;7(1):8718. Epub 2017/08/20. doi: 10.1038/s41598-
769 017-07152-5. PubMed PMID: 28821712; PubMed Central PMCID:
770 PMCPMC5562835.
- 771 21. Vieira CJ, Silva DJ, Barreto ES, Siqueira CE, Colombo TE, Ozanic K, et al.
772 Detection of Mayaro virus infections during a dengue outbreak in Mato Grosso,
773 Brazil. Acta Trop. 2015;147:12-6. Epub 2015/03/31. doi:
774 10.1016/j.actatropica.2015.03.020. PubMed PMID: 25817238.

- 775 22. Del Valle SY, McMahon BH, Asher J, Hatchett R, Lega JC, Brown HE, et al.
776 Summary results of the 2014-2015 DARPA Chikungunya challenge. BMC
777 infectious diseases. 2018;18(1):245. Epub 2018/05/31. doi: 10.1186/s12879-
778 018-3124-7. PubMed PMID: 29843621; PubMed Central PMCID:
779 PMCPMC5975673.
- 780 23. Silva NMD, Teixeira RAG, Cardoso CG, Siqueira Junior JB, Coelho GE,
781 Oliveira ESF. Chikungunya surveillance in Brazil: challenges in the context of
782 Public Health. Epidemiol Serv Saude. 2018;27(3):e2017127. Epub 2018/09/06.
783 doi: 10.5123/S1679-49742018000300003. PubMed PMID: 30183867.
- 784 24. Bogoch, II, Brady OJ, Kraemer MUG, German M, Creatore MI, Kulkarni MA,
785 et al. Anticipating the international spread of Zika virus from Brazil. Lancet.
786 2016;387(10016):335-6. Epub 2016/01/19. doi: 10.1016/S0140-
787 6736(16)00080-5. PubMed PMID: 26777915; PubMed Central PMCID:
788 PMCPMC4873159.
- 789 25. Nunes MR, Faria NR, Vasconcelos HB, Medeiros DB, Silva de Lima CP,
790 Carvalho VL, et al. Phylogeography of dengue virus serotype 4, Brazil, 2010-
791 2011. Emerging infectious diseases. 2012;18(11):1858-64. doi:
792 10.3201/eid1811.120217. PubMed PMID: 23092706; PubMed Central PMCID:
793 PMC3559147.
- 794 26. Faria NR, Kraemer MUG, Hill SC, Goes de Jesus J, Aguiar RS, Iani FCM, et al.
795 Genomic and epidemiological monitoring of yellow fever virus transmission
796 potential. Science. 2018;361(6405):894-9. Epub 2018/08/25. doi:
797 10.1126/science.aat7115. PubMed PMID: 30139911.
- 798 27. Olival KJ, Hosseini PR, Zambrana-Torrel C, Ross N, Bogich TL, Daszak P.
799 Host and viral traits predict zoonotic spillover from mammals. Nature.
800 2017;546(7660):646-50. Epub 2017/06/22. doi: 10.1038/nature22975.
801 PubMed PMID: 28636590; PubMed Central PMCID: PMCPMC5570460.
- 802 28. Weiss DJ, Nelson A, Gibson HS, Temperley W, Peedell S, Lieber A, et al. A
803 global map of travel time to cities to assess inequalities in accessibility in 2015.
804 Nature. 2018;553(7688):333-6. Epub 2018/01/11. doi: 10.1038/nature25181.
805 PubMed PMID: 29320477.
- 806 29. Naveca FG, Nascimento VAD, Souza VC, Nunes BTD, Rodrigues DSG,
807 Vasconcelos P. Multiplexed reverse transcription real-time polymerase chain

808 reaction for simultaneous detection of Mayaro, Oropouche, and Oropouche-like
809 viruses. *Memorias do Instituto Oswaldo Cruz*. 2017;112(7):510-3. Epub
810 2017/06/08. doi: 10.1590/0074-02760160062. PubMed PMID: 28591313;
811 PubMed Central PMCID: PMC5452489.

812 30. Lanciotti RS, Kosoy OL, Laven JJ, Panella AJ, Velez JO, Lambert AJ, et al.
813 Chikungunya virus in US travelers returning from India, 2006. *Emerging*
814 *infectious diseases*. 2007;13(5):764-7. doi: 10.3201/eid1305.070015. PubMed
815 PMID: 17553261; PubMed Central PMCID: PMC5452489.

816 31. Faria NR, Quick J, Claro IM, Theze J, de Jesus JG, Giovanetti M, et al.
817 Establishment and cryptic transmission of Zika virus in Brazil and the Americas.
818 *Nature*. 2017;546(7658):406-10. Epub 2017/05/26. doi: 10.1038/nature22401.
819 PubMed PMID: 28538727; PubMed Central PMCID: PMC5722632.

820 32. Quick J. G, N. D., Pullan, S. T., Claro, I. M., Smith, A. D., Gangavarapu, k.,
821 Oliveira, G., Robles-Sikisaka, R., Rogers, T. F., Beutler, N. A., Burton, D. R., Lewis-
822 Ximenez, L. L., de Jesus, J. G., Giovanetti, M., Hill, S., Black, A., Bedford, T., Carroll,
823 M. W., Nunes, M., Alcantara, L. C., Sabino, E. C., Baylis, S. A., Faria, N. R., Loose, M.,
824 Simpson, J. T., Pybus, O. G., Andersen, K. G., Loman, N. J. Multiplex PCR method for
825 MinION and Illumina sequencing of Zika and other virus genomes directly from
826 clinical samples. *Nature Protocols*. 2017;XX(X):XX.

827 33. Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. GenBank.
828 *Nucleic acids research*. 2016;44(D1):D67-72. doi: 10.1093/nar/gkv1276.
829 PubMed PMID: 26590407; PubMed Central PMCID: PMC4702903.

830 34. da Costa AC, Thézé, J., Komninakis, S. C. V., Sanz-Duro, R. L., Felinto, M. R.
831 L., Moura, L. C. C., Barroso, I. M. O., Santos, L. E. C., Nunes, M. A., L., Moura, A. A.,
832 Lourenco, J., Deng, X., Delwart, E. L., Guimaraes, M. R. A. S., Pybys, O. G., Sabino, E.
833 C., Faria, N. R. Spread of Chikungunya Virus East/Central/South African Genotype
834 in Northeast Brazil. *Emerging infectious diseases*. 2017;23(10).

835 35. Bruen TC, Philippe H, Bryant D. A simple and robust statistical test for
836 detecting the presence of recombination. *Genetics*. 2006;172(4):2665-81. doi:
837 10.1534/genetics.105.048975. PubMed PMID: 16489234; PubMed Central
838 PMCID: PMC1456386.

839 36. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-
840 analysis of large phylogenies. *Bioinformatics*. 2014;30(9):1312-3. doi:

841 10.1093/bioinformatics/btu033. PubMed PMID: 24451623; PubMed Central
842 PMCID: PMCPMC3998144.

843 37. Rambaut A, Lam, T. T., Fagundes de Carvalho, L., Pybus, O. G. Exploring the
844 temporal structure of heterochronous sequences using TempEst (formerly Path-
845 O-Gen). *Virus Evolution*. 2016;2(vew007).

846 38. Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A.
847 Bayesian phylogenetic and phylodynamic data integration using BEAST 1.10.
848 *Virus Evol.* 2018;4(1):vey016. Epub 2018/06/27. doi: 10.1093/ve/vey016.
849 PubMed PMID: 29942656; PubMed Central PMCID: PMCPMC6007674.

850 39. Drummond AJ, Rambaut A, Shapiro B, Pybus OG. Bayesian coalescent
851 inference of past population dynamics from molecular sequences. *Molecular*
852 *biology and evolution*. 2005;22(5):1185-92. doi: 10.1093/molbev/msi103.
853 PubMed PMID: 15703244.

854 40. Minin VN, Bloomquist EW, Suchard MA. Smooth skyride through a rough
855 skyline: Bayesian coalescent-based inference of population dynamics. *Molecular*
856 *biology and evolution*. 2008;25(7):1459-71. Epub 2008/04/15. doi:
857 10.1093/molbev/msn090. PubMed PMID: 18408232; PubMed Central PMCID:
858 PMC3302198.

859 41. Gill MS, Lemey P, Faria NR, Rambaut A, Shapiro B, Suchard MA. Improving
860 Bayesian population dynamics inference: a coalescent-based model for multiple
861 loci. *Molecular biology and evolution*. 2013;30(3):713-24. doi:
862 10.1093/molbev/mss265. PubMed PMID: 23180580; PubMed Central PMCID:
863 PMC3563973.

864 42. Drummond AJ, Ho SY, Phillips MJ, Rambaut A. Relaxed phylogenetics and
865 dating with confidence. *PLoS biology*. 2006;4(5):e88. doi:
866 10.1371/journal.pbio.0040088. PubMed PMID: 16683862; PubMed Central
867 PMCID: PMC1395354.

868 43. Lourenço J, Monteiro, M. L., Valdez, T., Rodrigues, J. M., Pybus, O. G., Faria,
869 N. R. Epidemiology of the Zika virus outbreak in the Cabo Verde Islands, West
870 Africa. *PLoS currents*. 2018. doi:
871 10.1371/currents.outbreaks.19433b1e4d007451c691f138e1e67e8c.

872 44. Salje H, Lessler J, Paul KK, Azman AS, Rahman MW, Rahman M, et al. How
873 social structures, space, and behaviors shape the spread of infectious diseases

874 using chikungunya as a case study. Proceedings of the National Academy of
875 Sciences of the United States of America. 2016;113(47):13420-5. Epub
876 2016/11/09. doi: 10.1073/pnas.1611391113. PubMed PMID: 27821727;
877 PubMed Central PMCID: PMCPMC5127331.

878 45. Chan EH, Sahai V, Conrad C, Brownstein JS. Using web search query data
879 to monitor dengue epidemics: a new model for neglected tropical disease
880 surveillance. PLoS Negl Trop Dis. 2011;5(5):e1206. Epub 2011/06/08. doi:
881 10.1371/journal.pntd.0001206. PubMed PMID: 21647308; PubMed Central
882 PMCID: PMCPMC3104029.

883 46. Team RC. R: A Language and Environment for Computing. Vienna, Austria:
884 R Foundation for Statistical Computing; 2014.

885 47. Tsetsarkin KA, McGee CE, Volk SM, Vanlandingham DL, Weaver SC, Higgs
886 S. Epistatic roles of E2 glycoprotein mutations in adaption of chikungunya virus
887 to Aedes albopictus and Ae. aegypti mosquitoes. PloS one. 2009;4(8):e6835.
888 Epub 2009/09/01. doi: 10.1371/journal.pone.0006835. PubMed PMID:
889 19718263; PubMed Central PMCID: PMCPMC2729410.

890 48. Tanabe ELL, Tanabe ISB, Santos ECD, Marques J, Borges AA, Lima MC, et
891 al. Report of East-Central South African Chikungunya virus genotype during the
892 2016 outbreak in the Alagoas State, Brazil. Revista do Instituto de Medicina
893 Tropical de Sao Paulo. 2018;60:e19. Epub 2018/04/26. doi: 10.1590/s1678-
894 9946201860019. PubMed PMID: 29694603; PubMed Central PMCID:
895 PMCPMC5956549.

896 49. Charlys da Costa A, Theze J, Komninakis SCV, Sanz-Duro RL, Felinto MRL,
897 Moura LCC, et al. Spread of Chikungunya Virus East/Central/South African
898 Genotype in Northeast Brazil. Emerging infectious diseases. 2017;23(10):1742-4.
899 Epub 2017/09/21. doi: 10.3201/eid2310.170307. PubMed PMID: 28930031;
900 PubMed Central PMCID: PMCPMC5621546.

901 50. Robinson M, Conan A, Duong V, Ly S, Ngan C, Buchy P, et al. A model for a
902 chikungunya outbreak in a rural Cambodian setting: implications for disease
903 control in uninfected areas. PLoS Negl Trop Dis. 2014;8(9):e3120. Epub
904 2014/09/12. doi: 10.1371/journal.pntd.0003120. PubMed PMID: 25210729;
905 PubMed Central PMCID: PMC4161325.

- 906 51. Perkins TA, Metcalf CJ, Grenfell BT, Tatem AJ. Estimating drivers of
907 autochthonous transmission of chikungunya virus in its invasion of the americas.
908 PLoS currents. 2015;7. Epub 2015/03/05. doi:
909 10.1371/currents.outbreaks.a4c7b6ac10e0420b1788c9767946d1fc. PubMed
910 PMID: 25737803; PubMed Central PMCID: PMC4339250.
- 911 52. Salje H, Cauchemez S, Alera MT, Rodriguez-Barraquer I, Thaisomboonsuk
912 B, Srikiatkachorn A, et al. Reconstruction of 60 Years of Chikungunya
913 Epidemiology in the Philippines Demonstrates Episodic and Focal Transmission.
914 The Journal of infectious diseases. 2016;213(4):604-10. Epub 2015/09/28. doi:
915 10.1093/infdis/jiv470. PubMed PMID: 26410592; PubMed Central PMCID:
916 PMCPMC4721913.
- 917 53. Dias JP, Costa, M. C. N., Campos, G. S., Paixao, E. S., Natividade, M. S.,
918 Barreto, F. R., Itaparica, M. S. C., Goes, C., Oliveira, F. L. S., Santana, E. B., Silva, N. S.
919 J., Brito, C. A. A., Rodrigues, L. C., Sardi, S. I., Saavedra, R. C., Teixeira, M. G.
920 Seroprevalence of Chikungunya Virus after Its Emergence in Brazil. Emerging
921 infectious diseases. 2018;24(4):617-24.
- 922 54. Bustos Carrillo F, Collado D, Sanchez N, Ojeda S, Lopez Mercado B, Burger-
923 Calderon R, et al. Epidemiological Evidence for Lineage-specific Differences in
924 the Risk of Inapparent Chikungunya Virus Infection. Journal of virology. 2018.
925 Epub 2018/11/23. doi: 10.1128/JVI.01622-18. PubMed PMID: 30463967.
- 926 55. Sissoko D, Moendandze A, Malvy D, Giry C, Ezzedine K, Solet JL, et al.
927 Seroprevalence and risk factors of chikungunya virus infection in Mayotte, Indian
928 Ocean, 2005-2006: a population-based survey. PloS one. 2008;3(8):e3066. Epub
929 2008/08/30. doi: 10.1371/journal.pone.0003066. PubMed PMID: 18725980;
930 PubMed Central PMCID: PMCPMC2518850.
- 931 56. Hassing RJ, Leparco-Goffart I, Tolou H, van Doornum G, van Genderen PJ.
932 Cross-reactivity of antibodies to viruses belonging to the Semliki forest
933 serocomplex. Euro surveillance : bulletin Europeen sur les maladies
934 transmissibles = European communicable disease bulletin. 2010;15(23). Epub
935 2010/06/16. PubMed PMID: 20546691.
- 936 57. Powers AM, Brault AC, Shirako Y, Strauss EG, Kang W, Strauss JH, et al.
937 Evolutionary relationships and systematics of the alphaviruses. Journal of
938 virology. 2001;75(21):10118-31. Epub 2001/10/03. doi:

939 10.1128/JVI.75.21.10118-10131.2001. PubMed PMID: 11581380; PubMed
940 Central PMCID: PMC114586.

941 58. Vasconcelos PFC, Travassos da Rosa, A. P. A., Degallier, N., Travassos da
942 Rosa, J. F. S., Pinheiro, F. R. Clinical and ecoepidemiological situation of human
943 arboviruses in Brazilian Amazonia. Journal of the Brazilian Association for the
944 Advancement of Science. 1992;43(2):117-24.

945 59. Serra OP, Cardoso BF, Ribeiro AL, Santos FA, Shlessarenko RD. Mayaro
946 virus and dengue virus 1 and 4 natural infection in culicids from Cuiaba, state of
947 Mato Grosso, Brazil. Memorias do Instituto Oswaldo Cruz. 2016;111(1):20-9.
948 Epub 2016/01/20. doi: 10.1590/0074-02760150270. PubMed PMID: 26784852;
949 PubMed Central PMCID: PMCPMC4727432.

950 60. Mourao MP, Bastos Mde S, de Figueiredo RP, Gimaque JB, Galusso Edos S,
951 Kramer VM, et al. Mayaro fever in the city of Manaus, Brazil, 2007-2008. Vector
952 borne and zoonotic diseases. 2012;12(1):42-6. Epub 2011/09/20. doi:
953 10.1089/vbz.2011.0669. PubMed PMID: 21923266; PubMed Central PMCID:
954 PMCPMC3249893.

955 61. Receveur MC, Grandadam M, Pistone T, Malvy D. Infection with Mayaro
956 virus in a French traveller returning from the Amazon region, Brazil, January,
957 2010. Euro surveillance : bulletin Europeen sur les maladies transmissibles =
958 European communicable disease bulletin. 2010;15(18). Epub 2010/05/13.
959 PubMed PMID: 20460093.

960 62. Azevedo RS, Silva EV, Carvalho VL, Rodrigues SG, Nunes-Neto JP, Monteiro
961 H, et al. Mayaro fever virus, Brazilian Amazon. Emerging infectious diseases.
962 2009;15(11):1830-2. Epub 2009/11/07. doi: 10.3201/eid1511.090461. PubMed
963 PMID: 19891877; PubMed Central PMCID: PMCPMC2857233.

964 63. LeDuc JW, Pinheiro FP, Travassos da Rosa AP. An outbreak of Mayaro
965 virus disease in Belterra, Brazil. II. Epidemiology. The American journal of
966 tropical medicine and hygiene. 1981;30(3):682-8. Epub 1981/05/01. PubMed
967 PMID: 6266264.

968 64. Causey OR, Maroja OM. Mayaro virus: a new human disease agent. III.
969 Investigation of an epidemic of acute febrile illness on the river Guama in Para,
970 Brazil, and isolation of Mayaro virus as causative agent. The American journal of

971 tropical medicine and hygiene. 1957;6(6):1017-23. Epub 1957/11/01. PubMed
972 PMID: 13487974.

973 65. da Costa VG, de Rezende Feres VC, Saivish MV, de Lima Gimaque JB,
974 Moreli ML. Silent emergence of Mayaro and Oropouche viruses in humans in
975 Central Brazil. International journal of infectious diseases : IJID : official
976 publication of the International Society for Infectious Diseases. 2017;62:84-5.
977 Epub 2017/07/26. doi: 10.1016/j.ijid.2017.07.016. PubMed PMID: 28739422.

978 66. Brunini S, Franca DDS, Silva JB, Silva LN, Silva FPA, Spadoni M, et al. High
979 Frequency of Mayaro Virus IgM among Febrile Patients, Central Brazil. Emerging
980 infectious diseases. 2017;23(6):1025-6. Epub 2017/05/19. doi:
981 10.3201/eid2306.160929. PubMed PMID: 28518022; PubMed Central PMCID:
982 PMC5443426.

983 67. Pauvolid-Correa A, Juliano RS, Campos Z, Velez J, Nogueira RM, Komar N.
984 Neutralising antibodies for Mayaro virus in Pantanal, Brazil. Memórias do
985 Instituto Oswaldo Cruz. 2015;110(1):125-33. Epub 2015/03/06. doi:
986 10.1590/0074-02760140383. PubMed PMID: 25742272; PubMed Central
987 PMCID: PMC4371226.

988 68. Zuchi N, Heinen LB, Santos MA, Pereira FC, Shlessarenko RD. Molecular
989 detection of Mayaro virus during a dengue outbreak in the state of Mato Grosso,
990 Central-West Brazil. Memórias do Instituto Oswaldo Cruz. 2014;109(6):820-3.
991 Epub 2014/08/21. PubMed PMID: 25141284; PubMed Central PMCID:
992 PMC4238776.

993 69. Grubaugh ND, Faria NR, Andersen KG, Pybus OG. Genomic Insights into
994 Zika Virus Emergence and Spread. Cell. 2018;172(6):1160-2. Epub 2018/03/10.
995 doi: 10.1016/j.cell.2018.02.027. PubMed PMID: 29522736.

996 70. Lourenco J, Maia de Lima M, Faria NR, Walker A, Kraemer MU, Villabona-
997 Arenas CJ, et al. Epidemiological and ecological determinants of Zika virus
998 transmission in an urban setting. eLife. 2017;6. Epub 2017/09/10. doi:
999 10.7554/eLife.29820. PubMed PMID: 28887877; PubMed Central PMCID:
1000 PMC5638629.

1001 71. Kraemer MU, Sinka ME, Duda KA, Mylne AQ, Shearer FM, Barker CM, et al.
1002 The global distribution of the arbovirus vectors *Aedes aegypti* and *Ae. albopictus*.

- 1003 eLife. 2015;4:e08347. Epub 2015/07/01. doi: 10.7554/eLife.08347. PubMed
- 1004 PMID: 26126267; PubMed Central PMCID: PMC4493616.
- 1005 72. Aragao CF, Cruz ACR, Nunes Neto JP, Monteiro HAO, da Silva EVP, da Silva
- 1006 SP, et al. Circulation of Chikungunya virus in Aedes aegypti in Maranhao,
- 1007 Northeast Brazil. Acta Trop. 2018;186:1-4. Epub 2018/06/23. doi:
- 1008 10.1016/j.actatropica.2018.06.022. PubMed PMID: 29932932.
- 1009 73. Souza TM, Azeredo EL, Badolato-Correa J, Damasco PV, Santos C,
- 1010 Petitinga-Paiva F, et al. First Report of the East-Central South African Genotype of
- 1011 Chikungunya Virus in Rio de Janeiro, Brazil. PLoS currents. 2017;9. Epub
- 1012 2017/03/14. doi:
- 1013 10.1371/currents.outbreaks.4200119978d62ccaa454599cd2735727. PubMed
- 1014 PMID: 28286701; PubMed Central PMCID: PMC5325710.
- 1015 74. Faria NR, da Costa AC, Lourenco J, Loureiro P, Lopes ME, Ribeiro R, et al.
- 1016 Genomic and epidemiological characterisation of a dengue virus outbreak among
- 1017 blood donors in Brazil. Scientific reports. 2017;7(1):15216. Epub 2017/11/11.
- 1018 doi: 10.1038/s41598-017-15152-8. PubMed PMID: 29123142; PubMed Central
- 1019 PMCID: PMC5680240.
- 1020 75. Nunes MR, Palacios G, Faria NR, Sousa EC, Jr., Pantoja JA, Rodrigues SG, et
- 1021 al. Air travel is associated with intracontinental spread of dengue virus serotypes
- 1022 1-3 in Brazil. PLoS Negl Trop Dis. 2014;8(4):e2769. doi:
- 1023 10.1371/journal.pntd.0002769. PubMed PMID: 24743730; PubMed Central
- 1024 PMCID: PMC3990485.
- 1025 76. White SK, Mavian C, Salemi M, Morris JG, Jr., Elbadry MA, Okech BA, et al.
- 1026 A new "American" subgroup of African-lineage Chikungunya virus detected in
- 1027 and isolated from mosquitoes collected in Haiti, 2016. PloS one.
- 1028 2018;13(5):e0196857. Epub 2018/05/11. doi: 10.1371/journal.pone.0196857.
- 1029 PubMed PMID: 29746539; PubMed Central PMCID: PMC5944945.
- 1030 77. Obolski U, Perez, P. N., Villabobna-Arenas, C. J., Faria, N. R., Lourenco, J.
- 1031 MVSE: an R-package that estimates a climate-driven mosquito-borne viral
- 1032 suitability index. bioRxiv. 2018. doi: <https://doi.org/10.1101/360701>.
- 1033 78. Cherian SS, Walimbe AM, Jadhav SM, Gandhe SS, Hundekar SL, Mishra AC,
- 1034 et al. Evolutionary rates and timescale comparison of Chikungunya viruses
- 1035 inferred from the whole genome/E1 gene with special reference to the 2005-07

1036 outbreak in the Indian subcontinent. Infection, genetics and evolution : journal of
 1037 molecular epidemiology and evolutionary genetics in infectious diseases.
 1038 2009;9(1):16-23. Epub 2008/10/23. doi: 10.1016/j.meegid.2008.09.004.
 1039 PubMed PMID: 18940268.
 1040




Figure 1




Figure 2




Figure 3









a**b****c****d****e****f****g****h**

Figure 4