

# **Sex and Estrous Cycle Effects on Anxiety- and Depression-related Phenotypes in Two-hit Developmental Stress Model**

**Ivana Jaric<sup>1#</sup>, Devin Rocks<sup>1#</sup>, Heining Cham<sup>2</sup>, Alice Herchek<sup>1</sup>, and Marija Kundakovic<sup>1\*</sup>**

<sup>1</sup>Department of Biological Sciences, Fordham University, Bronx, NY

<sup>2</sup>Department of Psychology, Fordham University, Bronx, NY

<sup>#</sup>These two authors contributed equally to this work

**\*Correspondence:**

Marija Kundakovic  
Fordham University  
Department of Biological Sciences  
441 E. Fordham Road  
Larkin Hall, Room 160  
Bronx, NY 10458  
Email: [mkundakovic@fordham.edu](mailto:mkundakovic@fordham.edu)

## Abstract:

Stress during sensitive developmental periods can adversely affect physical and psychological development and contribute to later-life mental disorders. In particular, adverse experiences during childhood dramatically increase the risk for the development of depression and anxiety disorders. Although women of reproductive age are twice as likely to develop anxiety and depression than men of the corresponding age, little is known about sex-specific factors that promote or protect against the development of psychopathology. To examine potential developmental mechanisms driving sex disparity in risk for anxiety and depression, we established a two-hit developmental stress model including maternal separation in early life followed by social isolation in adolescence. Our study shows complex interactions between early-life and adolescent stress, between stress and sex, and between stress and female estrogen status in shaping behavioral phenotypes of adult animals. In general, increased locomotor activity and body weight reduction were the only two phenotypes where two stressors showed synergistic activity. In terms of anxiety- and depression-related phenotypes, single exposure to early-life stress had the most significant impact and was female-specific. We show that early-life stress disrupts the protective role of estrogen in females, and promotes female vulnerability to anxiety- and depression-related phenotypes associated with the low-estrogenic state. We found plausible transcriptional and epigenetic alterations in psychiatric risk genes, *Nr3c1* and *Cacna1c*, that likely contributed to the stress-induced behavioral effects. In addition, two general transcriptional regulators, *Egr1* and *Dnmt1*, were found to be dysregulated in maternally-separated females and in animals exposed to both stressors, respectively, providing insights into possible transcriptional mechanisms that underlie behavioral phenotypes. Our findings provide a novel insight into developmental risk factors and biological mechanisms driving sex differences in depression and anxiety disorders, facilitating the search for more effective, sex-specific treatments for these disorders.

## Introduction

Stressful experiences during sensitive developmental periods can have a major impact on later-life physical and mental health (McEwen, 2008; Romeo and McEwen, 2006). In particular, adverse experiences during childhood, such as abuse or neglect, dramatically increase the risk for the development of psychiatric disorders including depression, anxiety disorders, and substance abuse (Heim and Nemeroff, 2001; Lupien et al., 2009). It has been proposed that stress in early life induces increased reactivity within the hypothalamic-pituitary-adrenal (HPA) axis, sensitizing an individual to future stressful situations and challenges that may precipitate psychiatric disorders (Lupien et al., 2009). It is notable, though, that only a subset of individuals that experience early-life trauma develop a psychiatric condition in their adult life, and it is of great interest to reveal the factors that promote both risk and resilience to the effects of early life adversity.

The majority of psychiatric conditions linked to early-life adversity show a substantial sex bias in prevalence and severity, yet little is known about sex-specific factors that promote or protect against the development of psychopathology (Altemus et al., 2014). One of the major reasons for the dearth of knowledge in this area has been a traditional focus on male subjects in experimental neuroscience research (Beery and Zucker, 2011; Will et al., 2017). Some insights have been provided by the studies of depression and anxiety disorders, which show striking sex bias being twice as frequent in women than in men (Kuehner, 2017; Li and Graham, 2017). The increased risk in females emerges with the onset of puberty and various findings have suggested that natural hormonal shifts predispose females to stress-related, mood and anxiety disorders (Altemus et al., 2014; Kuehner, 2017; Li and Graham, 2017; Steiner et al., 2003). How sex hormones and stress may interact to bring about increased vulnerability in females remains unknown.

To examine potential developmental mechanisms driving sex disparity in risk for anxiety and depression, we established a two-hit developmental stress model including maternal separation in early life followed by social isolation in adolescence. Repeated and prolonged separations of offspring from their mothers is one of the most frequently used rodent paradigms modeling aspects of childhood maltreatment in humans (Heim and Nemeroff, 2001; Kundakovic and Champagne, 2015). This model has been associated with disrupted brain development, and emotional and cognitive dysregulation in later life, showing translational validity (Heim and Nemeroff, 2001; Kundakovic and Champagne, 2015). However, the results have been inconsistent among different laboratories, largely due to different rodent strains used as well as different nature and timing of manipulation (Murthy and Gould, 2018). We use an established maternal separation protocol in C57BL/6 mice, combining maternal deprivation with disrupted maternal care, which was previously shown to induce depression-like behavior in early adolescent male and female animals (Kundakovic et al., 2013b). The rationale of the current study was to introduce another stress hit in adolescence and to examine whether increased female vulnerability to stress may emerge as females start experiencing cycling sex hormone levels following the onset of puberty. We chose a "suboptimal", mild adolescent stress paradigm, three-week social isolation, previously shown to be insufficient to trigger behavioral abnormalities but exerting a synergistic disruptive effect on depression-related behavior when combined with another psychiatric risk factor (Niwa et al., 2013).

Our study provides a comprehensive analysis of the effects of early-life stress and adolescent stress, individually and in combination, on anxiety- and depression-related behavioral

phenotypes in young adult male and female mice, accounting for the effects of the naturally cycling estrogen levels in females. Using several candidate genes of relevance to these behaviors, we also attempted to find molecular correlates that would link the effect of stressors to behaviors of interest. We examined both gene expression and DNA methylation in the ventral hippocampus, an area essential for emotional regulation, looking for a possible epigenetic link between stressful exposures and behavioral outcomes. We find complex interactions between two stressors, sex, and female estrogen status, and provide a new biological insight into the increased female vulnerability to stress and stress-related disorders.

## Material and Methods

**Study Design (Figure 1):** To assess the effects of early-life stress and adolescent stress, individually and in combination, on behavioral and molecular outcomes, the following treatment groups were generated: 1) No stress (**CON**); 2) Early life stress - maternal separation from postnatal day (PD) 1-14 (**MS**); 3) Social isolation in adolescence from PD35-56 (**SI**); and 4) Two-hit stress group - early life MS + SI in adolescence (**MS\_SI**). In all groups, behavioral testing was performed from PD57-68, during the animals' light cycle, and in the following order (from least to most stressful): open-field test; elevated plus maze; sucrose preference; and forced swim test. Animals were sacrificed two days after the last behavioral test (PD70). Molecular analyses were performed on littermates that were not behaviorally tested to avoid the effect of behavioral testing on gene expression and DNA methylation. To avoid possible litter effects, a maximum of two pups per sex per litter were tested on behavioral measures, and one pup per sex per litter were used for molecular analyses.

**Animals.** Male (N = 40) and female (N = 80) C57BL/6 mice were ordered from Jackson Laboratories and housed in same-sex groups of five for 2.5 weeks prior to mating to habituate to the facility. Mating pairs consisting of two females and one male were separately housed and yielded N = 72 litters which were counted and weighed at PD0. 18 litters were randomly assigned to each group (CON, MS, SI, and MS\_SI). Litters were weighed again on PD14, and individuals from the litters were weighed on weaning day, PD28, at which point they were housed in same-sex and same-group cages of five. Regardless of the treatment group, all animals were group-housed until PD35. Animals were kept on a 12:12 light/dark cycle (lights on at 8 a.m.) and given ad libitum access to food and water. All animal procedures were approved by the Institutional Animal Care and Use Committee at Fordham University.

**Early-life stress.** We used a previously established maternal separation protocol in C57BL/6 mice (Kundakovic et al., 2013b). Litters selected for early-life stress (MS and MS\_SI groups) underwent maternal separation for three hours every day for two weeks (PD1-14). During this time the dam was removed and placed in a temporary housing cage with food and water available. The time of day that maternal separation occurred was changed daily so that the stress was unpredictable. At a random time within the three-hour maternal separation window, dams were exposed to either restraint stress or forced swim to reduce the possibility of the dams giving compensatory maternal care to their litters following separation which was previously reported (Millstein and Holmes, 2007). In fact, we observed that this unpredictable stress disrupts maternal behavior for at least an hour after dams return to their pups; MS mothers spent more

time out of nest and showed less licking/grooming and arched-back nursing compared to control mothers. Restraint stress lasted 20 minutes and involved placing the dam in a conical tube that restricted movement. Forced swim lasted 2 minutes and involved placing the dam in a 2L glass beaker with 1L of water, after which the dams were dried and returned to the temporary housing cage.

**Adolescent stress.** Animals selected for adolescent stress (SI and MS\_SI groups) underwent social isolation for three weeks starting at PD35 and ending at PD56. Social isolation involved placing individual animals in separate cages without cagemates. Socially isolated animals were given ad libitum access to food and water. At PD56, all animals were again housed in same-sex, same-treatment groups.

**Open-Field Test.** On PD57, animals from each group were tested using the Open Field paradigm. At the time of testing the animal was placed in the corner of the 40cm x 40cm x 35cm glass box apparatus (Stoelting). Movement was tracked for 15 minutes using a camera and ANY-maze video tracking software (ANY-maze 5.1, Stoelting Co; IL). The total distance traveled, the amount of time the animal spent in the center of the maze versus the periphery, and the latency to the first center entry were recorded. The results of the first two minutes of the test were used for the analysis.

**Elevated Plus Maze Test.** On PD60, animals from each group were tested using the Elevated Plus Maze paradigm. At the time of testing the animal was placed in the center of the apparatus, which was a raised (40cm) plus-shaped platform with two 35cm x 5cm arms, one being protected by 15 cm walls (Stoelting). Movement was tracked for 5 minutes using a camera and ANY-Maze video tracking software. The total distance traveled, the amount of time the animal spent in the open arms of the maze versus the closed arms, and the latency to the first open arm entry were recorded. The results of the first two minutes of the test were used for the analysis.

**Sucrose Preference Test.** From PD62-65, animals from each group were individually housed and given access to two water bottles with one containing 1% sucrose solution. Water bottles were weighed daily to assess sucrose preference. The position of the water bottles was switched each day to eliminate the possibility of attaining erroneous results due to place preference. The percentage of the sucrose solution consumed over the total volume of fluid consumed was reported as sucrose preference.

**Forced Swim Test.** On PD68, animals from each group were tested using the Forced Swim paradigm. Animals were placed in 2L glass beakers filled with 1L of water. Movement was tracked for 6 minutes using a camera and ANY-Maze video tracking software. The amount of time the animal spent frozen (immobile) and the latency to freeze were recorded. To ensure the accuracy of data obtained by automated video tracking of this test, the data were additionally evaluated and confirmed by watching the video recordings.

**Estrous cycle stage identification.** To account for sex hormone status of female animals, vaginal smears were taken just prior to behavioral testing and sacrificing for all female animals. Smears were analyzed by cytology and estrous cycle stage was recorded following previously established protocols (McLean et al., 2012). Briefly, smears were allowed to dry and then stained

with crystal violet dye. Smears were then analyzed using light microscopy. Proestrus is characterized by a high number of nucleated epithelial cells. Estrus is characterized by a high number of cornified epithelial cells. Metestrus is characterized by mostly cornified epithelial cells, some nucleated epithelial cells, and some leukocytes. Diestrus is characterized by cornified epithelial cells, nucleated epithelial cells, and many more leukocytes compared to metestrus. For the purpose of behavioral data analysis (for each test except for the sucrose preference test), female animals were classified into two groups based on the estrogen status on the day of test: 1) *high-estrogen status* (proestrus, diestrus/proestrus transition, and proestrus/estrus transition), and 2) *low-estrogen status* (diestrus, estrus, metestrus, estrus/metestrus transition, and metestrus/diestrus transition).

**RNA and DNA Isolation.** From each group, a subset of male (n = 8/group) and female (n = 12/group) animals which did not undergo behavioral testing were sacrificed at PD70 for molecular analysis. We checked the estrous cycle stage at the time of sacrificing, and as much as possible, we balanced the number of high- and low-estrogenic females in each group. Immediately after sacrificing, whole brains were isolated, frozen, and then stored at -80°C. The ventral hippocampus was later dissected, the tissue was immediately homogenized and DNA and RNA were extracted simultaneously using the Qiagen AllPrep DNA/RNA Mini Kit following the manufacturer's instructions. This served as the starting material for all subsequent molecular analyses.

**Gene Expression Analysis.** The analysis of gene expression was performed as described previously (Kundakovic et al., 2013a). Briefly, cDNA was synthesized from 500 ng of total RNA using the SuperScript III First-Strand Synthesis (Invitrogen) kit following the manufacturer's protocol. qRT-PCR was then performed in QuantStudio 3 (Applied Biosystems) using Fast SYBR Green Master Mix (Applied Biosystems) and RT-PCR primers specific to the targeted genes (**Suppl. Table 1**). Relative mRNA expression was calculated using the standard  $\Delta\Delta CT$  method. *Ppia* gene was used as a reference gene, previously shown not to be affected by early-life stressors (Kundakovic et al., 2013a; Kundakovic et al., 2013b). All data are expressed relative to the male control group.

**DNA Methylation Analysis.** DNA methylation analysis was performed at single base resolution using the bisulfite pyrosequencing method as previously described (Kundakovic et al., 2013a). Briefly, 250 ng of genomic DNA was used for bisulfite conversion (EpiTect Bisulfite Kit, Qiagen) and 20 ng of bisulfite converted DNA was run in a PCR using the PyroMark PCR kit (Qiagen) and primers for genes of interest, one of which was biotinylated at the 5' end. Biotinylated PCR products were processed on a vacuum workstation and pyrosequencing was performed on an Advanced PyroMark Q24 pyrosequencer using the advanced CpG reagents (Qiagen) and a specific sequencing primer. PCR and pyrosequencing primers for the mouse *Nr3c1* and *Cacna1c* genes are presented in **Suppl. Table 2**. Methylation levels of single CpG sites were quantified using PyroMark Q24 Advanced software (Qiagen).

**Statistical Analysis.** For all analyses, we did two-way ANOVA with group and sex as factors. We first looked for group by sex interaction, and if the interaction was not significant, it was dropped from the model and we tested the main effects of group and sex. For the analyses of behavior (with the exception of sucrose preference), we did an additional two-way ANOVA

analysis in females in which we assessed the effects of group and estrogen status and their interaction. If the group by estrogen interaction was not significant, it was dropped from the model, and we assessed the main effects of group and estrogen status. All post-hoc tests were performed using Tukey's method. The analyses were performed in R version 3.5.1 and the graphs were generated using the R ggpubr package. Box plots show the first and third quartiles; the horizontal line is the median; the added, inner circles mark the mean with standard error for each box and small, outer circles mark possible outliers. The results were considered significant at significance level of  $P < 0.05$ .

## Results

**Body weight.** We monitored the effect of stressors on body weight throughout the study, including post-weaning (PD28), prior to social isolation (PD35), and after social isolation (PD56) time points (**Figure 2**). At all time points, males had higher body weight compared to females, and this relationship was not affected by any of the treatments (no significant group by sex interaction was found). Maternal separation led to significantly reduced body weight in both male and female offspring which was evident at PD28 ( $P < 0.001$ , **Fig. 2A**) and persisted through PD35 ( $P < 0.001$ , **Fig. 2B**) and PD56 ( $P < 0.001$ , **Fig. 2C**). At PD56, we found that social isolation in males and females also led to a significant reduction in body weight compared to control mice ( $P = 0.002$ , **Fig. 2C**). We found no significant difference between the effects of maternal separation and social isolation on body weight ( $P = 0.6$ , **Fig. 2C**). However, the body weight reduction was most profound in mice that underwent the double-stress paradigm, surpassing the effects of a single exposure to either maternal separation ( $P = 0.02$ ) or social isolation ( $P < 0.001$ , **Fig. 2C**).

**Anxiety-like behavior.** In all four groups of animals, we performed two well-established tests for anxiety-like behavior: open-field and elevated plus maze tests.

*Open Field.* In the open field, we found no significant group by sex interaction on total distance travelled; however, we found a significant effect of group on this variable ( $F (3, 148) = 2.87$ ,  $P = 0.04$ ) (**Fig. 3A**). Posthoc test revealed that this effect was driven by the double-stress (MS\_SI) group which was marginally more active than any other group including CON ( $P = 0.08$ ), MS ( $P = 0.09$ ), and SI ( $P = 0.08$ ). The average increase in the activity was higher in male MS\_SI ( $11.07 \pm 0.52$  m) vs. CON ( $9.27 \pm 0.60$  m) comparison than in female MS\_SI ( $11.15 \pm 0.41$  m) vs. CON ( $10.22 \pm 0.90$  m) comparison. There was no difference in activity between CON, MS, and SI groups ( $P > 0.99$  for all comparisons). No significant effect of group or estrogen, or an interaction between group and estrogen were found when females were considered separately from males (**Fig. 3B**).

When we analyzed the latency to the first entry into the center of the open field, no significant effect of group or sex nor their interaction was found with regard to this variable (**Fig. 3C**). However, we found a significant effect of estrogen status ( $F (1, 82) = 13.5$ ,  $P < 0.001$ ); high-estrogenic females showed shorter latency time to enter the center compared to low estrogenic females (**Fig. 3D**).

Finally, we analyzed time spent in the center of the open field. We found no significant group by sex interaction and no significant main effects of group or sex on this variable (**Fig. 3E**). However, when females were considered separately, we found a significant group by estrogen interaction ( $F (3, 79) = 5.47$ ,  $P = 0.002$ ) (**Fig. 3F**). In addition, both main effects of

group ( $F(3, 79) = 9.12, P < 0.001$ ) and estrogen status ( $F(1, 79) = 39.34, P < 0.001$ ) were significant. In general, high estrogenic females showed more time spent in the center of the open field than low estrogenic females. Interestingly, in posthoc test, the effect of group was found to be significant in high-estrogenic females only; MS ( $P < 0.001$ ), SI ( $P = 0.003$ ), and MS\_SI ( $P = 0.01$ ) high estrogenic females were all found to spend significantly less time in the center of the open field than control high-estrogenic females. Another interesting observation is that a significant difference in the time spent in the center between high and low-estrogenic females was found only in control ( $P < 0.001$ ) and SI ( $P = 0.01$ ) groups, whereas this difference was lost in groups that were exposed to early life stress: MS ( $P = 0.32$ ) and MS\_SI ( $P = 0.09$ ) groups (**Fig. 3F**).

*Elevated Plus Maze.* No significant effect of group or sex nor their interaction was found on total distance traveled in the elevated plus maze (**Fig. 4A**). There was a significant effect of estrogen status on activity in the maze ( $F(1, 82) = 4.28, P = 0.04$ ) with high-estrogenic females showing marginally higher activity than low-estrogenic females (**Fig. 4B**).

With regard to the latency to the first open arm entry, we found a significant group by sex interaction ( $F(3, 144) = 3.35, P = 0.02$ ) (**Fig. 4C**). The effect of group showed trend toward significance in males only ( $P = 0.07$ ) due to a longer latency to enter the open arm in SI male group compared to CON males ( $P = 0.1$ ). Furthermore, the significant group by sex interaction that we observed was due to a sex difference induced in SI group, in which SI males showed significantly delayed latency to enter the open arms compared to SI female group ( $P = 0.002$ ). No significant sex difference was found in CON, MS, or MS\_SI groups ( $P > 0.99$  for all comparisons). In addition, we found a significant effect of estrogen status in females ( $F(1, 82) = 4.39, P = 0.04$ ); on average, high estrogenic females took longer to enter the open arm than low estrogen females (**Fig. 4D**).

We found no significant group by sex interaction and no significant effect of group on time spent in the open arms of the elevated plus maze (**Fig. 4E**). However, we found a significant effect of sex ( $F(1, 148) = 17.78, P < 0.001$ ), with females spending, on average, more time in the center of the maze than males (**Fig. 4E**). This effect appears to be primarily driven by the estrogen status which was found to have a significant effect on time spent in the open arms in females ( $F(1, 82) = 131.85, P < 0.001$ ), with high estrogenic females spending on average much longer time in the open arms than low estrogenic females (**Fig. 4F**).

**Depression-like behavior.** Two well established tests for depression-like behavior were performed in all four groups of animals: sucrose preference and forced swim test.

*Sucrose preference test* represents a model of anhedonia in humans and we found a statistically significant interaction between the effects of group and sex on sucrose preference in our study ( $F(3, 145) = 2.87, P = 0.04$ ) (**Fig. 5A**). This interaction was primarily driven by a sex difference in sucrose consumption observed in MS group; MS females showed a significantly decreased sucrose preference when compared to MS males ( $P = 0.046$ ). Sex difference in sucrose preference did not exist in control group ( $P = 0.99$ ) or in other treatment groups, SI ( $P = 0.39$ ) and MS-SI ( $P > 0.99$ ) groups. Of note, we did not examine the effect of estrogen status in this test because the test spans three days or more than a half of the mouse estrous cycle.

*Forced swim test.* We analyzed time spent immobile in the forced swim test and found a significant group by sex interaction in this test ( $F(3, 145) = 3.07, P = 0.03$ ) (**Fig. 5B**). In posthoc

tests, we found no significant difference between groups and only trend for immobility time difference between males and females in the MS group ( $P = 0.13$ ). In females, we found no significant group by estrogen interaction and no significant main effect of group. However, we found a significant effect of the estrogen status ( $F (1, 82) = 211.80, P < 0.001$ ), with high-estrogenic females generally having significantly shorter immobility time in the forced swim test than low estrogenic females (Fig. 5C).

**Gene expression analysis.** We determined the effects of early-life stress and adolescent stress on expression of the following candidate genes in the ventral hippocampus: *Bdnf* (encoding brain-derived neurotrophic factor); *Nr3c1* (encoding glucocorticoid receptor); *Egr1* (encoding early growth response protein 1); *Cacna1c* (encoding calcium voltage-gated channel subunit alpha1C), *Gria3* (encoding glutamate ionotropic receptor AMPA type subunit 3); and *Dnmt1* (encoding DNA methyltransferase 1). The ventral hippocampus was selected because it is strongly implicated in emotion regulation and anxiety-related behavior (Bannerman et al., 2003; Fanselow and Dong, 2010; Kheirbek et al., 2013). The *Bdnf* and *Nr3c1* genes have been previously implicated in the effects of early-life stress and in psychopathology (Angelucci et al., 2005; Boulle et al., 2012; Kundakovic et al., 2015; Kundakovic et al., 2013b; McGowan et al., 2009; Roth et al., 2009; Weaver et al., 2004). *Egr1* has been reported to be sensitive to stress and encodes an immediate early gene product, a transcription factor which regulates expression of numerous genes including *Nr3c1* (Gutiérrez-Mecinas et al., 2011; Rusconi et al., 2016; Weaver et al., 2004; Xie et al., 2013). *Cacna1c* and *Gria3* are well-established psychiatric risk genes (de Sousa et al., 2017; Dedic et al., 2018; Henley and Wilkinson, 2016). Finally, *Dnmt1* encodes an epigenetic regulator that has been implicated in epigenetic effects of early-life stressors and in psychiatric disorders (Dong et al., 2015; Grayson and Guidotti, 2013; Kundakovic et al., 2013a).

We found no significant group by sex interaction and no significant effect of group or sex on the expression of *Bdnf* in the ventral hippocampus (Fig. 6A). With regards to *Nr3c1*, there was no significant group by sex interaction, but there were significant main effects of both group ( $F (3, 75) = 5.21, P = 0.003$ ) and sex ( $F (1, 75) = 32.97, P < 0.001$ ) on the expression of this gene (Fig. 6B). Posthoc test showed a significant difference between MS and CON groups only ( $P = 0.001$ ). This effect was driven by an increased *Nr3c1* expression in the MS female group ( $1.39 \pm 0.04$ ) compared to CON female ( $1.08 \pm 0.06$ ) and CON male ( $1.00 \pm 0.03$ ) groups, whereas the MS male group had *Nr3c1* expression ( $1.08 \pm 0.04$ ) comparable to controls (Fig. 6B). In general, female mice also had a higher *Nr3c1* expression in the ventral hippocampus than males ( $P < 0.001$ ) (Fig. 6B). *Egr1* showed a similar pattern of expression to *Nr3c1*; there was no significant group by sex interaction but there were significant main effects of group ( $F (3, 68) = 3.47, P = 0.02$ ) and sex ( $F (1, 68) = 4.66, P = 0.03$ ) on the expression of this gene (Fig. 6C). The only two groups that showed a statistically significant difference in *Egr1* expression were MS and CON groups ( $P = 0.02$ ) which, again, was driven by a higher expression in the MS female group ( $1.66 \pm 0.21$ ) compared to CON female ( $0.98 \pm 0.06$ ) and CON male ( $0.88 \pm 0.05$ ) groups and the MS male group had *Egr1* expression ( $1.03 \pm 0.07$ ) more comparable to controls (Fig. 6C). In general, females had a higher *Egr1* expression compared to males (Fig. 6C). The *Cacna1c* gene showed the most variable expression among the tested groups. We found a significant group by sex interaction for this gene ( $F (3, 72) = 14.75, P < 0.001$ ) (Fig. 6D). In females, we found significantly reduced expression of this gene in all three treatment groups including MS ( $P = 0.02$ ), SI ( $P < 0.001$ ), and MS\_SI ( $P = 0.02$ ) compared to control female animals. In males, though, we found a significant reduction in *Cacna1c* expression in CON vs. MS comparison ( $P <$

0.001) and in CON vs. SI comparison ( $P < 0.001$ ) but there was no significant difference between CON and MS\_SI groups ( $P = 0.63$ ). However, the double-stress MS\_SI male group had a significantly higher expression than both single-stress exposed male groups – MS ( $P < 0.001$ ) and SI ( $P < 0.001$ ) groups. Of all groups, only CON ( $P = 0.03$ ) and MS\_SI groups ( $P < 0.001$ ) showed a significant sex difference in *Cacna1c* expression; this difference was completely lost in MS and SI groups ( $P > 0.99$  for both comparisons) (Fig. 6D). Unlike what we observed with *Cacna1c*, we found no significant group by sex interaction and no significant effect of group or sex on the expression of *Gria3* in the ventral hippocampus (Fig. 6E). Finally, in terms of *Dnmt1* expression in the ventral hippocampus, there was no significant group by sex interaction but we found significant main effects of both group ( $F(3, 75) = 19.68, P < 0.001$ ) and sex ( $F(1, 75) = 9.60, P = 0.003$ ) (Fig. 6F). Two treatment groups SI ( $P = 0.003$ ) and MS\_SI ( $P < 0.001$ ) showed a significantly higher expression of *Dnmt1* than control group; there was no significant difference between MS and control group ( $P = 0.97$ ). Among the treatment groups, both SI ( $P = 0.01$ ) and MS\_SI groups ( $P < 0.001$ ) had a significantly higher expression of *Dnmt1* than MS group. The highest expression of *Dnmt1* was seen in the double-stress MS\_SI group, significantly surpassing the effects of a single exposure to social isolation in SI group ( $P = 0.01$ ) (Fig. 6F).

**DNA methylation analysis.** To address possible epigenetic mechanisms underlying gene expression changes that we have observed, we performed DNA methylation analysis of the two candidate genes: *Nr3c1* and *Cacna1c*. We analyzed 8 CpG sites located in the 5' regulatory region of the *Nr3c1* gene, containing the binding site for transcription factor Egr1 (or NGFI-A) overlapping CpG sites 7 and 8 (Fig. 7A) (Kundakovic et al., 2013b; Weaver et al., 2004). For CpG1 site, we found a significant group by sex interaction ( $F(3, 40) = 4.67, P = 0.01$ ) (Fig. 7A). Specifically, MS\_SI males had significantly higher methylation at this site than CON ( $P = 0.02$ ) and SI ( $P = 0.02$ ) males. In addition, MS\_SI males had higher methylation levels than MS\_SI females ( $P = 0.003$ ), whereas sex difference was not found in any other group. We found no significant main effects of group or sex nor their interaction when we analyzed CpG site 2 (Fig. 7A). We found a significant main effect of group ( $F(3, 43) = 3.98, P = 0.01$ ) on methylation levels of CpG site 3, which was driven by increased methylation in MS\_SI group compared to CON ( $P = 0.02$ ) and SI ( $P = 0.03$ ) groups (Fig. 7A). For CpG site 4, we found a significant effect of both group ( $F(3, 43) = 5.51, P = 0.003$ ) and sex ( $F(1, 43) = 7.30, P = 0.01$ ) (Fig. 7A). We found significantly lower CpG 4 methylation in MS ( $P = 0.003$ ), SI ( $P = 0.04$ ), and MS\_SI ( $P = 0.01$ ) groups compared to controls. In general, males had a higher methylation at CpG site 4 than females ( $P = 0.01$ ). We found significant effect of both group ( $F(3, 43) = 4.91, P = 0.01$ ) and sex ( $F(1, 43) = 5.77, P = 0.02$ ) on CpG5 methylation (Fig. 7A). This was driven by decreased methylation in SI group compared to CON ( $P = 0.005$ ) and MS\_SI ( $P = 0.03$ ) groups. In general, CpG 5 methylation was higher in males than in females. For CpG6, we found a significant effect of group ( $F(3, 43) = 4.95, P = 0.005$ ) and sex ( $F(1, 43) = 4.12, P = 0.049$ ) (Fig. 7A). Significant group difference was found in MS\_SI to SI comparison only, where SI group had significantly lower CpG6 methylation than MS\_SI group ( $P = 0.002$ ). Males had higher methylation levels than females. For CpG7 site, we found a significant group by sex interaction ( $F(3, 40) = 4.32, P = 0.01$ ). A significant effect of group was found in MS vs. CON group comparison in females only ( $P = 0.04$ ), where we found significantly reduced CpG7 methylation in MS females (Fig. 7A). CpG7 methylation was also marginally higher in MS\_SI male group when compared to CON male group ( $P = 0.09$ ). These results were consistent with observed sex difference, which

was statistically significant within MS\_SI group only ( $P = 0.001$ ) and showed trend toward significance in MS group ( $P = 0.06$ ), with CpG7 methylation being higher in males of both groups. Finally, we found a significant group by sex interaction with regards to methylation of CpG site 8 ( $F(3, 40) = 3.50, P = 0.02$ ) (Fig. 7A). In posthoc test, we found only significant decrease in CpG8 methylation in two female treatment groups – MS ( $P = 0.04$ ) and MS\_SI ( $P = 0.01$ ) – compared to CON female group. No significant sex difference was observed in any of the groups.

We also analyzed 4 CpG sites in the first intron of the *Cacna1c* gene (Fig. 7B). The analyzed region is within 1 kb from a transcription start site, and is located in the CpG island shore having high homology between the mouse and human genome. In addition, these sites were previously shown to have cell-type specific methylation patterns (Nishioka et al., 2013). For CpG1, we found a significant main effect of group ( $F(3, 43) = 3.43, P = 0.03$ ) which was driven by a decreased CpG1 methylation found in MS\_SI group compared to CON group ( $P = 0.02$ ) (Fig. 7B). For CpG2, we found a marginally significant effect of group ( $F(3, 43) = 2.79, P = 0.05$ ) which was again driven by reduced CpG2 methylation in MS\_SI group compared to CON group showing trend toward significance ( $P = 0.1$ ) (Fig. 7B). For CpG3, we found a similar trend with the main effect of group ( $F(3, 43) = 2.2, P = 0.1$ ) due to marginally significant reduction in CpG site 3 in MS\_SI group compared to controls ( $P = 0.07$ ) (Fig. 7B). There was also a significant effect of group on CpG4 methylation ( $F(3, 43) = 2.92, P = 0.04$ ), again driven by reduced methylation in MS\_SI group but this time compared to SI group ( $P = 0.04$ ) (Fig. 7B). We found no significant effect of sex on methylation of any of the CpG sites analyzed in the *Cacna1c* gene.

## Discussion

In this study, we examined the effects of early-life stress and adolescent stress, individually and in combination, on body weight, anxiety- and depression-related behaviors, and expression and DNA methylation of genes implicated in stress-related psychiatric disorders. Overall, we found that a single, early-life stress exposure had the most profound effect on anxiety- and depression-related phenotypes in adult animals. Interestingly, the 'second hit' of a suboptimal stress exposure in adolescence frequently attenuated the effects of early-life stress. Moreover, lasting effects of early-life stress are largely found to be female-specific and we have found significant effects of estrogen status on anxiety- and depression-like behavior in both control female animals and in those exposed to stress.

The current study is the continuation of our previous study which examined the effects of early-life maternal separation on early adolescent male and female offspring (Kundakovic et al., 2013b). The two developmental stressors that we used in the current study were sufficient to induce metabolic dysfunction in animals, as we have seen through the persistent reduction in body weight observed in all treatment groups. Interestingly, reduced body weight is a rare phenotype on which two stress hits worked synergistically, surpassing the effects of individual stress exposures. Another phenotype that emerged only in the double-hit stress group is hyperactivity in the open field test, with males being more strongly affected. While increased locomotor activity is a rather non-specific phenotype, which, interestingly, did not affect any of the indices of anxiety or depression-related behavior, this phenotype is seen after the treatment with psychostimulants and is a reminiscent of hyperactivity seen in disorders such as attention deficit hyperactivity disorder (ADHD). Psychotic disorders and ADHD have been linked to

developmental stress with males being more severely impacted (Bale et al., 2010), similar to what we observed in our study. Therefore, the double-stress paradigm that we constructed may be of relevance for studies of metabolic dysfunction and psychosis- or hyperactivity-related disorders but we found surprisingly small effects of this paradigm with regards to anxiety- and depression-related behaviors.

There are several possible explanations for the observed "buffering" effect of adolescent social isolation on the effect of early-life stress on anxiety- and depression-related phenotypes. First, three weeks of adolescent social isolation is a mild stressor. The only phenotype that we observed in the SI group is a marginally delayed latency to the open arm entry in males, which is as an index of increased anxiety-like behavior. Other than that, the behavioral effects of this paradigm were largely non-significant, in line with what was previously reported (Niwa et al., 2013). Our results suggest that mild stress in adolescence may decrease the risk for the development of anxiety and depression in individuals that experienced certain forms of early-life trauma. While this finding may be counterintuitive, it is consistent with the "mismatch hypothesis" of psychiatric disease (Nederhof, 2012; Schmidt, 2011). This hypothesis states that early-life environment induces adaptive phenotypic changes, preparing an individual for later-life environmental conditions (Nederhof, 2012). If the early-life and later-life environment match, the early-life stress may even be beneficial. However, the disorder may occur once the later-life conditions do not match stressful environment experienced in early life. As an example, it was shown that rats who receive low levels of maternal care (form of early-life stress) exhibit enhanced learning under stressful conditions but impaired learning under normal conditions in later life (Champagne et al., 2008). There are also additional "two-hit stress" studies that showed increased stress resilience in animals that experienced early-life stress (Biggio et al., 2014; Hsiao et al., 2016; Santarelli et al., 2017). However, while the "mismatch hypothesis" is plausible to explain the lack of anxiety- and depression-related phenotypes in our two-hit stress model, it is likely that a stronger stressor during adolescence or adulthood would, indeed, worsen the phenotype induced by early-life stress, as seen in some studies (Pena et al., 2017). In addition, in humans, genetic vulnerability or resilience to depression and anxiety disorders, certainly play an important role in how early- and later-life stress affect the risk for the development of these disorders.

As compared to our previous study which showed that maternal separation induces decreased sucrose preference, a form of anhedonia or depression-like behavior, in both male and female early adolescent mice (Kundakovic et al., 2013b), here we show that this effect persists into adulthood only in female animals. It is well known that anxiety and depression show explicit sex bias with women of reproductive age being twice as likely as men of the corresponding age to develop these disorders. The mechanisms underlying this sex disparity are unknown, and the studies of early-life stress have not necessarily addressed this (Bale et al., 2010), particularly because a lot of studies were focused on male animals. Our study provides an important insight into this sex disparity, providing a clue how early-life stress can be a sex-specific factor in driving vulnerability to anxiety and depression. Indeed, the early-life paradigm that we use, combining maternal separation with unpredictable maternal stress, turned out to be a good model to study increased female vulnerability to these disorders.

Among the most interesting sex-specific findings in our study is the effect of estrogen status and its interaction with early-life stress on anxiety- and depression-related behaviors in females. In fact, our most consistent finding is that the high-estrogen phase of the cycle is associated with remarkably lower indices of anxiety and depression-like behavior compared to

the low-estrogen phase. This was basically seen in all tests in which estrogen status was controlled for, including: open field (shorter latency and longer time spent in the center in high-estrogenic females); elevated plus maze (longer time spent in the open arms in high estrogenic females); and forced swim test (shorter time spent immobile in high estrogenic females). Our results are consistent with earlier studies in rats (Marcondes et al., 2001; Walf and Frye, 2006) and clearly indicate the importance of estrogen status for anxiety and mood regulation in females of reproductive age. Indeed, we can conclude that high estrogen levels have a protective, anxiolytic effect whereas a physiological drop in estrogen may increase risk for the occurrence of anxiety and depression symptoms in females of reproductive age. This is in line with human data which show that a large number of women with major depression experience worsening of their symptoms in the premenstrual, low-estrogen phase of the cycle, even when antidepressant therapy is effective during the remainder of the cycle (Altemus et al., 2014; Kornstein et al., 2005). Importantly, hormonal fluctuations, rather than low hormone levels *per se*, seem to be critical for both pathogenesis and treatment of depression, considering that post-menopausal women with stably low hormone levels appear to lose sensitivity to estrogen (Morrison et al., 2004) and show a similar risk for depression when compared to men of their age (Altemus et al., 2014; Deecker et al., 2008).

Our study provides an important, novel insight into how stress can interact with the ovarian cycle to facilitate the increased anxiety and depression vulnerability in females. When we focused our analysis on female animals, we found that developmental stressors had a significant negative impact on anxiety-like behavior (time spent in the center of the open-field) in our female population. However, when we looked further into the estrous cycle effects, we found that stressors increased anxiety-like behavior only during the high-estrogen phase of the cycle, when the anxiety levels are typically low. No effect of stress was found during the low-estrogen phase, when the anxiety levels are already high. In addition, the most significant effect was found in the early-life stress group. And, even more interestingly, maternally separated animals (MS and MS\_SI groups) showed largely blunted, insignificant difference in anxiety levels between low and high estrogen phases, suggesting that the anxiolytic effect of estrogen is predominantly lost in groups that experience early life stress. Whether this is the consequence of lower estrogen levels or lost sensitivity to estrogen in response to early-life adversity remains to be further explored. Nevertheless, this finding reveals a possible sex-specific mechanism through which early-life adversity may contribute to sex disparity in risk for anxiety and depression by suppressing the "protective" effect of estrogen in women of reproductive age.

We further explored gene expression in the ventral hippocampus with an intent to find molecular correlates of the behavioral effects that we have observed. For this analysis, we did not look at the estrous cycle effects but have explored more general sex-specific patterns of gene expression that may contribute to sex difference in vulnerability. Interestingly, although changes in hippocampal *Bdnf* expression have been found in many studies of early-life adversity (Boersma et al., 2014; Kundakovic et al., 2015; Kundakovic et al., 2013b; Onishchenko et al., 2008; Roth et al., 2009), we found no significant changes in the expression of this gene in our study. This may be due to the fact that we focused our analysis on the ventral hippocampus while previous studies examined the entire hippocampus. Considering that the ventral and dorsal hippocampi are functionally distinct regions (Fanselow and Dong, 2010), showing distinct gene expression profiles (Cembrowski et al., 2016) and distinct responses to chronic stress (Czeh et al., 2015; Pinto et al., 2015), changes previously reported may largely stem from the stressor-induced alterations in the dorsal hippocampus. However, in MS females, we found an

overexpression of *Nr3c1*, which is another gene that has been repeatedly associated with early-life stress in both animals and humans (Champagne et al., 2008; Kundakovic et al., 2013b; McGowan et al., 2009; Suderman et al., 2012; Weaver et al., 2004). This gene encodes the glucocorticoid receptor, directly involved in the negative feedback regulation of the HPA axis, providing one of the possible mechanisms for HPA axis dysregulation and consequent behavioral changes in the MS female group. Another interesting finding is that *Egr1* shows an almost identical expression pattern as *Nr3c1*. Considering that the *Nr3c1* regulatory region contains the binding site for *Egr1* (also known as NGFI-A) (Kundakovic et al., 2013b; Weaver et al., 2004), and *Egr1* is induced by stress (Rusconi et al., 2016; Xie et al., 2013), this provides a plausible link between coordinate up-regulation of these two genes and anxiety- and depression-related phenotypes in the early-life stress female group.

The *Cacna1c* gene was another important candidate gene in our study. This gene has recently received a lot of attention, as it is considered to be one of the strongest genetics hits in psychiatry, associated with risk for bipolar disorder, major depression, and schizophrenia (Ferreira et al., 2008; Green et al., 2010; Sklar et al., 2008). Moreover, embryonic deletion of this gene in the mouse forebrain results in increased anxiety-like behavior and increased susceptibility to chronic stress (Dedic et al., 2018). Some clinical and animal studies have also found a significant interaction between sex and *Cacna1c* genotype (Dao et al., 2010; Takeuchi et al., 2018; Witt et al., 2014); for instance, female, but not male, *Cacna1c* heterozygous mice have been shown to display decreased risk-taking behavior and increased anxiety in multiple tests (Dao et al., 2010). Our study has revealed that this gene is very sensitive to developmental stress and that, depending of the number of hits, the effects of stress can be sex-specific. First, we observed that each individual stressor, maternal separation and social isolation, decreased *Cacna1c* mRNA levels in both sexes. Once these two stressors are combined, though, females still show reduced expression, but this effect is lost in males. These results provide a good example where deficiency in a single gene cannot be simply associated with behavioral phenotype. For instance, we can speculate that the down-regulation of *Cacna1c* expression contributed to increased anxiety-like behavior observed in females of all groups and in SI males, with MS females being most significantly affected because these animals also show dysregulated expression of *Egr1* and *Nr3c1*. In addition, "normalized" *Cacna1c* expression is consistent with the lack of anxiety and depression-related phenotype in the double-stress male group. However, MS males also did not show any significant behavioral phenotype, despite reduced *Cacna1c* expression, suggesting that these animals may have expressed some protective or pro-resilient factors that "buffered" the effect of *Cacna1c* deficiency.

Finally, *Dnmt1* is the only gene which showed the largest expression changes in the double-hit stress group. Being an epigenetic regulator, *Dnmt1* over-expression may affect the activity of many downstream target genes and, therefore, contribute to behavioral phenotypes. One possibility is that *Dnmt1* up-regulation contributed to the hyperactivity seen in MS\_SI group. This is consistent with the finding that the psychostimulant methamphetamine, which induces hyperactivity in rodents, also induces *Dnmt1* mRNA levels in the rodent brain (Numachi et al., 2007). In addition, an increased *Dnmt1* expression has been found in psychosis (Veldic et al., 2005) and DNA methylation changes have been reported in many neuropsychiatric disorders including schizophrenia (Grayson and Guidotti, 2013) and ADHD (Chen et al., 2018; Heinrich et al., 2017), further providing a possible link between *Dnmt1* dysregulation and the hyperactivity-related phenotype observed in our double-hit stress group.

In the last part of our study, we explored whether DNA methylation, an epigenetic mechanism, may contribute to the effects of developmental stressors on gene expression and behavioral outcomes. There are now a quite large number of studies suggesting that early-life experiences may exert long-term effects on brain and behavior via epigenetic mechanisms. Among those studies, changes in DNA methylation of the *Nr3c1* gene were the first to be linked to altered gene expression, HPA dysfunction, and anxiety-and depression-related phenotypes induced by early-life adversity in rats (Weaver et al., 2004) and humans (McGowan et al., 2009). Here we examined the region in mouse which is highly homologous (92%) to the previously reported exon 17 in rats (Weaver et al., 2004), and also containing the binding site for Egr1 (or NGFI-A) transcription factor encompassing two CpG sites (Kundakovic et al., 2013b; Weaver et al., 2004). Changes in DNA methylation within or adjacent to the transcription factor binding sites are particularly posed to affect transcriptional activity by preventing or facilitating the binding of transcription factors (Klose and Bird, 2006). While our results show complex group- and sex-specific patterns, it is notable that both sites in the Egr1 binding motif show female-specific reduction in methylation in the MS group. This finding is consistent with higher *Nr3c1* and *Egr1* expression in this group, and provides one of the possible mechanisms for the induced expression of the glucocorticoid gene and consequent behavioral alterations in early-life stress female group.

In terms of *Cacna1c* gene, there are reports of altered or more variable DNA methylation patterns in bipolar disorder (Starnawska et al., 2016) and depression (Cordova-Palomera et al., 2015) patients, respectively, compared to healthy subjects. We analyzed the region close to the transcriptional start site, with high homology between mice and humans, and shown to be differentially methylated between neurons and glia cells (Nishioka et al., 2013), hence, of possible functional relevance. We found subtle changes in DNA methylation of this gene in the double-stress hit group only, which cannot explain group- and sex-specific differences in *Cacna1c* gene expression that we have observed. However, it is possible that other regions of the *Cacna1c* gene, not examined in this study, may show differential methylation, which remains to be further explored. In addition, we would like to note that, although our DNA methylation analysis was limited to two regions and only 12 CpG sites total, our results suggest that epigenetic dysregulation may be prominent in the double-stress group, in line with changed gene expression of an epigenetic regulator, Dnmt1.

## Conclusions

In conclusion, our study shows complex interactions between early-life and adolescent stress, between stress and sex, and between stress and female estrogen status in shaping behavioral phenotypes of adult animals. In our paradigm, early-life stress had the most significant impact on anxiety- and depression-related phenotypes and this was female-specific. We discovered that early-life stress disrupts the protective role of estrogen in females, and promotes female vulnerability to anxiety- and depression-related phenotypes associated with the low-estrogenic state. We found transcriptional and epigenetic alterations in psychiatric risk genes, *Nr3c1* and *Cacna1c*, that likely contributed to the stress-induced behavioral effects. In addition, two general transcriptional regulators, Egr1 and Dnmt1, were found to be dysregulated in maternally-separated females and in animals exposed to both stressors, respectively, providing insights into possible transcriptional mechanisms that underlie behavioral phenotypes. We envision future transcriptomics and epigenomics studies that will give more definitive answers

regarding molecular signatures and mechanisms that drive developmental stress-induced vulnerability or resilience to anxiety and depression-related phenotypes. Our study highlights the importance of accounting for the effects of sex and female hormone status in the studies of the impact of stress on brain and behavior, and provides a foundation for the future studies of molecular mechanisms underlying sex disparity in anxiety and depression. This knowledge will be extremely important to enable the development of more effective, sex-specific treatments for these disorders.

**Funding:** This study was supported by the NARSAD Young Investigator Grant from the Brain & Behavior Research Foundation to M.K.

**Author Contributions:** IJ and MK designed the study; IJ, DR, AH, and MK performed experiments; HC performed statistical analysis; IJ, DR, HC, and MK interpreted the data; IJ, HC, and MK constructed the figures; DR and MK wrote the article; MK conceived and directed the project; all authors revised and approved the final version of the paper.

**Conflict of Interest Statement:** The authors declare no competing financial interests.

## References:

Altemus, M., Sarvaiya, N., and Neill Epperson, C. (2014). Sex differences in anxiety and depression clinical perspectives. *Frontiers in neuroendocrinology* 35, 320-330.

Angelucci, F., Brene, S., and Mathe, A.A. (2005). BDNF in schizophrenia, depression and corresponding animal models. *Molecular psychiatry* 10, 345-352.

Bale, T.L., Baram, T.Z., Brown, A.S., Goldstein, J.M., Insel, T.R., McCarthy, M.M., Nemeroff, C.B., Reyes, T.M., Simerly, R.B., and Susser, E.S. (2010). Early life programming and neurodevelopmental disorders. *Biological psychiatry* 68, 314-319.

Bannerman, D.M., Grubb, M., Deacon, R.M., Yee, B.K., Feldon, J., and Rawlins, J.N. (2003). Ventral hippocampal lesions affect anxiety but not spatial learning. *Behavioural brain research* 139, 197-213.

Beery, A.K., and Zucker, I. (2011). Sex bias in neuroscience and biomedical research. *Neuroscience & Biobehavioral Reviews* 35, 565-572.

Biggio, F., Pisu, M.G., Garau, A., Boero, G., Locci, V., Mostallino, M.C., Olla, P., Utzeri, C., and Serra, M. (2014). Maternal separation attenuates the effect of adolescent social isolation on HPA axis responsiveness in adult rats. *European neuropsychopharmacology* 24, 1152-1161.

Boersma, G.J., Lee, R.S., Cordner, Z.A., Ewald, E.R., Purcell, R.H., Moghadam, A.A., and Tamashiro, K.L. (2014). Prenatal stress decreases Bdnf expression and increases methylation of Bdnf exon IV in rats. *Epigenetics* 9, 437-447.

Boulle, F., van den Hove, D.L., Jakob, S.B., Rutten, B.P., Hamon, M., van Os, J., Lesch, K.P., Lanfumey, L., Steinbusch, H.W., and Kenis, G. (2012). Epigenetic regulation of the BDNF gene: implications for psychiatric disorders. *Molecular psychiatry* 17, 584-596.

Cembrowski, M.S., Wang, L., Sugino, K., Shields, B.C., and Spruston, N. (2016). Hipposeq: a comprehensive RNA-seq database of gene expression in hippocampal principal neurons. *eLife* 5, e14997.

Champagne, D.L., Bagot, R.C., van Hasselt, F., Ramakers, G., Meaney, M.J., de Kloet, E.R., Joels, M., and Krugers, H. (2008). Maternal care and hippocampal plasticity: evidence for experience-dependent structural plasticity, altered synaptic functioning, and differential responsiveness to glucocorticoids and stress. *The Journal of neuroscience* 28, 6037-6045.

Chen, Y.C., Sudre, G., Sharp, W., Donovan, F., Chandrasekharappa, S.C., Hansen, N., Elnitski, L., and Shaw, P. (2018). Neuroanatomic, epigenetic and genetic differences in monozygotic twins discordant for attention deficit hyperactivity disorder. *Molecular psychiatry* 23, 683-690.

Cordova-Palomera, A., Fatjo-Vilas, M., Gasto, C., Navarro, V., Krebs, M.O., and Fananas, L. (2015). Genome-wide methylation study on depression: differential methylation and variable methylation in monozygotic twins. *Translational psychiatry* 5, e557.

Czeh, B., Varga, Z.K., Henningsen, K., Kovacs, G.L., Miseta, A., and Wiborg, O. (2015). Chronic stress reduces the number of GABAergic interneurons in the adult rat hippocampus, dorsal-ventral and region-specific differences. *Hippocampus* *25*, 393-405.

Dao, D.T., Mahon, P.B., Cai, X., Kovacsics, C.E., Blackwell, R.A., Arad, M., Shi, J., Zandi, P.P., O'Donnell, P., Bipolar Genome Study, C., *et al.* (2010). Mood disorder susceptibility gene CACNA1C modifies mood-related behaviors in mice and interacts with sex to influence behavior in mice and diagnosis in humans. *Biol Psychiatry* *68*, 801-810.

de Sousa, R.T., Loch, A.A., Carvalho, A.F., Brunoni, A.R., Haddad, M.R., Henter, I.D., Zarate, C.A., and Machado-Vieira, R. (2017). Genetic Studies on the Tripartite Glutamate Synapse in the Pathophysiology and Therapeutics of Mood Disorders. *Neuropsychopharmacology* *42*, 787-800.

Dedic, N., Pohlmann, M.L., Richter, J.S., Mehta, D., Czamara, D., Metzger, M.W., Dine, J., Bedenk, B.T., Hartmann, J., Wagner, K.V., *et al.* (2018). Cross-disorder risk gene CACNA1C differentially modulates susceptibility to psychiatric disorders during development and adulthood. *Molecular psychiatry* *23*, 533-543.

Deecker, D., Andree, T.H., Sloan, D., and Schechter, L.E. (2008). From menarche to menopause: exploring the underlying biology of depression in women experiencing hormonal changes. *Psychoneuroendocrinology* *33*, 3-17.

Dong, E., Dzitoyeva, S.G., Matrisciano, F., Tueting, P., Grayson, D.R., and Guidotti, A. (2015). Brain-derived neurotrophic factor epigenetic modifications associated with schizophrenia-like phenotype induced by prenatal stress in mice. *Biol Psychiatry* *77*, 589-596.

Fanselow, M.S., and Dong, H.W. (2010). Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron* *65*, 7-19.

Ferreira, M.A., O'Donovan, M.C., Meng, Y.A., Jones, I.R., Ruderfer, D.M., Jones, L., Fan, J., Kirov, G., Perlis, R.H., Green, E.K., *et al.* (2008). Collaborative genome-wide association analysis supports a role for ANK3 and CACNA1C in bipolar disorder. *Nature genetics* *40*, 1056-1058.

Grayson, D.R., and Guidotti, A. (2013). The dynamics of DNA methylation in schizophrenia and related psychiatric disorders. *Neuropsychopharmacology* *38*, 138-166.

Green, E.K., Grozeva, D., Jones, I., Jones, L., Kirov, G., Caesar, S., Gordon-Smith, K., Fraser, C., Forty, L., and Russell, E. (2010). The bipolar disorder risk allele at CACNA1C also confers risk of recurrent major depression and of schizophrenia. *Molecular psychiatry* *15*, 1016.

Gutiérrez-Mecinas, M., Trollope, A.F., Collins, A., Morfett, H., Hesketh, S.A., Kersanté, F., and Reul, J.M. (2011). Long-lasting behavioral responses to stress involve a direct interaction of glucocorticoid receptors with ERK1/2-MSK1-Elk-1 signaling. *Proceedings of the National Academy of Sciences* *108*, 13806-13811.

Heim, C., and Nemeroff, C.B. (2001). The role of childhood trauma in the neurobiology of mood and anxiety disorders: preclinical and clinical studies. *Biol Psychiatry* *49*, 1023-1039.

Heinrich, H., Grunitz, J., Stonawski, V., Frey, S., Wahl, S., Albrecht, B., Goecke, T.W., Beckmann, M.W., Kornhuber, J., and Fasching, P.A. (2017). Attention, cognitive control and motivation in ADHD: Linking event-related brain potentials and DNA methylation patterns in boys at early school age. *Scientific reports* *7*, 3823.

Henley, J.M., and Wilkinson, K.A. (2016). Synaptic AMPA receptor composition in development, plasticity and disease. *Nature reviews. Neuroscience* *17*, 337-350.

Hsiao, Y.M., Tsai, T.C., Lin, Y.T., Chen, C.C., Huang, C.C., and Hsu, K.S. (2016). Early life stress dampens stress responsiveness in adolescence: Evaluation of neuroendocrine reactivity and coping behavior. *Psychoneuroendocrinology* *67*, 86-99.

Kheirbek, M.A., Drew, L.J., Burghardt, N.S., Costantini, D.O., Tannenholz, L., Ahmari, S.E., Zeng, H., Fenton, A.A., and Hen, R. (2013). Differential control of learning and anxiety along the dorsoventral axis of the dentate gyrus. *Neuron* *77*, 955-968.

Klose, R.J., and Bird, A.P. (2006). Genomic DNA methylation: the mark and its mediators. *Trends in biochemical sciences* *31*, 89-97.

Kornstein, S.G., Harvey, A.T., Rush, A.J., Wisniewski, S.R., Trivedi, M.H., Svikis, D.S., McKenzie, N.D., Bryan, C., and Harley, R. (2005). Self-reported premenstrual exacerbation of depressive symptoms in patients seeking treatment for major depression. *Psychological medicine* *35*, 683-692.

Kuehner, C. (2017). Why is depression more common among women than among men? *The Lancet Psychiatry* *4*, 146-158.

Kundakovic, M., and Champagne, F.A. (2015). Early-life experience, epigenetics, and the developing brain. *Neuropsychopharmacology* *40*, 141-153.

Kundakovic, M., Gudsnu, K., Franks, B., Madrid, J., Miller, R.L., Perera, F.P., and Champagne, F.A. (2013a). Sex-specific epigenetic disruption and behavioral changes following low-dose in utero bisphenol A exposure. *Proceedings of the National Academy of Sciences of the United States of America* *110*, 9956-9961.

Kundakovic, M., Gudsnu, K., Herbstman, J.B., Tang, D., Perera, F.P., and Champagne, F.A. (2015). DNA methylation of BDNF as a biomarker of early-life adversity. *Proceedings of the National Academy of Sciences of the United States of America* *112*, 6807-6813.

Kundakovic, M., Lim, S., Gudsnu, K., and Champagne, F.A. (2013b). Sex-specific and strain-dependent effects of early life adversity on behavioral and epigenetic outcomes. *Frontiers in psychiatry* *4*, 78.

Li, S.H., and Graham, B.M. (2017). Why are women so vulnerable to anxiety, trauma-related and stress-related disorders? The potential role of sex hormones. *The Lancet Psychiatry* *4*, 73-82.

Lupien, S.J., McEwen, B.S., Gunnar, M.R., and Heim, C. (2009). Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature reviews. Neuroscience* *10*, 434-445.

Marcondes, F.K., Miguel, K.J., Melo, L.L., and Spadari-Bratfisch, R.C. (2001). Estrous cycle influences the response of female rats in the elevated plus-maze test. *Physiol Behav* 74, 435-440.

McEwen, B.S. (2008). Understanding the potency of stressful early life experiences on brain and body function. *Metabolism: clinical and experimental* 57 *Suppl 2*, S11-15.

McGowan, P.O., Sasaki, A., D'Alessio, A.C., Dymov, S., Labonte, B., Szyf, M., Turecki, G., and Meaney, M.J. (2009). Epigenetic regulation of the glucocorticoid receptor in human brain associates with childhood abuse. *Nature neuroscience* 12, 342-348.

McLean, A.C., Valenzuela, N., Fai, S., and Bennett, S.A. (2012). Performing vaginal lavage, crystal violet staining, and vaginal cytological evaluation for mouse estrous cycle staging identification. *Journal of visualized experiments : JoVE*, e4389.

Millstein, R.A., and Holmes, A. (2007). Effects of repeated maternal separation on anxiety- and depression-related phenotypes in different mouse strains. *Neuroscience and biobehavioral reviews* 31, 3-17.

Morrison, M.F., Kallan, M.J., Ten Have, T., Katz, I., Tweedy, K., and Battistini, M. (2004). Lack of efficacy of estradiol for depression in postmenopausal women: a randomized, controlled trial. *Biol Psychiatry* 55, 406-412.

Murthy, S., and Gould, E. (2018). Early Life Stress in Rodents: Animal Models of Illness or Resilience? *Frontiers in behavioral neuroscience* 12, 157.

Nederhof, E. (2012). The mismatch hypothesis of psychiatric disease. *Physiol Behav* 106, 689-690.

Nishioka, M., Shimada, T., Bundo, M., Ukai, W., Hashimoto, E., Saito, T., Kano, Y., Sasaki, T., Kasai, K., Kato, T., *et al.* (2013). Neuronal cell-type specific DNA methylation patterns of the Cacna1c gene. *International journal of developmental neuroscience* 31, 89-95.

Niwa, M., Jaaro-Peled, H., Tankou, S., Seshadri, S., Hikida, T., Matsumoto, Y., Cascella, N.G., Kano, S., Ozaki, N., Nabeshima, T., *et al.* (2013). Adolescent stress-induced epigenetic control of dopaminergic neurons via glucocorticoids. *Science* 339, 335-339.

Numachi, Y., Shen, H., Yoshida, S., Fujiyama, K., Toda, S., Matsuoka, H., Sora, I., and Sato, M. (2007). Methamphetamine alters expression of DNA methyltransferase 1 mRNA in rat brain. *Neuroscience letters* 414, 213-217.

Onishchenko, N., Karpova, N., Sabri, F., Castren, E., and Ceccatelli, S. (2008). Long-lasting depression-like behavior and epigenetic changes of BDNF gene expression induced by perinatal exposure to methylmercury. *J Neurochem* 106, 1378-1387.

Pena, C.J., Kronman, H.G., Walker, D.M., Cates, H.M., Bagot, R.C., Purushothaman, I., Issler, O., Loh, Y.E., Leong, T., Kiraly, D.D., *et al.* (2017). Early life stress confers lifelong stress susceptibility in mice via ventral tegmental area OTX2. *Science* 356, 1185-1188.

Pinto, V., Costa, J.C., Morgado, P., Mota, C., Miranda, A., Bravo, F.V., Oliveira, T.G., Cerqueira, J.J., and Sousa, N. (2015). Differential impact of chronic stress along the hippocampal dorsal-ventral axis. *Brain structure & function* *220*, 1205-1212.

Romeo, R.D., and McEwen, B.S. (2006). Stress and the adolescent brain. *Annals of the New York Academy of Sciences* *1094*, 202-214.

Roth, T.L., Lubin, F.D., Funk, A.J., and Sweatt, J.D. (2009). Lasting epigenetic influence of early-life adversity on the BDNF gene. *Biol Psychiatry* *65*, 760-769.

Rusconi, F., Grillo, B., Ponzoni, L., Bassani, S., Toffolo, E., Paganini, L., Mallei, A., Braida, D., Passafaro, M., Popoli, M., *et al.* (2016). LSD1 modulates stress-evoked transcription of immediate early genes and emotional behavior. *Proceedings of the National Academy of Sciences of the United States of America* *113*, 3651-3656.

Santarelli, S., Zimmermann, C., Kalideris, G., Lesuis, S.L., Arloth, J., Uribe, A., Dournes, C., Balsevich, G., Hartmann, J., and Masana, M. (2017). An adverse early life environment can enhance stress resilience in adulthood. *Psychoneuroendocrinology* *78*, 213-221.

Schmidt, M.V. (2011). Animal models for depression and the mismatch hypothesis of disease. *Psychoneuroendocrinology* *36*, 330-338.

Sklar, P., Smoller, J.W., Fan, J., Ferreira, M.A., Perlis, R.H., Chambert, K., Nimgaonkar, V.L., McQueen, M.B., Faraone, S.V., Kirby, A., *et al.* (2008). Whole-genome association study of bipolar disorder. *Molecular psychiatry* *13*, 558-569.

Starnawska, A., Demontis, D., Pen, A., Hedemand, A., Nielsen, A., Staunstrup, N., Grove, J., Als, T., Jarram, A., and O'brien, N. (2016). CACNA1C hypermethylation is associated with bipolar disorder. *Translational psychiatry* *6*, e831.

Steiner, M., Dunn, E., and Born, L. (2003). Hormones and mood: from menarche to menopause and beyond. *Journal of affective disorders* *74*, 67-83.

Suderman, M., McGowan, P.O., Sasaki, A., Huang, T.C., Hallett, M.T., Meaney, M.J., Turecki, G., and Szyf, M. (2012). Conserved epigenetic sensitivity to early life experience in the rat and human hippocampus. *Proceedings of the National Academy of Sciences of the United States of America* *109 Suppl 2*, 17266-17272.

Takeuchi, H., Tomita, H., Taki, Y., Kikuchi, Y., Ono, C., Yu, Z., Nouchi, R., Yokoyama, R., Kotozaki, Y., and Nakagawa, S. (2018). A Common CACNA1C Gene Risk Variant has Sex-Dependent Effects on Behavioral Traits and Brain Functional Activity. *Cerebral Cortex*.

Veldic, M., Guidotti, A., Maloku, E., Davis, J.M., and Costa, E. (2005). In psychosis, cortical interneurons overexpress DNA-methyltransferase 1. *Proceedings of the National Academy of Sciences of the United States of America* *102*, 2152-2157.

Walf, A.A., and Frye, C.A. (2006). A review and update of mechanisms of estrogen in the hippocampus and amygdala for anxiety and depression behavior. *Neuropsychopharmacology* *31*, 1097-1111.

Weaver, I.C., Cervoni, N., Champagne, F.A., D'Alessio, A.C., Sharma, S., Seckl, J.R., Dymov, S., Szyf, M., and Meaney, M.J. (2004). Epigenetic programming by maternal behavior. *Nature neuroscience* 7, 847-854.

Will, T.R., Proaño, S.B., Thomas, A.M., Kunz, L.M., Thompson, K.C., Ginnari, L.A., Jones, C.H., Lucas, S.-C., Reavis, E.M., and Dorris, D.M. (2017). Problems and Progress regarding Sex Bias and Omission in Neuroscience Research. *eNeuro* 4, ENEURO.0278-0217.2017.

Witt, S.H., Kleindienst, N., Frank, J., Treutlein, J., Muhleisen, T., Degenhardt, F., Jungkunz, M., Krumm, B., Cichon, S., Tadic, A., *et al.* (2014). Analysis of genome-wide significant bipolar disorder genes in borderline personality disorder. *Psychiatric genetics* 24, 262-265.

Xie, L., Korkmaz, K.S., Braun, K., and Bock, J. (2013). Early life stress-induced histone acetylations correlate with activation of the synaptic plasticity genes Arc and Egr1 in the mouse hippocampus. *J Neurochem* 125, 457-464.

## Figure Legends:

**Figure 1. Study Design:** two stress hits included early-life maternal separation (MS, PD1-14) and adolescent social isolation (SI, PD35-56). This design generated four groups of animals: no stress controls (CON), MS only, SI only, and combined two-hit stress MS\_SI group. The majority of animals underwent behavioral testing from PD57-68 and their non-tested littermates were sacrificed for molecular analyses of the ventral hippocampus at PD70.

**Figure 2. The effect of maternal separation and social isolation on body weight.** Body weight was examined after weaning (A), prior to social isolation (B), and before behavioral testing (C). Note that PD28 and PD35 show the effects of MS only and the groups are merged to represent animals that did not undergo MS (CON & SI groups) vs. animals that underwent MS (MS & MS\_SI groups). At PD56, social isolation was completed and four groups of animals are shown separately reflecting the effects of both MS and SI hits (N=15-23/sex/group). Symbols in the upper left corner of each graph show significant main effects of group and sex or their interaction. Significant results of posthoc tests are presented in detail. CON, control; MS, maternal separation; SI, social isolation; MS\_SI, combined maternal separation and social isolation; \*, P < 0.05; \*\*, P < 0.01; \*\*\*, P < 0.001.

**Figure 3. Sex-specific and estrous cycle-dependent effects of maternal separation and social isolation on the open field behavior.** Total distance traveled (A, B), latency to the first center entry (C, D) and time spent in the center (E, F) were analyzed, either in all animals with group and sex as factors (upper panels - A, C, E) or in females with group and estrogen status as factors (lower panels - B, D, F); (N=15-23/sex/group). Symbols in the upper left corner of each graph show significant main effects of group and sex/estrogen or their interaction. Significant results of posthoc tests are presented in detail. CON, control; MS, maternal separation; SI, social isolation; MS\_SI, combined maternal separation and social isolation; \*, P < 0.05; \*\*, P < 0.01; \*\*\*, P < 0.001.

**Figure 4. Sex- specific effects of maternal separation and social isolation in the elevated plus maze test.** Total distance traveled (A, B), latency to the first open arm entry (C, D) and time spent in the open arms (E, F) were analyzed, either in all animals with group and sex as factors (upper panels - A, C, E) or in females with group and estrogen status as factors (lower panels - B, D, F); (N=15-23/sex/group). Symbols in the upper left corner of each graph show significant main effects of group and sex/estrogen or their interaction. Significant results of posthoc tests are presented in detail. CON, control; MS, maternal separation; SI, social isolation; MS\_SI, combined maternal separation and social isolation; \*, P < 0.05; \*\*, P < 0.01; \*\*\*, P < 0.001.

**Figure 5. Sex-specific effects of maternal separation and social isolation on depression-related phenotypes.** Results of sucrose preference (A) and the forced swim (B-C) tests are shown. Sucrose preference was analyzed in all animals with group and sex as factors (A). For the forced swim test, time spent immobile was analyzed in all animals with group and sex as factors (B) and in females with group and estrogen status as factors (C); (N=15-23/sex/group). Symbols in the upper left corner of each graph show significant main effects of group and sex/estrogen or their interaction. Significant results of posthoc tests are presented in detail. CON, control; MS,

maternal separation; SI, social isolation; MS\_SI, combined maternal separation and social isolation; \*, P < 0.05; \*\*, P < 0.01; \*\*\*, P < 0.001.

**Figure 6. Sex-specific effects of maternal separation and social isolation on gene expression in the ventral hippocampus.** The expression of *Bdnf* (A), *Nr3c1* (B), *Egr1* (C), *Cacna1c* (D), *Gria3* (E), and *Dnmt1* (F) was analyzed in the ventral hippocampus, using group and sex as factors (N=12 females/group, N=8 males/group). Symbols in the upper left corner of each graph show significant main effects of group and sex or their interaction. Significant results of posthoc tests are presented in detail. CON, control; MS, maternal separation; SI, social isolation; MS\_SI, combined maternal separation and social isolation; \*, P < 0.05; \*\*, P < 0.01; \*\*\*, P < 0.001.

**Figure 7. Sex-specific effects of maternal separation and social isolation on DNA methylation of the *Nr3c1* and *Cacna1c* regulatory regions.** **A.** Schematic representation of the *Nr3c1* gene, the sequence of the analyzed CpG sites in the *Nr3c1* 5' regulatory region, and methylation levels of individual sites, analyzed using group and sex as factors. Underlined sequence represents *Egr1* binding site. **B.** Schematic representation of the *Cacna1c* gene, the sequence of the analyzed CpG sites in the *Cacna1c* regulatory region, and methylation levels of individual sites, analyzed using group and sex as factors (N=6 sex/group). (Note: both genes have alternative first exons which are shown as dashed rectangles). Symbols in the upper left corner of each graph show significant main effects of group and sex or their interaction. Significant results of posthoc tests are presented in detail. CON, control; MS, maternal separation; SI, social isolation; MS\_SI, combined maternal separation and social isolation; \*, P < 0.05; \*\*, P < 0.01; \*\*\*, P < 0.001.

Figure 1





Figure 2

● Group ● Sex ●● Group x Sex


A

PD28



B

PD35



C

PD56




Figure 3

● Group ● Sex ●● Group x Sex ●● Estrogen ●●● Group x Estrogen




Figure 4

● Group ● Sex ●● Group x Sex ●● Estrogen ●●● Group x Estrogen




Figure 5

● Group ● Sex ●● Group x Sex ●● Estrogen ●●● Group x Estrogen




Figure 6

● Group ● Sex ● Group x Sex





Figure 7


● Group ● Sex ●● Group x Sex

A

Mouse *Nr3c1* gene



bioRxiv preprint doi: <https://doi.org/10.1101/530311>; this version posted January 25, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.



B

Mouse *Cacna1c* gene

