

# **Opposing signaling pathways regulate morphology in response to temperature in the fungal pathogen *Histoplasma capsulatum***

Lauren Rodriguez, Mark Voorhies, Sarah Gilmore\*, Sinem Beyhan\*\*, Anthony Myint#, Anita

Sil

Department of Microbiology and Immunology, University of California San Francisco, San Francisco, CA 94143-0414

\*Current address: Gilead Sciences, 333 Lakeside Drive, Foster City, CA 94404

\*\*Current address: J. Craig Venter Institute, 4120 Capricorn Lane, La Jolla, CA 92047

#Current address: Los Angeles, CA

Corresponding author

Anita Sil, Department of Microbiology and Immunology, University of California San Francisco, San Francisco, CA 94143-0414

Phone: (415) 502-1805

Fax: (415) 476-8201

Email: [sil@cgl.ucsf.edu](mailto:sil@cgl.ucsf.edu)

## ABSTRACT

Phenotypic switching between two opposing cellular states is a fundamental aspect of biology, and fungi provide facile systems to analyze the interactions between regulons that control this type of switch. A long-standing mystery in fungal pathogens of humans is how thermally dimorphic fungi switch their developmental form in response to temperature. These fungi, including the subject of this study, *Histoplasma capsulatum*, are temperature-responsive organisms that utilize unknown regulatory pathways to couple their cell shape and associated attributes to the temperature of their environment. *H. capsulatum* grows as a multicellular hypha in the soil that switches to a pathogenic yeast form in response to the temperature of a mammalian host. These states can be triggered in the laboratory simply by growing the fungus either at room temperature (where it grows as hyphae) or at 37°C (where it grows as yeast). Prior work revealed that 15-20% of transcripts are differentially expressed in response to temperature, but it is unclear which transcripts are linked to specific phenotypic changes such as cell morphology or virulence. To elucidate temperature-responsive regulons, we previously identified four transcription factors (Ryp1-4) that are required for yeast-phase growth at 37°C; in each *ryp* mutant, the fungus grows constitutively as hyphae regardless of temperature and the cells fail to express genes that are normally induced in response to growth at 37°C. Here we perform the first genetic screen to identify genes required for hyphal growth of *H. capsulatum* at room temperature and find that disruption of the signaling mucin *MSB2* results in a yeast-locked phenotype. RNAseq experiments reveal that *MSB2* is not required for the majority of gene expression changes that occur when cells are shifted to room temperature. However, a small subset of temperature-responsive genes is dependent on *MSB2* for its expression, thereby

implicating these genes in the process of filamentation. Disruption or knockdown of an Msb2-dependent MAP kinase (*HOG2*) and an APSES transcription factor (*STU1*) prevents hyphal growth at room temperature, validating that the Msb2 regulon contains genes that control filamentation. Notably, the Msb2 regulon shows conserved hyphal-specific expression in other dimorphic fungi, suggesting that this work defines a small set of genes that are likely to be conserved regulators and effectors of filamentation in multiple fungi. In contrast, a few yeast-specific transcripts, including virulence factors that are normally expressed only at 37°C, are inappropriately expressed at room temperature in the *msb2* mutant, suggesting that expression of these genes is coupled to growth in the yeast form rather than to temperature. Finally, we find that the yeast-promoting transcription factor Ryp3 associates with the *MSB2* promoter and inhibits *MSB2* transcript expression at 37°C, whereas Msb2 inhibits accumulation of Ryp transcripts and proteins at room temperature. These findings indicate that the Ryp and Msb2 circuits antagonize each other in a temperature-dependent manner, thereby allowing temperature to govern cell shape and gene expression in this ubiquitous fungal pathogen of humans.

## Introduction

The ability of microbes to sense and respond to the environment is key for survival in a variety of niches. Thermally dimorphic fungi make a dichotomous developmental choice depending on their environment. In the soil, they grow in a multicellular, hyphal form (also referred to here as the filamentous form) that also produces vegetative spores. When the soil is disturbed, hyphal fragments and associated spores can be inhaled by mammalian hosts. The body temperature of the host is a sufficient signal to switch the growth pattern of the fungus to a specialized host form (unicellular yeasts in the case of *Histoplasma capsulatum*) that expresses

virulence factors associated with survival in the host and pathogenesis. In the laboratory, *H. capsulatum* grows in a hyphal form at room temperature (RT) and switches to a yeast form at 37°C, providing an excellent system to study the regulation of developmental switches that are relevant to disease progression during infection.

*H. capsulatum* is a fungus endemic to the Ohio and Mississippi River valleys where 60-80% of inhabitants test positive for exposure during their lifetime [1]. The transition to the yeast form is thought to facilitate the lifestyle of the fungus in the host, where it replicates within macrophages and causes disseminated disease. Although *H. capsulatum* is ubiquitous, little is known about the molecular regulators and effectors that enable the bidirectional switch between the infectious soil form and the pathogenic host form. A number of studies [2-7] have shown that a significant fraction of the genome is differentially expressed between *H. capsulatum* hyphae grown at RT and *H. capsulatum* yeasts grown at 37°C, including known virulence factors that are specifically expressed at mammalian body temperature. However, little is known about the role of the majority of differentially expressed transcripts in either the biology of the organism or the switch between developmental forms.

In previous work, to elucidate how such a large morphological and transcriptional restructuring occurs in response to temperature, we utilized forward genetic screens to identify four transcription factors, *RYPI-4*, that are required for yeast-phase growth. Mutation of any of the *RYP* genes results in constitutive hyphal growth independent of temperature. Additionally, each of these transcription factors is required for the normal transcriptional response to temperature [2, 7, 8]: the transcriptional profile of the *ryp* mutants at 37°C looks drastically different from wild-type yeast cells but very similar to wild-type hyphae grown at RT [2]. We also found that Ryp transcripts and proteins accumulate to higher levels at 37°C than at RT [2, 4,

7, 8], which presumably contributes to their ability to trigger a temperature-dependent program of gene expression.

Here we elucidate pathways that function at RT to antagonize the Ryp regulatory circuit and prevent yeast-phase growth. We perform a genetic screen for yeast-locked mutants and identify the signaling mucin *MSB2* as required for hyphal formation at RT. Orthologs of Msb2 have been studied in *Saccharomyces cerevisiae* and *Candida albicans* where this transmembrane protein stimulates a number of signaling pathways, including the high osmolarity glycerol (HOG) pathway in response to osmotic stress, as well as the filamentous growth pathway in response to nutrient limitation [9-15]. We examine the role of *MSB2* in *H. capsulatum* as an environmental sensor that controls the morphologic response to RT growth. We find that *MSB2* is required for the expression of a small set of temperature-responsive genes, and that this regulon shows conserved expression in the filamentous form of other dimorphic fungi. To validate that this *MSB2* regulon contains genes that promote hyphal growth, we disrupt both a MAP kinase and APSES transcription factor whose transcriptional induction at RT is dependent on *MSB2* and show that the resultant mutants are defective for normal hyphal growth in response to temperature. Additionally, we provide the first molecular evidence that the Ryp circuit and the Msb2 circuit are mutually antagonistic in a temperature-dependent manner, resulting in the ability of temperature to control cell shape and associated characteristics.

## Results

### The *MSB2* gene is required for hyphal growth at RT in *Histoplasma capsulatum*

Previous studies have identified and characterized genes required for the yeast phase of *H. capsulatum* and other thermally dimorphic fungi [2, 7, 8, 16]. These studies have shed light

on the regulation of yeast phase growth; however the regulation of filamentation remains largely unclear. Similarly, how RT growth inhibits the yeast program is also not understood. We sought to shed light on these mechanisms by identifying mutants that inappropriately maintain the yeast state and fail to form hyphae at RT. We used *Agrobacterium*-mediated insertional mutagenesis and screened for mutants that retained their yeast morphology after growth for 6 weeks at RT. One of the mutants we obtained failed to transition to hyphal growth after a shift to RT, which normally induces filamentation (Fig. 1A). Instead, the mutant continued to divide as yeast (Fig. S1). Whole-genome sequencing of the mutant led to the identification of the *Agrobacterium* T-DNA insertion site in the promoter region of the *MSB2* gene (Fig. 1B). Introduction of a wild-type copy of the *MSB2* gene on an episomal plasmid (hereafter referred to as “*msb2* + *MSB2*” or the “complemented strain”) restored the ability of the mutant to form hyphae at RT (Fig. 1C). Knockdown of *MSB2* by RNA interference (RNAi) in a wild-type background recapitulated the yeast-locked phenotype during growth at RT (Fig. 1D). Overexpression of *MSB2* at 37°C resulted in disruption of the normal yeast morphology and the appearance of polarized projections reminiscent of hyphae (Fig. 1E). Taken together, these data indicate that *MSB2* is necessary and sufficient to trigger filamentous growth.

**Fig. 1. *MSB2* is necessary and sufficient for filamentation of *H. capsulatum* in response to temperature.**

(A) Cell morphology of wild-type *H. capsulatum* and mutant after 4 days at RT or 37°C in HMM + GlcNAc media. (B) Genome sequencing alignment indicating the transposon insertion site upstream of the *MSB2* gene. This alteration results in transcript initiation in the T-DNA insertion (purple box), thereby introducing an upstream ORF (pink box). Thus the resultant

*MSB2* transcript is non-functional because it does not yield a normal translation product. (C) Cell morphology of wild-type *H. capsulatum*, *msb2*, and *MSB2* complementation strain shown after 8 days at RT. (D) *MSB2* RNAi strain cell morphology is shown after 6 week incubation at RT and the control strain after 4 days at RT. qRT-PCR analysis showed significant depletion of *MSB2* transcript in the RNAi knockdown strain compared to the vector control ( $P < 0.001$ ). (E) *MSB2* overexpression was driven by the *GAPDH* promoter on an ectopic plasmid. qRT-PCR analysis of *MSB2* expression level showed a significant increase in the overexpression (OE) strain compared to control at 37°C ( $P < 0.01$ ). Cells are shown after growth at 37°C for 12 days. (F) Wild-type, *msb2*, and *msb2+MSB2* were grown on HMM solid media +/- 1M sorbitol to observe any osmotic stress response that is reflected in the colony morphology at 37°C. HMM colonies were imaged at day 8 after plating and HMM +1M Sorbitol were imaged at day 17.

The function of *Msb2* in *H. capsulatum* was previously unknown, though it has orthologs in the distant fungal species *Candida albicans* [10], and *Saccharomyces cerevisiae*, where its function has been more thoroughly examined [9, 11-15]. The *S. cerevisiae MSB2* ortholog acts as a sensor in the osmotic stress response pathway. The response to high osmolarity in *H. capsulatum* has not been thoroughly characterized. To determine if *H. capsulatum MSB2* plays a role in responding to high osmolarity, we plated serial dilutions of wild-type, *msb2*, and the complemented strain on solid HMM media with 1M sorbitol (Fig. 1F). Interestingly, we observed that the wild-type and complemented strains exhibited a wrinkly colony morphology in the presence of sorbitol at 37°C, consistent with some degree of filamentous cells in the colony. In contrast, the *msb2* mutant gave rise to smooth colonies on high sorbitol, correlating with

yeast-form cells. These data suggest that the morphology of *H. capsulatum* changes in response to high osmolarity and that this altered cell shape is dependent on Msb2.

# **The transcriptional changes elicited by RT growth are largely independent of *MSB2***

Cells lacking *MSB2* are unable to form hyphae at RT. To understand the contribution of *MSB2* to the myriad transcriptional changes that accompany RT growth, we surveyed the transcriptome of wild-type, *msb2*, and complemented strains as they were shifted from 37°C to RT. Yeast cultures were grown in normal yeast-promoting conditions (37°C, 5% CO<sub>2</sub>) and transferred to RT without CO<sub>2</sub> to promote filamentation. This timecourse was performed in two different carbon sources, glucose (Fig.S2) and N-Acetylglucosamine (GlcNAc) (Fig.2A). We chose GlcNAc because we previously showed that it expedites the transition from yeast to hyphae at RT [17].

Wildtype and the complemented strain produced filaments after 2 days at RT in GlcNAc, but the *msb2* mutant remained in the yeast form over the 8-day timecourse. RNA was isolated from wild-type, *msb2*, and the complemented strains, and we performed RNAseq analysis at day zero (37°C) and 2, 6, and 8 days after the transition to RT. Fig. 2B is a scatter plot showing transcript abundances between two biological replicates of wild-type grown at 37°C, and as expected there is a high Pearson correlation between the samples ( $r = 0.99$ ) and a corresponding low average deviation from the line of best fit ( $rmsd = 0.36$ ). In contrast, wild-type hyphae harvested eight days after transfer to RT had an expression profile that was quite distinct from wild-type yeast at 37°C (Fig.2C,  $r = 0.86$ ,  $rmsd = 1.08$ ), consistent with the previously observed differences between mature yeast and hyphal cultures [2-7]. Surprisingly, the transcriptional program of the *msb2* mutant, which remained locked in the yeast form even at 8 days post-shift, looked very similar to that of wild-type hyphae (Fig. 2E,  $r = 0.97$ ,  $rmsd = 0.57$ ), and very different from wild-type 37°C yeast cultures (Fig. 2D,  $r = 0.86$ ,  $rmsd = 1.12$ ) in spite of their common morphology.



We conclude from these data that the majority of the temperature-dependent transcriptional program is independent of *MSB2*. Since the *msb2* mutant is yeast-locked, the small set of Msb2-dependent genes may be linked to the ability of *H. capsulatum* to form hyphae.

**Fig. 2. The majority of the temperature-dependent changes in gene expression is independent of Msb2.**

(A) Experimental setup for RNAseq timecourse. Cells shown were grown at 37°C with 5% CO<sub>2</sub> in HMM media with N-acetylglucosamine (GlcNAc) as the carbon source. Once shifted to RT, cells were collected at day 2, 6, and 8 post temperature shift. Images were taken at each time point to compare cell morphology of wild-type, *msb2*, and complemented (*msb2*+*MSB2*) strains. (B-E) Scatter plots of relative transcript abundances as log<sub>2</sub>(cpm). (B) Wild-type (WT) 37°C replicate 1 (day 0) vs replicate 2 (day 0). (C) WT 37°C (day 0) vs WT RT (day 8). (D) WT 37°C (day 0) vs *msb2* RT (day 8). (E) WT RT (day 8) vs *msb2* RT (day 8). The morphology of each sample is depicted with a cartoon schematic of yeast or hyphae.

**Transcriptional profiling identified a small set of Msb2-dependent genes with varying functions**

As described above, to assess the role of Msb2 in the transcriptional response to temperature, we examined the transcriptional profile of wild-type, mutant, and complemented strains before and after shifting cultures from 37°C to RT in two different media. Liquid cultures were split at t = 0 to give biological replicates, and the entire experiment was carried out on two distinct occasions (batches). Genotype and conditions for each profiled sample are given in S1

Data, with biological replicates collected both at the same time and on different days. There was some variability based on the particular batch, and the transcriptional program in GlcNAc vs. glucose was similar but not identical, with differences that were largely independent of the temperature-dependent changes. To extract the temperature and *MSB2* dependence of each gene from our full data set, we fit a linear model with independent terms for each time point and genotype with additional terms to control for effects due to media (GlcNAc vs. glucose) or batch. Estimated read counts and fit terms are given in S2 Data. We considered genes significantly differential with respect to a fit term if they had at least a two fold change and a p-value less than 0.05 after multiple hypothesis correction. Normalized expression levels of significantly differential genes are given in S3 Data and rendered as a heatmap in Fig. 3B. Consistent with the single sample comparisons (Fig. 2B-E), over a quarter of the transcriptome had significant differential expression over the transition (1870 genes up (11 + 1694 + 165; Fig. 3A) and 1115 genes down (40 + 1067 + 8; Fig. 3A) when comparing the wild-type yeast-phase transcriptome at day 0 (37°C) vs. the wild-type hyphal transcriptome at day 8 (RT)). In contrast, when comparing wild-type day 8 with the *msb2* mutant at day 8, only 3% of the transcriptome was significantly differential (93 genes up in the *msb2* mutant compared to wild-type (11 + 42 + 40; Fig. 3A) and 249 genes down in the *msb2* mutant compared to wild-type (165 + 76 + 8; Fig. 3A), again meaning that the vast majority of transcriptional changes upon shift to room temperature are intact in the *msb2* mutant. For the small subset of genes whose expression is dependent on *Msb2*, we observed some correlation between morphology and *Msb2*-dependence, with either (1) increased expression in the yeast-locked *msb2* mutant vs. wild-type hyphae *and* increased expression in wild-type yeast cells vs. wild-type filaments (40 genes; Fig. 3A) or (2) decreased expression in the yeast-locked *msb2* mutant vs. wild-type hyphae *and* decreased expression in

wild-type yeast cells vs. wild-type filaments (165 genes; Fig. 3A). This pattern was more common than transcripts that were up in the yeast-locked *msb2* mutant vs. wild-type hyphae and up in wild-type hyphae vs. wild-type yeast (11 genes; Fig. 3A) or transcripts that were up in wild-type hyphae vs. wild-type yeast and up in yeast-locked *msb2* mutant vs. wild-type hyphae (8 genes; Fig. 3A). This observation is consistent with the hypothesis that genes correlated with (and perhaps required for) hyphal morphology should be upregulated in wild-type hyphae over wild-type yeast as well as upregulated in wild-type hyphae over the yeast-locked *msb2* mutant. Similarly, genes that correlate with yeast morphology should have the opposite expression pattern. Therefore, we further inspected these jointly regulated "filament-associated" (165 genes; Fig. 3A) and "yeast-associated" (40 genes; Fig. 3A) gene sets for potential morphological regulators, noting that a subset of these genes could control morphology and others could be effectors associated with a particular morphologic state.

### **Fig. 3. Classification of transcripts by dependence on *MSB2* and temperature.**

(A) Scatter plot of limma-fit parameters (as described in the Methods) for genotype (WT or *msb2* mutant) vs. time (day 0 or day 8). Transcripts significantly differential (at FDR = 5%) with at least a 2-fold change on one of these parameters are shown as larger, colored circles. Colored numbers indicate the number of transcripts in each class. (B) Heatmap of transcripts passing the significance and fold change criteria in A. Transcripts are grouped by the classification scheme of A, as indicated by the colored bar to the right of the heatmap. (C) Expanded view of the "filament-associated" class (165 genes) from 3A and B. (D) Expanded view of the "yeast-associated" class (40 genes) from 3A and B. Asterisks indicate putative knottin transcripts described in the text.

Analysis of the “filament-associated” gene set (Fig. 3C, S4 Data) revealed a number of orthologs of developmental regulators in other fungi. Among these was *FBA1*, which is a hyphal-specific gene that is upregulated in wild-type *H. capsulatum* at RT but not in *msb2*. It has an ortholog (*flbA*) in *Aspergillus sp.* whose function has been elucidated as a regulator of conidiation [18]. While the cell morphologies of *Aspergillus spp* are distinct from those of *H. capsulatum* (for example, *Aspergillus spp* lack a yeast form) this result may suggest a central role for this gene as a regulator of morphological development in a wide range of fungi. Other potential regulators in the Msb2-dependent filament-associated gene set include the mitogen-activated protein kinase *HOG2* and the APSES transcription factor *STUI*; we assessed the role of these regulators in filamentation as described below.

Other genes in the filament-associated set have been flagged previously as showing hyphal-enriched expression in *H. capsulatum*. *MS8* and *MS95* have been shown to be highly hyphal-enriched in *H. capsulatum* in previous datasets through transcriptional profiling and ribosomal footprint analysis [4, 19, 20]. Here we observed that *MS8* and *MS95* are upregulated as early as 2 days at RT in a Msb2-dependent manner. Thus their expression is reliably correlated with growth at RT, but their molecular function and how it pertains to hyphal growth remains unclear. Notably, disruption of *MS8* has been reported to result in abnormally shaped hyphal cells [20]. Similarly, *TYRI*, which encodes a putative tyrosinase, has been observed as a highly abundant hyphal-enriched transcript in multiple studies [2-7]. Our data indicate that the transcriptional induction of *TYRI* after growth at room temperature is dependent on Msb2.

Of the 40 yeast-associated genes (Fig. 3D, S5 Data), 16 (40%) were previously identified as targets of the Ryp transcription factors, meaning that their promoters are associated with

Ryp1, 2, 3, and/or 4. This is a significant enrichment over chance ( $p = 1.5e-7$ ). Strikingly, all 16 are Ryp1-associated, with Ryp2, Ryp3, and Ryp4 associating with different subsets of these promoters. Since the Ryp program drives yeast-phase growth, and the *msb2* mutant is yeast-locked, these data suggest the hypothesis that the Ryp pathway is inappropriately active in the *msb2* mutant at room temperature. We previously observed that many known virulence factors are Ryp-associated [2]; consistent with this observation, the 16 Ryp-associated yeast-specific genes include 2 virulence factors, *CBP1* [21-24] and *CTR3* [25], as well as two genes required for efficient lysis of macrophages, *LDF1* and a gene previously designated as UA35-G3 [26]. Several others are putative secreted factors of unknown function, including the paralogs Yps21 [27] and GH17/Cfp4 [28, 29] as well as ucsf.hc\_01.G217B.05476 which, along with UA35-G3, is predicted to be a secreted knottin (cystine knot proteins that show yeast-specific expression and may play a role in virulence in *H. capsulatum*) [4]. Notably, these yeast-associated genes, which are normally expressed by wild-type yeast cells at 37°C, are expressed in the yeast-locked *msb2* mutant even at room temperature, suggesting that their expression can be unlinked to temperature but remains linked to cell morphology.

# **The MAP kinase *HOG2* is downstream of *MSB2* and is involved in the temperature-dependent regulation of filamentation**

The function of Msb2 has been studied in a number of fungi [9-15, 30, 31]. These studies have supported the role of Msb2 as a transmembrane signaling molecule and more specifically, as one of the upstream signaling proteins in the *S. cerevisiae* high osmolarity glycerol (HOG) pathway [15, 32-34]. This pathway signals through the mitogen-activated protein kinase (MAPK) *HOG1* [35], which undergoes phosphorylation and translocation to the nucleus where it activates a transcriptional response to produce glycerol and restore osmotic balance. Given this

precedent, we were interested in the possibility that Msb2 signals through a MAPK in *H. capsulatum*. Using phylogenetic analysis, we identified four MAPKs in *H. capsulatum* (Fig. 4A) and named them, in three cases, for their orthologs in other fungi (*HMK1*, *HOG1*, *SLT2*). The fourth MAPK is a paralog of Hog1 that we named Hog2. To determine if any of these were induced in hyphal cells, we examined the gene expression of these four MAPKs in our transcriptional profiling dataset. Whereas *SLT2*, *HMK1*, and *HOG1* did not show transcriptional induction in wild-type cells undergoing filamentation, *HOG2* was markedly induced in wild-type cells in an Msb2-dependent fashion (Fig. 4B). Additionally, *HOG2* was classified as a filament-associated transcript (Fig. 3C), and thus is a good candidate for a regulator of hyphal growth. To test whether *HOG2* is required for filamentation, we constructed a *HOG2* RNAi strain and assessed the morphology of these cells at RT. Isolates that showed efficient knockdown were yeast-locked, indicating that *HOG2* is required for filamentation at RT (Fig. 4C). Additionally, in the *MSB2* overexpression strain, which exhibits disruption of yeast morphology and filamentation at 37°C (Fig. 1E), we observed a significant increase in *HOG2* expression level compared to the control strain (Fig. 4D). Taken together, these data indicate that *HOG2* expression is increased in an Msb2-dependent manner and that *HOG2*, like *MSB2*, is necessary for filamentation in response to temperature.

**Fig. 4. *HOG2* is a MAP kinase involved in filamentation in *H. capsulatum*.**

(A) FastTree maximum likelihood phylogeny of the 4 MAP kinases of *H. capsulatum* (shown in red) and their relation to other fungal MAP kinases. (B) Heatmap of *HMK2*, *HOG2*, *HOG1*, and *SLT2* data extracted from the RNAseq dataset. Values are log<sub>2</sub> ratios from single GlcNAc cultures of WT or *msb2* at multiple time points after shift to RT, relative to a WT 37°C sample.

(C-D) qRT-PCR data of *HOG2* transcript levels. (C) *HOG2* transcript levels in *HOG2* RNAi strain (P<0.01). Cell morphology of control and *HOG2* knockdown strains shown after 8 days at RT. (D) *HOG2* transcript levels in *MSB2* overexpression (OE) strain grown for 10 days at 37°C (P<0.05).

## **The APSES transcription factor *STUI* is necessary and sufficient for filamentation**

The *Msb2*-dependent filament-associated genes also include the transcription factor *STUI* (Fig. 3C). *Stu1* orthologs have been thoroughly studied in species across the ascomycetes. The *Candida albicans* ortholog, *EFG1*, has long been known to promote filamentation, regulate other developmental transitions, and control gene expression [36-41]. *StuA*, the *Stu1* ortholog in *Aspergillus spp.*, has been extensively studied as a developmental regulator [42, 43]. There are five APSES transcription factors in *H. capsulatum* [44], and *Stu1* is the only one of the five that is transcriptionally induced as yeast-cells transition to filaments (Fig. 5A). As noted above, *msb2* mutant cells fail to induce *STUI* (Fig. 3C, Fig. 5A) but *STUI* expression is restored in the complemented strain at RT (Fig. 5B). To confirm *MSB2* is required for *STUI* expression in independent samples, *STUI* transcript level was measured by qRT-PCR in the control or *msb2* RNAi strain at RT and 37°C. We observed that *STUI* expression was significantly (p<0.001) depleted in the *msb2* knockdown strain at RT (Fig. 5C). To assess whether *STUI* is required for filamentation, we disrupted the *STUI* gene using CRISPR/Cas 9. We observed markedly reduced filamentation in the *stu1* mutant at RT (Fig. 5D), indicating that *STUI* is necessary for proper filamentation. Conversely, overexpression of *STUI* in wild-type *H. capsulatum* was sufficient to cause inappropriate filamentation at 37°C (Fig. 5E). Overexpression of *STUI* in the *msb2* mutant background was sufficient to restore the ability of the *msb2* mutant cells to form filaments at RT,



and, like wild-type, inappropriate filamentation was observed at 37°C (Fig. 5E). Furthermore, cells harboring an *MSB2* overexpression plasmid not only displayed inappropriate filamentation at 37°C (Fig. 1E), but also showed inappropriate transcriptional induction of *STUI* at 37°C (Fig. 5F). Additionally, the *HOG2* RNAi strain, which is yeast-locked at RT (Fig. 4A), fails to induce *STUI* expression at RT (Fig. 5G). Taken together, our data strongly support the model that *MSB2* signals through *HOG2* and *STUI* to stimulate filamentation at RT (Fig. 5H).

**Fig. 5. The conserved transcription factor *STUI* drives filamentation.**

(A) Heatmap showing transcription levels of APSES transcription factors in WT vs *msb2* extracted from the RNAseq dataset. Values are log<sub>2</sub> ratios from single GlcNAc cultures of WT or *msb2* at multiple time points after shift to RT, relative to a WT 37°C sample. (B-C) qRT-PCR analysis of *STUI* transcript level (B) in WT compared to *MSB2* complemented strain after two weeks at RT (not significant); (C) in control compared to *msb2* RNAi strain after 2 weeks at RT (P<0.001). (D) Cell morphology of *stu1* disruption strain and WT after growth at RT for 8 days. (E) Cell morphology of strains produced using a control plasmid or *STUI* overexpression (OE) plasmid in wild-type or *msb2* background. Strains were grown at 37°C or RT. Control strains were imaged after 2 days 37°C or 5 days at RT and overexpression strains were imaged after 4 days at 37°C or RT. (F-G) qRT-PCR analysis of *STUI* transcript level in (F) *MSB2* overexpression strain (P<0.001); (G) *HOG2* RNAi strain (P<0.01). (H) Model showing the genetic relationship between *MSB2*, *HOG2*, and *STUI*.

**The “filament-associated” genes have conserved hyphal expression in ascomycete thermally dimorphic pathogens**



As described above, our analysis defined filament-associated genes, which are expressed in an *Msb2*-dependent manner as wild-type cells undergo filamentation. Similarly, we defined yeast-associated genes, which are expressed in wild-type yeast cells at 37°C, and inappropriately maintain their expression in the *msb2* mutant at RT. As we demonstrated here, the filament-associated genes include regulators of filamentation such as *HOG2* and *STUI*, and the yeast-associated genes include previously identified virulence factors. We reasoned that if these morphological regulators and effectors have conserved function in other dimorphic fungi, then the "filament-associated" and "yeast-associated" regulons identified from our data should show conserved morphology-specific expression in these fungi.

In the case of filament-associated genes, this is indeed the case for three additional *Histoplasma* strains (G186AR, H88, and H143) and three additional thermally dimorphic ascomycetes (*Blastomyces dermatitidis*, *Penicillium marneffei*, and *Candida albicans*) (Fig. 6). In all cases where RNAseq data was available [3, 4, 45-47] (S6 Data), the distribution of yeast/hyphal expression for the orthologs of the filament-associated genes was biased towards showing hyphal expression rather than mimicking the distribution of the global transcriptome ( $p < .05$ , Wilcoxon test; S7 Data). In contrast, in *Ophiostoma novo-ulmi* (the causative agent of Dutch elm disease and the only other dimorphic ascomycete for which we were able to obtain RNAseq data [48]), the distribution of yeast/hyphal expression for the filament-associated regulon was indistinguishable from the rest of the transcriptome. It may be relevant that the dimorphic transition of *O. novo-ulmi* is both temperature-independent and rapid, with cells fully transitioning from yeast to hyphae within 27 hours.

**Fig. 6. *Msb2*-dependent filament-associated genes are conserved across fungi.**

Distributions of  $\log_2$ (yeast/hyphal) expression ratios for orthologs of filament-associated genes from Fig. 3C (blue) or all other detected transcripts (black) for dimorphic ascomycetes with available RNAseq data. See Supp\_comparative\_sources.tdt data for data sources.

Notable genes with conserved hyphal expression in most or all of the thermally dimorphic fungi for which we could detect orthologs include *TYRI* (unique to the Ajellomycetacea), *MS95*, and *STUI* (S7 Data). We note that in *C. albicans*, *STUI* is homologous to the paralogs *EFH1* and *EFG1*. Whereas the *EFG1* transcript is not enriched in *C. albicans* hyphae relative to yeast cells in the RNAseq dataset we analyzed, the Efg1 protein has been shown previously to undergo post-translational regulation that affects its role in *C. albicans* hyphal morphology [49].

In contrast to the filament-associated genes, the yeast-associated genes had conserved morphologic expression only within *Histoplasma* using the same criteria as above. This may be due to *MSB2*-repressed yeast effectors that are unique to *Histoplasma* with respect to the dimorphic fungi listed above (e.g., *CBPI*).

### **The Msb2 and Ryp programs oppose each other in response to temperature**

Our data clearly show that *MSB2* and downstream genes such as *HOG2* and *STUI* are necessary and sufficient to drive a hyphal program. We previously identified four transcription factors, Ryp1-4, that are required for yeast-phase growth at 37°C, and show enhanced expression at 37°C over room temperature [2, 7, 8]. Since the *msb2* mutant is yeast-locked even at room temperature, we hypothesized that the *RYP* transcripts might be inappropriately expressed at RT in the *msb2* mutant, thereby allowing yeast-phase growth to persist. We compared *RYP* transcript levels by qRT-PCR in wild-type and *msb2* mutant strains at 37°C and RT (Fig. 7A). We

determined that all four *RYP* transcripts are inappropriately maintained at higher levels in the *msb2* mutant at RT. This phenotype could be reversed by introducing a wild-type copy of *MSB2* into the mutant strain (Fig. 7B).

**Fig. 7. The Ryp and Msb2 programs antagonize each other in a temperature-dependent manner.**

(A-B) qRT-PCR analysis of *RYP1-4* transcript levels. (A) in WT strain compared to *msb2* after two weeks at RT: *RYP1* ( $P<0.01$ ); *RYP2* ( $P<0.05$ ); *RYP3* ( $P<0.01$ ); *RYP4* ( $P<0.001$ ); (B) in WT strain compared to *MSB2* complemented strain after two weeks at RT. (C) Western blots performed on protein extracted from cells grown either at 37°C or grown at RT for two weeks. GAPDH is a loading control. (D) Location of Ryp3 binding site upstream of *MSB2* transcript. Smoothed Ryp3 ChIP/input ratio ([2], GEO accession number GSE47341) is plotted for WT (dark blue) and *ryp3* control (light blue). Blue triangles indicate locations of Ryp2/Ryp3 associated Motif B [2]. (E) Microarray data [2] showing *MSB2* and *STUI* transcript levels in WT at RT and *ryp3* at 37°C, relative to corresponding transcript level in WT at 37°C. (F) Model showing the temperature-dependent relationship between the Ryps and Msb2.

As described above, we also noted that a number of the “yeast-associated” genes that are inappropriately expressed in the *msb2* mutant at RT are transcriptional targets of Ryp proteins. This observation implied that Ryp protein is inappropriately present at RT in the *msb2* mutant. We examined Ryp1, 2, and 3 proteins by Western blot and determined that these proteins are inappropriately expressed at RT in the *msb2* mutant (Fig. 7C). These data indicate that Msb2 is required to inhibit Ryp transcript and protein expression at RT.

We wondered if there was an analogous antagonistic relationship at 37°C, where Ryp proteins are required to inhibit *MSB2* transcript levels. We examined our previously published chromatin immunoprecipitation (ChIP-chip) data [2] and found a predicted Ryp3 binding site upstream of the *MSB2* open reading frame (Fig. 7D), suggesting that Ryp3 might repress *MSB2* transcript expression at 37°C. We examined our published microarray data for the *ryp3* knockdown strain [2]. In these experiments, as expected, the *MSB2* transcript showed upregulation at RT relative to 37°C in wild-type cells. Notably, the *MSB2* transcript levels at 37°C in the absence of *RYP3* were at levels similar to the wild-type strain at RT (Fig. 7E), indicating that *RYP3* is required to prevent transcriptional induction of *MSB2* at 37°C. Moreover, *RYP3* is also required to inhibit *STUI* expression at 37°C (Fig. 7E), indicating that the Ryp program inhibits expression of multiple members of the Msb2 hyphal program at 37°C. Taken together, we conclude that the Ryp program and the Msb2 program oppose each other in a temperature-dependent manner such that the Ryp program dominates at 37°C whereas the Msb2 program dominates at RT, ultimately allowing regulation of cell shape by temperature (Fig. 7F).

## Discussion

Thermally dimorphic fungal pathogens display an exquisite response to temperature that governs their ability to thrive in either the environment of the soil or the mammalian body. At environmental temperatures, these fungi grow in a multicellular hyphal form that is suited to survival in the soil. In response to mammalian body temperature, these organisms switch to a host form (unicellular yeast cells in the case of *Histoplasma*) that expresses virulence factors that are important for disease progression. Although the existence of this response is thought to be critical to the pathogenesis of endemic fungi such as *Histoplasma*, *Blastomyces*,

*Paracoccidioides*, and *Coccidioides* spp.[50], the molecular basis of this dichotomous response to temperature is not understood. Here we uncover a pathway required for the filamentation response to environmental temperature, and show that this pathway acts in a mutually antagonistic manner with the regulators that promote the morphologic response to mammalian body temperature. Thus regulation of thermal dimorphism represents a binary choice, where the fungus transitions between two different states in response to temperature by means of opposing regulatory pathways.

A number of cell fate decisions in both microbes and multicellular organisms are controlled by bistable switches, where feedback mechanisms drive the system into one of two possible dynamic steady states. In the case of *Histoplasma* thermal dimorphism, we showed previously that the Ryp proteins form an interlocking circuit that promotes yeast-phase growth [2]. All four Ryp proteins bind upstream of Ryp1, 2, and 4, which is thought to create a positive feedback loop that contributes to the marked increase of Ryp protein levels at 37°C. Additionally, ribosomal profiling data suggest that the translational efficiency of Ryp1 and Ryp2 is higher at 37°C than at room temperature [4]. Notably, Ryp3 associates with the upstream region of the *MSB2* gene at 37°C, which is associated with decreased accumulation of the *MSB2* transcript [2] (Fig. 7). Thus several mechanisms cooperate to enhance Ryp protein levels at 37°C. In contrast, at room temperature, the abundance of Ryp proteins is markedly decreased. During room temperature growth, *MSB2* is required for the morphologic switch from yeast-phase cells to hyphal cells, and molecularly, *MSB2* is required for the reduction in Ryp transcript and protein levels (Fig. 7). Notably, overexpression of *MSB2* is sufficient to trigger morphologic changes even at 37°C (Fig. 1), suggesting that levels of *MSB2* are a critical determinant of cell shape. Taken together, our data show that in wild-type cells, inhibition of the Ryp pathway by an Msb2-

dependent mechanism is likely to be a key step in the switch from yeast to filaments in response to temperature. Notably, in *Candida spp*, an antagonistic regulatory relationship between Wor1 (an ortholog of the *H. capsulatum* Ryp1 protein) and Efg1 (an ortholog of the *H. capsulatum* Stu1 protein) is critical for controlling the switch between distinct developmental states [36, 51-53].

*In vitro* experiments show that morphologic transitions in the thermally dimorphic fungi occur over the timescale of days rather than hours, suggesting that sustained changes in temperature are required to shift cell shape. It is attractive to speculate that *H. capsulatum* is integrating a temperature signal over time to enable specific responses to the homeostatic temperature environment of the host while ignoring transient spikes in environmental temperature. In molecular terms, the precise temporal parameters that govern a switch between dominance of the Ryp pathway and dominance of the Msb2 pathway are unknown. Further experimentation may reveal that sustained but not transient temperature shifts are required for one antagonistic program to fully disrupt and reset the other. Ultimately, multiple molecular mechanisms could contribute to the role of temperature in these cell-fate decisions.

This study is the first to elucidate components of the pathway that transduce the morphogenetic response to environmental temperature in *H. capsulatum*. The role of Msb2, a transmembrane signaling mucin, in temperature response is particularly intriguing. Msb2 has long been known to regulate the filamentous growth and osmosensing pathways in *Saccharomyces cerevisiae* [11, 14] as well as to influence morphogenesis in *C. albicans* [10] and appressorium development in *Ustilago maydis* [30]. Additionally, recent work showed that Msb2 regulates responses to temperature stress in *C. albicans* [54]. It has become clear that proteolytic cleavage of the extracellular domain of Msb2 is a critical step in signaling [12, 55]. It is

attractive to speculate that this type of cleavage could be triggered by temperature shift and contribute to Msb2-dependent responses in *H. capsulatum*.

A number of studies have shown that the transcriptional profile of 37°C-grown yeast cells and RT-grown hyphae are quite distinct [2-7], although the function of the majority of differentially expressed transcripts is unknown. One of the most informative aspects of the *MSB2* analysis is that the compact Msb2 regulon distinguishes genes that are key to filamentation or virulence from the high background of differentially expressed factors. As shown in this work, the *msb2* mutant appears morphologically as yeast at room temperature, and we initially predicted that its transcriptional profile would resemble that of wild-type yeast cells. In actuality, during RT growth, the *msb2* mutant transcriptome resembles that of wild-type hyphae despite its yeast-phase morphology. Specifically, the majority of the ~1870 genes that are differentially expressed by wild-type hyphae at RT are also expressed by *msb2* mutant yeast at RT (Fig. 3A), and the majority of the ~1100 genes that are differentially expressed by wild-type yeast cells at 37°C fail to maintain their expression at RT in the yeast-form *msb2* mutant (Fig. 3A). The fact that the *msb2* mutant grows as yeast at RT but looks transcriptionally like hyphae is in contrast to the *ryp* mutants, which appear both morphologically and transcriptionally as hyphae at 37°C [2, 7]. Interestingly, there is a set of yeast-phase genes whose expression is decreased after shift to room temperature in wild-type cells, but inappropriately maintained in the *msb2* mutant. This set of 40 genes is enriched for direct transcriptional targets of the Ryp transcription factors, suggesting that the Ryp proteins, which are present at inappropriately high levels at RT in the *msb2* mutant, are able to activate expression of a subset of their targets. Importantly, our experiments reveal that expression of these genes appears to be tightly coupled to growth in the yeast form rather than growth at 37°C. It seems highly significant that this gene set contains

known virulence factors and small, secreted proteins that may be well positioned to manipulate events within host cells. Thus the transcriptional analysis of the *msb2* mutant defined 40 yeast-associated transcripts that are of high interest for potential roles in virulence.

In an analogous fashion, the *msb2* mutant also defined a core set of 167 filament-associated transcripts that fail to be expressed in the yeast-locked mutant at RT when compared to wild-type cells, suggesting that only a fraction of the RT transcriptional program is required for the developmental process of filamentation. This result is analogous to findings in *C. albicans*, where the majority of the transcriptional response to filamentation conditions is dispensable for hyphal formation [47, 56]. We reasoned that the *H. capsulatum* *MSB2* regulon was likely to contain regulators and effectors of the hyphal growth program, and indeed confirmed that *HOG2* and *STUI* are required for filamentation. Notably, a recent publication examining APSES transcription factors in *H. capsulatum* showed that knockdown of *STUI* resulted in a defect in aerial hyphae production [44], which agrees with our data indicating that *STUI* is involved in hyphal growth. A number of other putative regulatory factors are members of the *MSB2* regulon, and their functions remain to be analyzed. When this *Msb2*-dependent filamentous gene set was compared to other *Histoplasma* strains and more distantly related fungi, we found that, on average, this gene set displayed conserved, filament-enriched expression. This suggests that the *MSB2* regulon is conserved across a diverse group of fungi and that the genes involved in filamentation are shared in multiple fungal species. Notably, this filament-associated gene set may highlight factors that have important regulatory or effector roles in the hyphal growth program of other fungi. We are particularly interested in ucsf\_hc.01\_1.G217B.09523, which lacks an ortholog in *C. albicans*, but was present with conserved hyphal enrichment in all other fungi in our dataset, including *O. novo-ulmi*. This gene is orthologous to fungal



*Metarhizium anisopliae* Crp2 and plant *Arabidopsis thaliana* atRZ-1a, sharing their N-terminal RRM\_1 RNA binding domain and glycine rich C-terminus. Crp2 and atRZ-1a are thought to provide cold resistance by acting as RNA chaperones, and are sufficient to confer this resistance when heterologously expressed in *Saccharomyces cerevisiae* [57] and an *Escherichia coli* cold-sensitive mutant [58], respectively. Intriguingly, two other RRM\_1 family members with broad RNA binding activity have distinct effects on hyphal growth in the basidiomycete dimorphic fungus *Ustilago maydis*: Rrm4 mediates proper polar growth filaments via long-distance transport of mRNAs along hyphae while Grp1 may act as an RNA chaperone as well as an accessory component in endosomal mRNA transport [59]. The intriguing relationship between expression of putative RNA-binding factors and thermal dimorphism has yet to be fully explored, but may shed light on how adaptation of fungi to different environments is linked to morphologic changes.

Finally, the identification of regulators of the environmental differentiation program has provided insight into critical antagonists of the host program. Elucidating the molecular switches that control the host program may lead to novel tactics to target virulence strategies in thermally dimorphic fungi.

## Materials and Methods

### *H. capsulatum* Strains

*Histoplasma capsulatum* G217B *ura5<sup>-</sup>* (WU15; referred to in this work as wild-type) was used for the insertional mutagenesis screen, which was performed using the *Agrobacterium tumefaciens* strain LBA1100 containing the plasmid pRH5b as described [7]. All strain manipulations were done in the *H. capsulatum* G217B *ura5<sup>-</sup>* background. Strains used in this

paper can be found in Table S3.

## Media

Wild-type G217B, complementation, and RNAi strains were grown in liquid or solid *Histoplasma* Macrophage Medium (HMM) [60]. G217B *ura5<sup>-</sup>* and insertional mutants were grown in HMM supplemented with 2.5mg/ml uracil. Broth cultures were grown at 37°C with 5% CO<sub>2</sub> on an orbital shaker at 120–150 rpm. For yeast-form colonies, plates were grown in a humidified incubator at 37°C with 5% CO<sub>2</sub>. Plates grown at room temperature were wrapped individually in Parafilm and placed in a 25°C standing incubator in a Biosafety Level 3 facility. Cells were thawed fresh from frozen stock and passaged on HMM agarose plates every 1–2 weeks for up to 2 months. 100mM N-acetylglucosamine (Sigma-Aldrich) was used as the primary carbon source in HMM where indicated.

## Screen for yeast-locked mutants

A small pilot screen was performed to identify yeast-locked mutants. *H. capsulatum* G217B *ura5<sup>-</sup>* yeast were co-cultured with *Agrobacterium* carrying plasmid pRH5b, which has a T-DNA element with a hygromycin marker, allowing for selection of insertion mutants as previously described [7]. The transformants were plated at a dilution aiming for 50 colonies per plate and were incubated at 37°C, 5% CO<sub>2</sub> until colonies appeared. After 25 days, the plates were transferred to a room temperature incubator in the Biosafety Level 3 laboratory. 92 colonies were observed for colony morphology. After 13 days, 2 of the 92 colonies retained yeast morphology while the others had filamentous projections.

## Mapping the T-DNA insertion in the *msb2* mutant

The location of the T-DNA insertion was determined from whole-genome sequencing. The SG1 strain (the *msb2* mutant) and the control strain were grown up from frozen glycerol stocks onto HMM agarose plates supplemented with uracil. Liquid cultures were inoculated for each strain and passaged twice at 1:25 dilution in HMM + uracil. Two days after the second passage, the cells were collected via centrifugation. Genomic DNA of the strains was isolated using the Gentra Puregene Yeast Kit (Qiagen). Libraries were made using the Nextera XT DNA Library Prep Kit (Illumina).

The libraries were multiplexed with 7 other *Histoplasma* genomic DNA libraries and sequenced on an Illumina HiSeq 4000, yielding 21,038,128 paired-end 101-mer reads for ~106x coverage of the ~40MB G217B genome. Bowtie2 was used to map these read pairs to a joint index of the T-DNA plasmid sequence (pRH5b) plus the 11/30/2004 version of the G217B genome assembly from the Genome Sequencing Center at Washington University (GSC) as mirrored at <http://histo.ucsf.edu/downloads/>. The location of the T-DNA insert was determined by filtering for discordantly mapped read pairs with one mate in the T-DNA and one mate in the reference genome. Discordant pairs consistently mapped to the T-DNA left border and HISTO\_ZL.Contig1131 or the T-DNA right border and HISTO\_DA.Contig93, indicating a single insertion event resulting in a genomic rearrangement. A crude model of the HISTO\_ZL.Contig1131/T-DNA/HISTO\_DA.Contig93 hybrid sequence was constructed based on the discordantly mapped reads, erring on the side of including extra sequence from HISTO\_ZL.Contig1131 and HISTO\_DA.Contig93. Bowtie2 was then used to remap the reads to the model sequence, allowing the exact junction coordinates to be determined from the alignment of reads spanning the genomic/T-DNA boundaries. This gave a final model of the hybrid sequence (Fig. 1B, bottom), which was confirmed by Bowtie mapping of RNAseq reads

from the SG1 mutant to the hybrid sequence (Fig. 1B, green coverage lines).

It is evident from this RNAseq mapping that *MSB2* is transcribed in the SG1 mutant, with transcription initiating from a site in the T-DNA. The resultant extended 5' region of the transcript introduces a new open reading frame (pink box in Fig. 1B), which we hypothesize acts as an upstream ORF that inhibits Msb2 translation.

# **Construction of plasmids**

Sub-cloning was performed in *E. coli* DH5 $\alpha$ . All knockdown constructs were generated by amplifying a 500 bp region to target a particular gene for RNA interference as previously described [2]. The *MSB2* gene was targeted by amplifying G217B genomic DNA using OAS5486-87 (primers are described in Table S2). The fragment was cloned into the pDONR/Zeo vector using Gateway Cloning Technology (Invitrogen), generating the plasmid pLR04. The final episomal RNAi vector was produced by recombining pSB23 [2] with pLR04, generating plasmid pLR11. The *HOG2* RNAi strains were generated with the same methodology. The 500 bp region used to target *HOG2* was amplified using OAS5005-06 and the PCR product was cloned into pDONR/Zeo, generating plasmid pSB420. This plasmid was then recombined with pSB23 to produce pSB430.

The overexpression construct for *STUI* was produced by placing the *ACT1* promoter upstream of the *STUI* coding sequence. This construct was generated by amplifying the *ACT1* promoter using OAS2594 and OAS2587, *STUI* using OAS2586 and OAS2592, and the *CATB* terminator using OAS2591 and OAS2593. The fragments were joined using the overlap extension PCR cloning method [61]. The construct was recombined with pDONR/Zeo using Gateway Cloning Technology (Invitrogen) to produce the plasmid pTM1. pTM1 was recombined with the destination plasmid pDG33 to produce the final plasmid pTM2. The

overexpression construct for *MSB2* was produced by placing the GAPDH promoter upstream of the *MSB2* coding sequence. OAS5476-77 was used to amplify the coding sequence by PCR. Using Gateway Cloning Technology (Invitrogen), the PCR fragment was cloned into the pDONR/Zeo plasmid to produce pLR09. This plasmid was recombined with pSB234, which is an episomal plasmid carrying the GAPDH promoter upstream of the Gateway cassette, making it compatible with the Gateway cloning system (Invitrogen). pLR02 and pSB234 were recombined to produce pLR09.

The *MSB2* complementation construct was generated by using the primers OAS5711-12 to amplify the gene and its native promoter. This fragment was cloned into pDONR/Zeo using Gateway Cloning Technology (Invitrogen) to produce the plasmid pLR01. pLR01 was recombined with pDG33 to produce pLR09.

The plasmids used in this study are described in Table S1 and primers are described in Table S2. To generate RNAi, complementation, or overexpression strains, episomal plasmids were linearized, electroporated into the relevant yeast strain, and plated onto HMM agarose plates without uracil to select for *URA5*-containing transformants. The strains used in this study are described in Table S3.

# **Construction of the *stu1* mutant strain**

We made an episomal version (pBJ219) of an *Agrobacterium*-based Cas9 targeting plasmid [62] (kind gift of Bruce Klein). pBJ219 is described in a separate manuscript (Joehnk and Sil, in preparation). In brief, a guide RNA targeting the first exon of *STU1* was introduced into pBJ219, which carries a fungal codon-optimized Cas9 along with ribozyme sequences required for the excision of the RNA sequence [62]. The plasmid was introduced into the G217B *ura5<sup>-</sup>* strain by electroporation and transformants were selected on HMM plates lacking uracil. Single colonies

were isolated from individual transformants, genomic DNA was prepared, and the genomic sequence at the *STUI* locus was analyzed using TIDE, an online tool measuring the frequency of indels in a mixed population [63]. After three sequential passages, we colony-purified an isolate with an insertion of an A at position 23 relative to the ATG, resulting in an early stop codon after amino acid 13.

## **Imaging**

Imaging was performed on a Zeiss AxioCam MRM microscope using DIC at 100X magnification.

## **Culture conditions for expression profiling**

Samples were prepared in two distinct experiments and sequenced in three distinct sequencing runs. In the first experiment, two biological replicates each of *Histoplasma capsulatum* G217B *ura5* (WU15) or an *msb2* mutant derived from WU15 (SG1) were grown in liquid culture with glucose or GlcNAc as the carbon source at 37°C. After taking  $t = 0$  samples, the cultures were moved to RT and additional samples were taken at 2, 6, 8, or 10 days. One sample was not usable for sequencing, giving a total of  $2 \times 2 \times 2 \times 5 - 1 = 39$  samples. The GlcNAc samples were sequenced in part of a lane of one HiSeq 4000 run (batch 1) and the glucose samples were sequenced in part of a lane of a second run (batch 2). Because these samples were grown at the same time, they were treated as a single batch in our limma analysis. In the second experiment, two biological replicates each of WU15, SG1, or SG1 complemented with the wild-type *MSB2* gene (SG1cMSB2) (resulting from transformation with pLR08) were grown in liquid culture with glucose or GlcNAc as the carbon source at 37°C. In this second experiment, WU15 and SG1 were transformed with the empty vector control (pLH211) so that the parental, mutant, and complemented strains were comparable. After taking  $t = 0$  samples, the cultures were moved to

RT and additional samples were taken at 2, 6, or 8 days, for a total of  $2 \times 3 \times 2 \times 4 = 48$  samples.

These samples were sequenced in a full lane of a HiSeq 4000 run (batch 3).

# **Fungal cell collection**

*H. capsulatum* cultures were collected by filtration with a disposable filtration apparatus either in the Biosafety Level 2\* laboratory for cultures grown at 37°C or in the Biosafety Level 3 laboratory for cultures grown at RT. At each time point, 10 mL of fungal culture was passed through a disposable filter apparatus (Thermo Fisher), the cells were scraped off with a cell scraper, placed into a conical tube with 1 mL Trizol reagent (Qiazol) and flash frozen in liquid nitrogen. Samples were stored at -80°C until all time points were collected.

# **RNA extraction**

Total RNA was extracted from fungal cells using a Trizol-based RNA extraction protocol. Frozen, resuspended pellets of cells were incubated at RT for 5 minutes to thaw. The lysate was subjected to bead beating (Mini-Beadbeater, Biospec Products) followed by a chloroform extraction. The aqueous phase was then transferred to an Epoch RNA column where the filter was washed with 3 M NaOAc and then with 10 mM TrisCl in 80% EtOH. DNase (Purelink) treatment was used to remove any residual DNA and the filters were washed again with NaOAc and TrisCl before eluting the RNA in nuclease-free water. RNA quality was determined with a High Sensitivity DNA Bioanalyzer chip (Agilent).

# **mRNA isolation**

20 µg of total RNA was purified for mRNA through polyA selection with Oligo-dT Dynabeads (Thermo Fisher) as described in the manufacturer's protocol. Ribosomal RNA depletion was confirmed with an RNA 6000 Nano Bioanalyzer chip (Agilent Technologies). Samples were acceptable for library preparation at 0-4% rRNA.

## RNAseq library preparation

Libraries for RNAseq were prepared using the NEB Next Ultra Directional RNA Library Prep Kit (New England Biolabs). Individual libraries were uniquely barcoded with NEBNext Multiplex Oligos for Illumina sequencing platform (New England Biolabs). Average fragment size and presence of excess adapter was determined with High Sensitivity DNA bioanalyzer chip from Agilent Technologies. Libraries had an average fragment length of 300-500 bp. The concentration of the individual libraries was quantified through Qubit dsDNA High Sensitivity Assay (Invitrogen). 5 ng of each library was pooled into three final libraries (batches in S1 Data) and run on a High Sensitivity DNA Bioanalyzer chip to determine the average fragment size of the final pooled samples. The final libraries were submitted to the UCSF Center for Advanced Technology for sequencing on a Illumina HiSeq4000 sequencer.

## RNAseq data analysis

For quality control, data exploration, and to quantify yields (mapping statistics in S1 Data), samples were aligned with Bowtie version 1.1.2 [64] to the 11/30/2004 version of the G217B genome assembly from the Genome Sequencing Center at Washington University as mirrored at <http://histo.ucsf.edu/downloads/>, invoked as:

```
zcat SAMPLE.fastq.gz | bowtie -tS -p 4 -k 1 HcG217B - \
| samtools view -bS -o SAMPLE.bam
```

Transcript abundances were quantified based on version ucsf\_hc.01\_1.G217B of the *H. capsulatum* G217B transcriptome (S5 Data of [4],



[http://histo.ucsf.edu/downloads/ucsf\\_hc.01\\_1.G217B.transcripts.fasta](http://histo.ucsf.edu/downloads/ucsf_hc.01_1.G217B.transcripts.fasta))

Relative abundances (reported as TPM values [65]) and estimated counts (est\_counts) of each transcript in each sample were estimated by alignment free comparison of k-mers between the preprocessed reads and assembled transcripts using KALLISTO version 0.43.0 [66], invoked as:

```
kallisto quant -i ucsf_hc.01_1.G217B.idx -t 4 -b 100 \
--single --rf-stranded -l 250 -s 50 \
-o SAMPLE.kallisto SAMPLE.fastq.gz
```

Further analysis was restricted to transcripts with TPM  $\geq 10$  in at least one sample.

Differentially expressed genes were identified by linear regression on four independent factors using LIMMA version 3.26.1 [67, 68].

Specifically, KALLISTO est\_counts were rounded to the nearest integer and imported as counts per million (CPM) in a DGEList in R. Samples were TMM normalized with calcNormFactors and VROOM [69] was used to estimate the mean/variance trend.

Samples were classified on four factors (S1 Data): batch (1 or 3), genotype (wild-type or *msb2*), media (GlcNAc or glucose), and time (0, 2d, 6d, or 8d). For each transcript, the observed counts were fit to a model assuming a mean transcript abundance plus independent corrections for each factor using lmFit. Shrinkage of the individual gene variances over the full data was applied

with eBayes, and significantly differential genes were identified by extracting genes with a significantly non-zero genotype and/or t = 8d coefficient (at 5% FDR) with an effect size of at least 2x (absolute log2 fold change >= 1) using topTable.

The core R invocations were:

```
dge <- DGEList(counts)
dge <- calcNormFactors(dge)
time <- as.factor(time)
genotype <- as.factor(genotype)
media <- as.factor(media)
batch <- as.factor(batch)
d <- model.matrix(~time+genotype+media+batch)
v <- voom(dge, d)
fit <- lmFit(v, d)
fit <- eBayes(fit)
topTable(fit, coef="genotypeMSB2", n = 20000, lfc=1, p.value = .05)
topTable(fit, coef="time8", n = 20000, lfc=1, p.value = .05)
```

## Comparative Transcriptome Analysis

For all ascomycete yeast/hyphal dimorphs with available RNAseq data (S6 Data), reads were downloaded from the SRA. Corresponding transcriptome sequences were downloaded from the sources indicated in S6 Data and indexed for kallisto. Transcripts were quantified for each

sample with kallisto, invoked as:

kallisto quant -i TRANSCRIPTOME.idx -t 4 EXTRA RUNS

where TRANSCRIPTOME.idx is the appropriate index file, RUNS are gzipped FASTQ files for all runs corresponding to the given sample, and EXTRA are additional flags for appropriate experiment-specific handling of single-ended and strand-specific reads, as given in the "extra flags" column of S6 Data. Further analysis was restricted to transcripts with TPM  $\geq 10$  in at least one sample. Log<sub>2</sub>(TPM) values were averaged across biological replicates for each morphology and then subtracted to give log<sub>2</sub>(yeast/hyphae) differential expression values (S7 Data). *Histoplasma* ortholog groups were taken from S13 Data from [4]. Othologous genes between G217B and the remaining species were determined by InParanoid version 1.35 [70]. We tested the null hypothesis that log<sub>2</sub>(yeast/hyphae) values for orthologs of the G217B filament-associated genes were drawn from the same distribution as the remaining genes with G217B orthologs using a two sided Wilcoxon rank sum test. Kernel density estimates of these distributions are plotted in Fig. 6 using R's density function.

### **Expression level by qRT-PCR**

Relative gene expression was observed using qRT-PCR. RNA extraction was performed as described above and cDNA was prepared by priming 2 µg DNase-treated total RNA with oligodT/pdN9 and dNTPs. Samples were shifted to RT and treated with 1 µl RNaseOUT Recombinant Ribonuclease Inhibitor (ThermoFisher) and 0.5 µl Thermo Scientific Maxima H Minus Reverse Transcriptase. Negative controls included samples that were processed without the addition of reverse transcriptase. Samples with reverse transcriptase were diluted 1:50 and

samples without reverse transcriptase were diluted 1:10. qPCR reactions were set up using 96-well plates with FastStart Universal SYBR Green Master (Roche) and 1.5  $\mu$ M primers (found in Table S2), and plates were read using a Mx3000P machine (Stratagene) and analyzed using MxPro software (Stratagene).

### **Protein extraction**

Organic fractions from Trizol RNA extractions were stored at -20°C until protein extraction was performed. To remove DNA from the organic fraction, 100% ethanol was added and samples were centrifuged to pellet DNA. The phenol-ethanol supernatant containing the proteins was transferred to a fresh tube containing isopropanol and centrifuged at 4°C to pellet the protein precipitate. The pellet was washed with 0.3M guanidine thiocyanate in 95% ethanol and centrifuged at 4°C. This wash step was repeated for a total of 3 times. 100% ethanol was added to the protein, which was then vortexed and incubated for an additional 20 minutes at RT. The protein was pelleted by centrifugation and air-dried at RT. Finally, the pellet was resuspended in urea lysis buffer (9M Urea, 25mM Tris-HCl, 1mM EDTA, 1% SDS, 0.7M  $\beta$ -mercaptoethanol). If necessary, the pellet was incubated at 50°C for 10-20 minutes to successfully resuspend it in the urea buffer. Once resuspended, the sample was boiled, centrifuged, and the supernatant, containing the protein was transferred to a new tube for quantification by the Pierce 660 nm assay with added ionic detergent compatibility reagent (ThermoFisher).

### **Expression level by Western blot**

Western blotting was performed with polyclonal peptide antibodies against either Ryp1, Ryp2, or Ryp3; antibodies were described previously [2]. Following quantification of protein, 10-30  $\mu$ g was resuspended in a total of 20  $\mu$ L of urea lysis buffer and 6  $\mu$ L NuPAGE LDS Sample Buffer

(Novex). The samples were boiled and electrophoresed on a 10-well NuPAGE 4-12% BT SDS-PAGE gel (Novex) in MOPS running buffer at 150V. The protein was then transferred to a nitrocellulose membrane at ~40 V for 2 hours. The membrane was incubated with blocking solution (1 g milk powder in 100 mL wash buffer (0.1% Tween-20 in PBS)) for an hour and then incubated in the primary antibody in wash buffer overnight at 4°C. Primary antibody dilutions were:  $\alpha$ -Ryp1 (1:10,000),  $\alpha$ -Ryp2 (1:2,500),  $\alpha$ -Ryp3 (1:5,000),  $\alpha$ -GAPDH (1:1,000). The blot was washed and secondary antibody (for Ryp proteins: Goat  $\alpha$ -rabbit HRP (GenScript) 1:1,000, for GAPDH: Goat  $\alpha$ -mouse HRP (ThermoFisher) 1:1,000) was added to the blot for 1 hour at RT followed by another wash. Protein bands were detected using chemiluminescence according to the manufacturer's instructions (SuperSignal West Pico kit (ThermoFisher)).

### MAP Kinase Phylogeny

MAPK protein sequences were obtained from SGD (<https://www.yeastgenome.org/>; HOG1=YLR113W, SMK1=YPR054W, KDX1=YKL161C, STL2=YHR030C, KSS1=YGR040W, FUS3=YBL016W), AspGD (<http://www.aspgd.org/>; mpkC=AN4668), and HistoBase (<http://histo.ucsf.edu/downloads/>; HOG1=ucsf\_hc.01\_1.G217B.05737, HOG2=ucsf\_hc.01\_1.G217B.11562, HMK1=ucsf\_hc.01\_1.G217B.10227, SLT2=ucsf\_hc.01\_1.G217B.02764) and aligned with PROBCONS 1.12 [71]. A phylogeny was generated from the PROBCONS alignment with FastTree 2.1.7 [72].

### Statistical Analysis

qRT-PCR data analysis was performed as described [73] with significant differences determined by the t-test using propagated standard errors. Growth curves were graphed with Prism (GraphPad Software).

## Other software and libraries

We wrote custom scripts and generated plots in R 3.2.3 [74] and Python 2.7.12, using Numpy 1.11.0 [75] and Matplotlib 1.5.1 [76]. Jupyter notebooks [77] and JavaTreeView [78] were used for interactive data exploration.

## Acknowledgements

We are grateful to Chad Rappleye and Bruce S. Klein for generously sharing reagents and information. We thank Bevin English for helpful advice and suggestions, and Alexander Johnson and Joseph DeRisi for their guidance. Eric Chow and the UCSF Center for Advanced Technology provided invaluable advice on library preparation and sequencing. We also thank Davina Hocking Murray for help with figure design and preparation, as well as Hiten Madhani, Suzanne Noble, and members of the Sil and Noble labs for helpful discussions.

## Supporting Figure Legends

### **Fig. S1. *msb2* yeast grown at RT continue to divide.**

A growth curve of WT and mutant yeast at 37°C is compared to mutant yeast grown at RT.

### **Fig. S2. Cells grown for RNAseq timecourse in glucose as carbon source do not transition to hyphae by 8 days after the shift to RT.**

Cells were grown at 37°C (d0) or RT (d2-8) and imaged to observe morphology.

## Supporting Data Legends

All supporting data is formatted as Excel-compatible tab-delimited text. Expression profile data sets (S2, S3, S4, S5, and S7) are additionally compatible with the JavaTreeView extended CDT file format.

**S1 Data. Sequenced samples.** Columns are unique sample name; the four parameters used in limma fitting, *viz.*: genotype (G217B = wild type, SG1 = *msb2* mutant, SG1cMSB2 = complemented *msb2* mutant), time (days after transfer to RT), media (GlcNAc or glucose), and batch (where batches 1 and 2 were treated as a single batch while fitting); as well as total read depth and number of reads mapped to the G217B assembly by Bowtie.

## **S2 Data. Kallisto estimated counts and limma fit values for time course expression profiles.**

Each row is a transcript, with the UNIQID column giving the ucsf\_hc.01\_1.G217B systematic gene name. The NAME column gives short names taken from Data S13 of [4] with additional names and corrections based on human curation. The next 87 columns give KALLISTO estimated counts for each transcript in each sample (with column headings corresponding to sample names in Data S1). Additional annotation columns are G217B predicted gene name, *H. capsulatum* conserved enrichment, and signal peptide, all taken from Data S13 of [4]; and ChIP-chip based association with the Ryp transcription factors, taken from Table S2 of [2]. The final 7 columns give the limma fit parameters, in units of log<sub>2</sub>(CPM), and the corresponding limma probabilities that the non-intercept parameters differ from 0 are given as p(parameter).

**S3 Data. Genes significantly differential in response to temperature and/or *msb2* mutation.**

Each row is a transcript, with UNIQID and NAME, and annotation columns as in S2 Data with an additional “class” column indicating the eight expression classes from Fig 3A and 3B. The remaining columns give log2(CPM) values for each sample (with column heading corresponding to sample names in Data S1). These values are TMM normalized and each row is mean centered, such that positive and negative values correspond to samples with expression above or below the average expression for a given transcript. As in Fig 3B, transcripts are grouped by class and sorted by the limma fit MSB2 dependence within each class.

**S4 Data. Yeast associated genes.** Subset of S3 Data corresponding to genes significantly up in both WT/*msb2* and t = 0/t = 8d.

**S5 Data. Filament associated genes.** Subset of S3 Data corresponding to genes significantly down in both WT/*msb2* and t = 0/t = 8d.

**S6 Data. Data sources for comparative transcriptome analysis.** Each row corresponds to a profile plotted in Fig. 6. Columns give a shorthand name, taxonomic details (genus, species, strain), data accession (SRA) and reference (PMID), sequencing layout (paired or single), extra flags supplied to KALLISTO (see Methods), the specific SRA sample accessions used for the yeast or hyphae expression profiles, and the source of the transcriptome FASTA file used to build the KALLISTO index.



914    **S7 Data. Comparative transcriptome profiles.** Each row corresponds to an *H. capsulatum*  
915    G217B transcript with UNIQID, NAME, G217B predicted gene, and class as in Data S3.  
916    Ortholog columns give the corresponding transcript names for other species (empty for cases  
917    with no mapped ortholog). Y/H columns give KALLISTO estimated log2(yeast/hyphae)  
918    expression ratios corresponding to sample names in Data S6.  
919

## References

1. Manos NE, Ferebee SH, Kerschbaum WF. Geographic variation in the prevalence of histoplasmin sensitivity. *Dis Chest*. 1956;29(6):649-68. Epub 1956/06/01. PubMed PMID: 13317782.
2. Beyhan S, Gutierrez M, Voorhies M, Sil A. A temperature-responsive network links cell shape and virulence traits in a primary fungal pathogen. *PLoS Biol*. 2013;11(7):e1001614. Epub 2013/08/13. doi: 10.1371/journal.pbio.1001614. PubMed PMID: 23935449; PubMed Central PMCID: PMC3720256.
3. Edwards JA, Chen C, Kemski MM, Hu J, Mitchell TK, Rappleye CA. Histoplasma yeast and mycelial transcriptomes reveal pathogenic-phase and lineage-specific gene expression profiles. *BMC Genomics*. 2013;14:695. Epub 2013/10/12. doi: 10.1186/1471-2164-14-695. PubMed PMID: 24112604; PubMed Central PMCID: PMC3852720.
4. Gilmore SA, Voorhies M, Gebhart D, Sil A. Genome-Wide Reprogramming of Transcript Architecture by Temperature Specifies the Developmental States of the Human Pathogen Histoplasma. *PLoS Genet*. 2015;11(7):e1005395. Epub 2015/07/16. doi: 10.1371/journal.pgen.1005395. PubMed PMID: 26177267; PubMed Central PMCID: PMC4503680.
5. Hwang L, Hocking-Murray D, Bahrami AK, Andersson M, Rine J, Sil A. Identifying phase-specific genes in the fungal pathogen Histoplasma capsulatum using a genomic shotgun microarray. *Mol Biol Cell*. 2003;14(6):2314-26. Epub 2003/06/17. doi: 10.1091/mbc.E03-01-0027. PubMed PMID: 12808032; PubMed Central PMCID: PMC194881.
6. Inglis DO, Voorhies M, Hocking Murray DR, Sil A. Comparative transcriptomics of infectious spores from the fungal pathogen Histoplasma capsulatum reveals a core set of

944 transcripts that specify infectious and pathogenic states. *Eukaryot Cell*. 2013;12(6):828-52. Epub  
945 2013/04/09. doi: 10.1128/EC.00069-13. PubMed PMID: 23563482; PubMed Central PMCID:  
946 PMCPMC3675998.

947 7. Nguyen VQ, Sil A. Temperature-induced switch to the pathogenic yeast form of  
948 *Histoplasma capsulatum* requires Ryp1, a conserved transcriptional regulator. *Proc Natl Acad*  
949 *Sci U S A*. 2008;105(12):4880-5. Epub 2008/03/15. doi: 10.1073/pnas.0710448105. PubMed  
950 PMID: 18339808; PubMed Central PMCID: PMCPMC2290814.

951 8. Webster RH, Sil A. Conserved factors Ryp2 and Ryp3 control cell morphology and  
952 infectious spore formation in the fungal pathogen *Histoplasma capsulatum*. *Proc Natl Acad Sci U*  
953 *S A*. 2008;105(38):14573-8. Epub 2008/09/16. doi: 10.1073/pnas.0806221105. PubMed PMID:  
954 18791067; PubMed Central PMCID: PMCPMC2567189.

955 9. Bender A, Pringle JR. A Ser/Thr-rich multicopy suppressor of a *cdc24* bud emergence  
956 defect. *Yeast*. 1992;8(4):315-23. Epub 1992/04/01. doi: 10.1002/yea.320080409. PubMed  
957 PMID: 1514328.

958 10. Roman E, Cottier F, Ernst JF, Pla J. Msb2 signaling mucin controls activation of Cek1  
959 mitogen-activated protein kinase in *Candida albicans*. *Eukaryot Cell*. 2009;8(8):1235-49. Epub  
960 2009/06/23. doi: 10.1128/EC.00081-09. PubMed PMID: 19542310; PubMed Central PMCID:  
961 PMCPMC2725568.

962 11. Cullen PJ, Sabbagh W, Jr., Graham E, Irick MM, van Olden EK, Neal C, et al. A  
963 signaling mucin at the head of the Cdc42- and MAPK-dependent filamentous growth pathway in  
964 yeast. *Genes Dev*. 2004;18(14):1695-708. Epub 2004/07/17. doi: 10.1101/gad.1178604. PubMed  
965 PMID: 15256499; PubMed Central PMCID: PMCPMC478191.

12. Vadaie N, Dionne H, Akajagbor DS, Nickerson SR, Krysan DJ, Cullen PJ. Cleavage of the signaling mucin Msb2 by the aspartyl protease Yps1 is required for MAPK activation in yeast. *J Cell Biol.* 2008;181(7):1073-81. Epub 2008/07/02. doi: 10.1083/jcb.200704079. PubMed PMID: 18591427; PubMed Central PMCID: PMC2442203.
13. Pitoniak A, Birkaya B, Dionne HM, Vadaie N, Cullen PJ. The signaling mucins Msb2 and Hkr1 differentially regulate the filamentation mitogen-activated protein kinase pathway and contribute to a multimodal response. *Mol Biol Cell.* 2009;20(13):3101-14. Epub 2009/05/15. doi: 10.1091/mbc.E08-07-0760. PubMed PMID: 19439450; PubMed Central PMCID: PMC2704161.
14. O'Rourke SM, Herskowitz I. A Third Osmosensing Branch in *Saccharomyces cerevisiae* Requires the Msb2 Protein and Functions in Parallel with the Sho1 Branch. *Molecular and Cellular Biology.* 2002;22(13):4739-49. doi: 10.1128/mcb.22.13.4739-4749.2002.
15. Tatebayashi K, Tanaka K, Yang HY, Yamamoto K, Matsushita Y, Tomida T, et al. Transmembrane mucins Hkr1 and Msb2 are putative osmosensors in the SHO1 branch of yeast HOG pathway. *EMBO J.* 2007;26(15):3521-33. Epub 2007/07/14. doi: 10.1038/sj.emboj.7601796. PubMed PMID: 17627274; PubMed Central PMCID: PMC1949007.
16. Nemecek JC, Wuthrich M, Klein BS. Global control of dimorphism and virulence in fungi. *Science.* 2006;312(5773):583-8. doi: 10.1126/science.1124105. PubMed PMID: 16645097.
17. Gilmore SA, Naseem S, Konopka JB, Sil A. N-acetylglucosamine (GlcNAc) triggers a rapid, temperature-responsive morphogenetic program in thermally dimorphic fungi. *PLoS*

988 Genet. 2013;9(9):e1003799. Epub 2013/09/27. doi: 10.1371/journal.pgen.1003799. PubMed  
989 PMID: 24068964; PubMed Central PMCID: PMC3778022.

990 18. Wieser J, Lee BN, Fondon J, 3rd, Adams TH. Genetic requirements for initiating asexual  
991 development in *Aspergillus nidulans*. Curr Genet. 1994;27(1):62-9. Epub 1994/12/01. PubMed  
992 PMID: 7750148.

993 19. Tian X, Shearer G, Jr. Cloning and analysis of mold-specific genes in the dimorphic  
994 fungus *Histoplasma capsulatum*. Gene. 2001;275(1):107-14. Epub 2001/09/28. PubMed PMID:  
995 11574158.

996 20. Tian X, Shearer G. The Mold-Specific MS8 Gene Is Required for Normal Hypha  
997 Formation in the Dimorphic Pathogenic Fungus *Histoplasma capsulatum*. Eukaryotic Cell.  
998 2002;1(2):249-56. doi: 10.1128/ec.1.2.249-256.2002.

999 21. Patel JB, Batanghari JW, Goldman WE. Probing the yeast phase-specific expression of  
1000 the CBP1 gene in *Histoplasma capsulatum*. J Bacteriol. 1998;180(7):1786-92. Epub 1998/04/16.  
1001 PubMed PMID: 9537376; PubMed Central PMCID: PMC3778022.

1002 22. Isaac DT, Berkes CA, English BC, Murray DH, Lee YN, Coady A, et al. Macrophage  
1003 cell death and transcriptional response are actively triggered by the fungal virulence factor Cbp1  
1004 during *H. capsulatum* infection. Mol Microbiol. 2015;98(5):910-29. Epub 2015/08/20. doi:  
1005 10.1111/mmi.13168. PubMed PMID: 26288377; PubMed Central PMCID: PMC3778022.

1006 23. English BC, Van Prooyen N, Ord T, Ord T, Sil A. The transcription factor CHOP, an  
1007 effector of the integrated stress response, is required for host sensitivity to the fungal intracellular  
1008 pathogen *Histoplasma capsulatum*. PLoS Pathog. 2017;13(9):e1006589. Epub 2017/09/28. doi:  
1009 10.1371/journal.ppat.1006589. PubMed PMID: 28953979; PubMed Central PMCID:  
1010 PMC3778022.

1011 24. Batanghari JW, Deepe GS, Jr., Di Cera E, Goldman WE. Histoplasma acquisition of  
1012 calcium and expression of CBP1 during intracellular parasitism. *Mol Microbiol.* 1998;27(3):531-  
1013 9. Epub 1998/03/07. PubMed PMID: 9489665.

1014 25. Shen Q, Beucler MJ, Ray SC, Rappleye CA. Macrophage activation by IFN-gamma  
1015 triggers restriction of phagosomal copper from intracellular pathogens. *PLoS Pathog.*  
1016 2018;14(11):e1007444. Epub 2018/11/20. doi: 10.1371/journal.ppat.1007444. PubMed PMID:  
1017 30452484.

1018 26. Isaac DT, Coady A, Van Prooyen N, Sil A. The 3-hydroxy-methylglutaryl coenzyme A  
1019 lyase HCL1 is required for macrophage colonization by human fungal pathogen *Histoplasma*  
1020 *capsulatum*. *Infect Immun.* 2013;81(2):411-20. Epub 2012/11/28. doi: 10.1128/IAI.00833-12.  
1021 PubMed PMID: 23184522; PubMed Central PMCID: PMC3553812.

1022 27. Abidi FE, Roh H, Keath EJ. Identification and characterization of a phase-specific,  
1023 nuclear DNA binding protein from the dimorphic pathogenic fungus *Histoplasma capsulatum*.  
1024 *Infect Immun.* 1998;66(8):3867-73. Epub 1998/07/23. PubMed PMID: 9673274; PubMed  
1025 Central PMCID: PMC108439.

1026 28. Chandrashekar R, Curtis KC, Rawot BW, Kobayashi GS, Weil GJ. Molecular cloning  
1027 and characterization of a recombinant *Histoplasma capsulatum* antigen for antibody-based  
1028 diagnosis of human histoplasmosis. *J Clin Microbiol.* 1997;35(5):1071-6. Epub 1997/05/01.  
1029 PubMed PMID: 9114383; PubMed Central PMCID: PMC232705.

1030 29. Holbrook ED, Kemski MM, Richer SM, Wheat LJ, Rappleye CA. Glycosylation and  
1031 immunoreactivity of the *Histoplasma capsulatum* Cfp4 yeast-phase exoantigen. *Infect Immun.*  
1032 2014;82(10):4414-25. Epub 2014/08/13. doi: 10.1128/IAI.01893-14. PubMed PMID: 25114108;  
1033 PubMed Central PMCID: PMC4187849.

- 1034 30. Lanver D, Mendoza-Mendoza A, Brachmann A, Kahmann R. Sho1 and Msb2-related  
1035 proteins regulate appressorium development in the smut fungus *Ustilago maydis*. *Plant Cell*.  
1036 2010;22(6):2085-101. Epub 2010/07/01. doi: 10.1105/tpc.109.073734. PubMed PMID:  
1037 20587773; PubMed Central PMCID: PMC2910971.
- 1038 31. Perez-Nadales E, Di Pietro A. The membrane mucin Msb2 regulates invasive growth and  
1039 plant infection in *Fusarium oxysporum*. *Plant Cell*. 2011;23(3):1171-85. Epub 2011/03/29. doi:  
1040 10.1105/tpc.110.075093. PubMed PMID: 21441438; PubMed Central PMCID:  
1041 PMCPMC3082261.
- 1042 32. Tanaka K, Tatebayashi K, Nishimura A, Yamamoto K, Yang HY, Saito H. Yeast  
1043 osmosensors Hkr1 and Msb2 activate the Hog1 MAPK cascade by different mechanisms. *Sci*  
1044 *Signal*. 2014;7(314):ra21. Epub 2014/02/27. doi: 10.1126/scisignal.2004780. PubMed PMID:  
1045 24570489.
- 1046 33. Yamamoto K, Tatebayashi K, Saito H. Binding of the Extracellular Eight-Cysteine Motif  
1047 of Opy2 to the Putative Osmosensor Msb2 Is Essential for Activation of the Yeast High-  
1048 Osmolarity Glycerol Pathway. *Mol Cell Biol*. 2015;36(3):475-87. Epub 2015/11/26. doi:  
1049 10.1128/MCB.00853-15. PubMed PMID: 26598606; PubMed Central PMCID:  
1050 PMCPMC4719420.
- 1051 34. Zuzuarregui A, Li T, Friedmann C, Ammerer G, Alepuz P. Msb2 is a Ste11 membrane  
1052 concentrator required for full activation of the HOG pathway. *Biochim Biophys Acta*.  
1053 2015;1849(6):722-30. Epub 2015/02/18. doi: 10.1016/j.bbagr.2015.02.001. PubMed PMID:  
1054 25689021.

1055 35. Brewster JL, Gustin MC. Positioning of cell growth and division after osmotic stress  
1056 requires a MAP kinase pathway. *Yeast*. 1994;10(4):425-39. Epub 1994/04/01. doi:  
1057 10.1002/yea.320100402. PubMed PMID: 7941729.

1058 36. Hernday AD, Lohse MB, Fordyce PM, Nobile CJ, DeRisi JL, Johnson AD. Structure of  
1059 the transcriptional network controlling white-opaque switching in *Candida albicans*. *Mol*  
1060 *Microbiol*. 2013;90(1):22-35. Epub 2013/07/17. doi: 10.1111/mmi.12329. PubMed PMID:  
1061 23855748; PubMed Central PMCID: PMC3888361.

1062 37. Leng P, Lee PR, Wu H, Brown AJ. Efg1, a morphogenetic regulator in *Candida albicans*,  
1063 is a sequence-specific DNA binding protein. *J Bacteriol*. 2001;183(13):4090-3. Epub  
1064 2001/06/08. doi: 10.1128/JB.183.13.4090-4093.2001. PubMed PMID: 11395474; PubMed  
1065 Central PMCID: PMC95293.

1066 38. Noble SM, Gianetti BA, Witchley JN. *Candida albicans* cell-type switching and  
1067 functional plasticity in the mammalian host. *Nat Rev Microbiol*. 2017;15(2):96-108. doi:  
1068 10.1038/nrmicro.2016.157. PubMed PMID: 27867199; PubMed Central PMCID:  
1069 PMC5957277.

1070 39. Si H, Hernday AD, Hirakawa MP, Johnson AD, Bennett RJ. *Candida albicans* white and  
1071 opaque cells undergo distinct programs of filamentous growth. *PLoS Pathog*.  
1072 2013;9(3):e1003210. doi: 10.1371/journal.ppat.1003210. PubMed PMID: 23505370; PubMed  
1073 Central PMCID: PMC3591317.

1074 40. Sohn K, Urban C, Brunner H, Rupp S. EFG1 is a major regulator of cell wall dynamics in  
1075 *Candida albicans* as revealed by DNA microarrays. *Mol Microbiol*. 2003;47(1):89-102. Epub  
1076 2002/12/21. PubMed PMID: 12492856.



41. Stoldt VR, Sonneborn A, Leuker CE, Ernst JF. Efg1p, an essential regulator of morphogenesis of the human pathogen *Candida albicans*, is a member of a conserved class of bHLH proteins regulating morphogenetic processes in fungi. *EMBO J.* 1997;16(8):1982-91. Epub 1997/04/15. doi: 10.1093/emboj/16.8.1982. PubMed PMID: 9155024; PubMed Central PMCID: PMC1169801.
42. Miller KY, Toennis TM, Adams TH, Miller BL. Isolation and transcriptional characterization of a morphological modifier: the *Aspergillus nidulans* stunted (*stuA*) gene. *Mol Gen Genet.* 1991;227(2):285-92. Epub 1991/06/01. PubMed PMID: 2062309.
43. Miller KY, Wu J, Miller BL. StuA is required for cell pattern formation in *Aspergillus*. *Genes Dev.* 1992;6(9):1770-82. Epub 1992/09/01. PubMed PMID: 1516832.
44. Longo LVG, Ray SC, Puccia R, Rappleye CA. Characterization of the APSES-family transcriptional regulators of *Histoplasma capsulatum*. *FEMS Yeast Res.* 2018. Epub 2018/08/14. doi: 10.1093/femsyr/foy087. PubMed PMID: 30101348.
45. Yang E, Wang G, Woo PC, Lau SK, Chow WN, Chong KT, et al. Unraveling the molecular basis of temperature-dependent genetic regulation in *Penicillium marneffei*. *Eukaryot Cell.* 2013;12(9):1214-24. doi: 10.1128/EC.00159-13. PubMed PMID: 23851338; PubMed Central PMCID: PMC3811563.
46. Munoz JF, Gauthier GM, Desjardins CA, Gallo JE, Holder J, Sullivan TD, et al. The Dynamic Genome and Transcriptome of the Human Fungal Pathogen *Blastomyces* and Close Relative *Emmonsia*. *PLoS Genet.* 2015;11(10):e1005493. doi: 10.1371/journal.pgen.1005493. PubMed PMID: 26439490; PubMed Central PMCID: PMC4595289.
47. Azadmanesh J, Gowen AM, Creger PE, Schafer ND, Blankenship JR. Filamentation Involves Two Overlapping, but Distinct, Programs of Filamentation in the Pathogenic Fungus

1100 *Candida albicans*. G3 (Bethesda). 2017;7(11):3797-808. doi: 10.1534/g3.117.300224. PubMed  
1101 PMID: 28951491; PubMed Central PMCID: PMC5677161.

1102 48. Comeau AM, Dufour J, Bouvet GF, Jacobi V, Nigg M, Henrissat B, et al. Functional  
1103 annotation of the *Ophiostoma novo-ulmi* genome: insights into the phytopathogenicity of the  
1104 fungal agent of Dutch elm disease. *Genome Biol Evol*. 2014;7(2):410-30. doi:  
1105 10.1093/gbe/evu281. PubMed PMID: 25539722; PubMed Central PMCID: PMC50166.

1106 49. Wang A, Raniga PP, Lane S, Lu Y, Liu H. Hyphal chain formation in *Candida albicans*:  
1107 Cdc28-Hgc1 phosphorylation of Efg1 represses cell separation genes. *Mol Cell Biol*.  
1108 2009;29(16):4406-16. doi: 10.1128/MCB.01502-08. PubMed PMID: 19528234; PubMed Central  
1109 PMCID: PMC2725740.

1110 50. Sil A, Andrianopoulos A. Thermally Dimorphic Human Fungal Pathogens--Polyphyletic  
1111 Pathogens with a Convergent Pathogenicity Trait. *Cold Spring Harb Perspect Med*.  
1112 2014;5(8):a019794. Epub 2014/11/12. doi: 10.1101/cshperspect.a019794. PubMed PMID:  
1113 25384771; PubMed Central PMCID: PMC4526722.

1114 51. Anderson MZ, Porman AM, Wang N, Mancera E, Huang D, Cuomo CA, et al. A  
1115 Multistate Toggle Switch Defines Fungal Cell Fates and Is Regulated by Synergistic Genetic  
1116 Cues. *PLoS Genet*. 2016;12(10):e1006353. doi: 10.1371/journal.pgen.1006353. PubMed PMID:  
1117 27711197; PubMed Central PMCID: PMC5053522.

1118 52. Zordan RE, Miller MG, Galgoczy DJ, Tuch BB, Johnson AD. Interlocking transcriptional  
1119 feedback loops control white-opaque switching in *Candida albicans*. *PLoS Biol*.  
1120 2007;5(10):e256. doi: 10.1371/journal.pbio.0050256. PubMed PMID: 17880264; PubMed  
1121 Central PMCID: PMC1976629.

- 1122 53. Sriram K, Soliman S, Fages F. Dynamics of the interlocked positive feedback loops  
1123 explaining the robust epigenetic switching in *Candida albicans*. *J Theor Biol.* 2009;258(1):71-88.  
1124 doi: 10.1016/j.jtbi.2009.01.008. PubMed PMID: 19490874.
- 1125 54. Saraswat D, Kumar R, Pande T, Edgerton M, Cullen PJ. Signalling mucin Msb2  
1126 Regulates adaptation to thermal stress in *Candida albicans*. *Mol Microbiol.* 2016;100(3):425-41.  
1127 Epub 2016/01/11. doi: 10.1111/mmi.13326. PubMed PMID: 26749104; PubMed Central  
1128 PMCID: PMCPMC4955288.
- 1129 55. Cullen PJ. Post-translational regulation of signaling mucins. *Curr Opin Struct Biol.*  
1130 2011;21(5):590-6. doi: 10.1016/j.sbi.2011.08.007. PubMed PMID: 21889329; PubMed Central  
1131 PMCID: PMCPMC3189326.
- 1132 56. Martin R, Albrecht-Eckardt D, Brunke S, Hube B, Hunniger K, Kurzai O. A core  
1133 filamentation response network in *Candida albicans* is restricted to eight genes. *PLoS One.*  
1134 2013;8(3):e58613. doi: 10.1371/journal.pone.0058613. PubMed PMID: 23516516; PubMed  
1135 Central PMCID: PMCPMC3597736.
- 1136 57. Fang W, St Leger RJ. RNA binding proteins mediate the ability of a fungus to adapt to  
1137 the cold. *Environ Microbiol.* 2010;12(3):810-20. doi: 10.1111/j.1462-2920.2009.02127.x.  
1138 PubMed PMID: 20050869.
- 1139 58. Kim YO, Kang H. The role of a zinc finger-containing glycine-rich RNA-binding protein  
1140 during the cold adaptation process in *Arabidopsis thaliana*. *Plant Cell Physiol.* 2006;47(6):793-8.  
1141 doi: 10.1093/pcp/pcj047. PubMed PMID: 16608866.
- 1142 59. Olgeiser L, Haag C, Boerner S, Ule J, Busch A, Koepke J, et al. The key protein of  
1143 endosomal mRNP transport Rrm4 binds translational landmark sites of cargo mRNAs. *EMBO*

1144 Rep. 2019;20(1). doi: 10.15252/embr.201846588. PubMed PMID: 30552148; PubMed Central  
1145 PMCID: PMCPMC6322384.

1146 60. Worsham PL, Goldman WE. Quantitative plating of *Histoplasma capsulatum* without  
1147 addition of conditioned medium or siderophores. *Journal of medical and veterinary mycology* :  
1148 bi-monthly publication of the International Society for Human and Animal Mycology.  
1149 1988;26(3):137-43. Epub 1988/06/01. PubMed PMID: 3171821.

1150 61. Bryksin AV, Matsumura I. Overlap extension PCR cloning: a simple and reliable way to  
1151 create recombinant plasmids. *Biotechniques*. 2010;48(6):463-5. Epub 2010/06/24. doi:  
1152 10.2144/000113418. PubMed PMID: 20569222; PubMed Central PMCID: PMCPMC3121328.

1153 62. Kujoth GC, Sullivan TD, Merkhofer R, Lee TJ, Wang H, Brandhorst T, et al.  
1154 CRISPR/Cas9-Mediated Gene Disruption Reveals the Importance of Zinc Metabolism for  
1155 Fitness of the Dimorphic Fungal Pathogen *Blastomyces dermatitidis*. *MBio*. 2018;9(2). doi:  
1156 10.1128/mBio.00412-18. PubMed PMID: 29615501; PubMed Central PMCID:  
1157 PMCPMC5885028.

1158 63. Brinkman EK, Chen T, Amendola M, van Steensel B. Easy quantitative assessment of  
1159 genome editing by sequence trace decomposition. *Nucleic Acids Res*. 2014;42(22):e168. doi:  
1160 10.1093/nar/gku936. PubMed PMID: 25300484; PubMed Central PMCID: PMCPMC4267669.

1161 64. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment  
1162 of short DNA sequences to the human genome. *Genome Biol*. 2009;10(3):R25. doi: 10.1186/gb-  
1163 2009-10-3-r25. PubMed PMID: 19261174; PubMed Central PMCID: PMCPMC2690996.

1164 65. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or  
1165 without a reference genome. *BMC Bioinformatics*. 2011;12:323. Epub 2011/08/06. doi:

1166 10.1186/1471-2105-12-323. PubMed PMID: 21816040; PubMed Central PMCID:  
1167 PMCPMC3163565.

1168 66. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq  
1169 quantification. Nat Biotechnol. 2016;34(5):525-7. Epub 2016/04/05. doi: 10.1038/nbt.3519.  
1170 PubMed PMID: 27043002.

1171 67. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential  
1172 expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res.  
1173 2015;43(7):e47. Epub 2015/01/22. doi: 10.1093/nar/gkv007. PubMed PMID: 25605792;  
1174 PubMed Central PMCID: PMCPMC4402510.

1175 68. Smyth GK. Linear models and empirical bayes methods for assessing differential  
1176 expression in microarray experiments. Stat Appl Genet Mol Biol. 2004;3:Article3. Epub  
1177 2006/05/02. doi: 10.2202/1544-6115.1027. PubMed PMID: 16646809.

1178 69. Law CW, Chen Y, Shi W, Smyth GK. voom: Precision weights unlock linear model  
1179 analysis tools for RNA-seq read counts. Genome Biol. 2014;15(2):R29. Epub 2014/02/04. doi:  
1180 10.1186/gb-2014-15-2-r29. PubMed PMID: 24485249; PubMed Central PMCID:  
1181 PMCPMC4053721.

1182 70. Remm M, Storm CE, Sonnhammer EL. Automatic clustering of orthologs and in-  
1183 paralogs from pairwise species comparisons. J Mol Biol. 2001;314(5):1041-52. doi:  
1184 10.1006/jmbi.2000.5197. PubMed PMID: 11743721.

1185 71. Do CB, Mahabhashyam MS, Brudno M, Batzoglou S. ProbCons: Probabilistic  
1186 consistency-based multiple sequence alignment. Genome Res. 2005;15(2):330-40. doi:  
1187 10.1101/gr.2821705. PubMed PMID: 15687296; PubMed Central PMCID: PMCPMC546535.

1188 72. Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood trees for  
1189 large alignments. PLoS One. 2010;5(3):e9490. doi: 10.1371/journal.pone.0009490. PubMed  
1190 PMID: 20224823; PubMed Central PMCID: PMC2835736.

1191 73. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time  
1192 quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods. 2001;25(4):402-8. doi:  
1193 10.1006/meth.2001.1262. PubMed PMID: 11846609.

1194 74. Team RC. R: A language and environment for statistical computing. Vienna,  
1195 Austria 2015.

1196 75. Van der Walt S, Colbert SC, Varoquaux G. The NumPy Array: A Structure for Efficient  
1197 Numerical Computation. Computing in Science and Engineering. 2011;13:22-30.

1198 76. Hunter JD. Matplotlib: A 2D Graphics Environment. Computing in Science &  
1199 Engineering. Computing in Science and Engineering. 2007;9(3):90-5.

1200 77. Perez F, Granger BE. IPython: A System for Interactive Scientific Computing.  
1201 Computing in Science and Engineering. 2007;9:21-9.

1202 78. Saldanha AJ. Java Treeview--extensible visualization of microarray data. Bioinformatics.  
1203 2004;20(17):3246-8. doi: 10.1093/bioinformatics/bth349. PubMed PMID: 15180930.

1204






Figure 1










Figure 2







Figure 3




A


bioRxiv preprint doi: <https://doi.org/10.1101/546853>; this version posted February 11, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.



B



C



D




Figure 4




Figure 5






Figure 6





Figure 7

# Growth curves





Temp: 37°C

RT

Timepoint: day 0

day 2

day 6

day 8

WT

*msb2*

*msb2+MSB2*

