

1 DNA barcode reference libraries for the monitoring 2 of aquatic biota in Europe: Gap-analysis and 3 recommendations for future work

4
5
6 Hannah Weigand¹, Arne J. Beermann², Fedor Čiampor³, Filipe O. Costa^{4,5}, Zoltán Csabai⁶,
7 Sofia Duarte^{4,5}, Matthias F. Geiger⁷, Michał Grabowski⁸, Frédéric Rimet⁹, Björn Rulik⁷, Malin
8 Strand¹⁰, Nikolaus Szucsich¹¹, Alexander M. Weigand^{1,2}, Endre Willassen¹², Sofia A. Wyler¹³,
9 Agnès Bouchez⁹, Angel Borja¹⁴, Zuzana Čiamporová-Zaťovičová³, Sónia Ferreira¹⁵, KD
10 Dijkstra¹⁶, Ursula Eisendle¹⁷, Jörg Freyhof¹⁸, Piotr Gadawski⁸, Wolfram Graf¹⁹, Arne
11 Haegerbaeumer²⁰, Berry B. van der Hoorn¹⁶, Bella Japoshvili²¹, Lujza Keresztes²², Emre
12 Keskin²³, Florian Leese², Jan Macher¹⁶, Tomasz Mamos⁸, Guy Paz²⁴, Vladimir Pešić²⁵,
13 Daniela Maric Pfannkuchen²⁶, Martin Andreas Pfannkuchen²⁶, Benjamin W. Price²⁷, Buki
14 Rinkevich²⁴, Marcos A. L. Teixeira^{4,5}, Gábor Várbíró²⁸, Torbjørn Ekrem^{29*}

15

16 *Corresponding author

17

18 Affiliations

19 ¹Musée National d'Histoire Naturelle, 25 Rue Münster, 2160 Luxembourg, Luxembourg. E-
20 mail: hannah.weigand@mnhn.lu, alexander.weigand@mnhn.lu

21 ²University of Duisburg-Essen, Faculty of Biology, Aquatic Ecosystem Research,
22 Universitaetsstr. 5, 45141 Essen, Germany. E-mail: arne.beermann@uni-due.de,
23 florian.leese@uni-due.de

24 ³Slovak Academy of Sciences, Plant Science and Biodiversity Centre, Zoology Lab,
25 Dúbravská cesta 9, 84523, Bratislava, Slovakia. E-mail: f.ciampor@savba.sk,
26 zuzana.zatovicova@savba.sk

27 ⁴Centre of Molecular and Environmental Biology (CBMA), University of Minho, Campus de
28 Gualtar, 4710-057 Braga, Portugal.

29 ⁵Institute of Science and Innovation for Bio-Sustainability (IB-S), University of Minho,
30 Campus de Gualtar, 4710-057 Braga, Portugal. E-mail: fcosta@bio.uminho.pt,
31 sduarte@bio.uminho.pt, mark-us_teixeira@hotmail.com

32 ⁶University of Pécs, Faculty of Sciences, Department of Hydrobiology, Ifjúság útja 6, H7624
33 Pécs, Hungary. E-mail: csabai@gamma.ttk.pte.hu

34 ⁷Zoologisches Forschungsmuseum Alexander Koenig, Leibniz Institute for Animal
35 Biodiversity, Adenauerallee 160, 53113 Bonn, Germany. E-mail: m.geiger@leibniz-zfmk.de,
36 B.Rulik@leibniz-zfmk.de

37 ⁸University of Lodz, Faculty of Biology and Environmental Protection, Department of
38 Invertebrate Zoology and Hydrobiology, Banacha 12/16, 90-237 Łódź, Poland. E-mail:
39 michal.grabowski@biol.uni.lodz.pl, tomasz.mamos@biol.uni.lodz.pl,
40 piotr.gadawski@biol.uni.lodz.pl

41 ⁹INRA, Université Savoie Mont Blanc, UMR Carrel, FR-74200 Thonon-les-Bains, France.
42 Email: frederic.rimet@inra.fr, agnes.bouchez@inra.fr

43 ¹⁰Swedish University of Agricultural Sciences, Swedish Species Information Centre, Uppsala,
44 Sweden. E-mail: Malin.Strand@slu.se

45 ¹¹Natural History Museum Vienna, Burgring 7, 1010 Vienna, Austria. E-mail:
46 nikolaus.szucsich@nhm-wien.ac.at

47 ¹²University of Bergen, University Museum of Bergen, NO-5007 Bergen, Norway. E-mail:
48 endre.willassen@uib.no

49 ¹³info fauna - Centre Suisse de Cartographie de la Faune (CSCF), Avenue de Bellevaux 51,
50 2000 Neuchâtel, Switzerland. E-mail: sofia.wyler@unine.ch

51 ¹⁴AZTI – Marine Research Division, Herrera Kaia, Portualdea z/g, 20110 Pasaia, Gipuzkoa,
52 Spain, Spain. E-mail: aborja@azti.es

53 ¹⁵CIBIO/InBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos,
54 Universidade do Porto, Campus Agrário de Vairão, 4485-661 Vairão, Portugal. E-mail:
55 hiporame@gmail.com

56 ¹⁶Naturalis Biodiversity Center, PO box 9517, 2300 RA Leiden, the Netherlands. E-mail:
57 kd.dijkstra@naturalis.nl, berry.vanderhoorn@naturalis.nl, jan.macher@naturalis.nl

58 ¹⁷University of Salzburg, Department of Biosciences, Hellbrunnerstraße 34, 5020 Salzburg,
59 Austria. E-mail: Ursula.Eisendle@sbg.ac.at

60 ¹⁸Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), 12587 Berlin, Germany.
61 E-mail: j.freyhof@igb-berlin.de

62 ¹⁹University of Natural Resources and Life Sciences, Vienna, Institute of Hydrobiology and
63 Aquatic Ecosystem Management (IHG), Gregor-Mendel-Straße 33/DG, 1180 Vienna,
64 Austria. E-mail: wolfram.graf@boku.ac.at

65 ²⁰Bielefeld University, Department of Animal Ecology, Konsequenz 45, 33615 Bielefeld,
66 Germany. E-mail: a.haegerbaeumer@uni-bielefeld.de

67 ²¹Ilia State University, Institute of Zoology, 3/5 Cholokashvili ave, 0179, Tbilisi, Georgia. E-
68 mail: bela_japoshvili@iliauni.edu.ge

69 ²²Babeş-Bolyai University, Faculty of Biology and Geology, Center of Systems Biology,
70 Biodiversity and Bioresources, Cliniclor 5-7, 400006, Cluj Napoca, Romania. E-mail:
71 keresztes2012@gmail.com

72 ²³Ankara University, Agricultural Faculty, Department of Fisheries and Aquaculture,
73 Evolutionary Genetics Laboratory (eGL), Ankara, Turkey. E-mail: keskin@ankara.edu.tr

74 ²⁴Israel Oceanographic and Limnological Research, National Institute of Oceanography,
75 Haifa 31080, Israel. E-mail: buki@ocean.org.il, guy@ocean.org.il

76 ²⁵University of Montenegro, Department of Biology, Cetinjski put bb., 20000 Podgorica,
77 Montenegro. E-mail: vladopesic@gmail.com

78 ²⁶Rudjer Boskovic Institute, Center for marine research, G. Paliaga 5, Rovinj, Croatia. E-mail:
79 dmaric@irb.hr, Pfannkuchen@cim.irb.hr

80 ²⁷Natural History Museum, Cromwell Road, London, UK. E-mail: b.price@nhm.ac.uk

81 ²⁸MTA Centre for Ecological Research, Danube Research Institute, Department of Tisza
82 River Research, Bem square 18/C, H4026 Debrecen, Hungary. E-mail:
83 varbiro.gabor@okologia.mta.hu

84 ²⁹Norwegian University of Science and Technology, NTNU University Museum, Department
85 of Natural History, NO-7491 Trondheim, Norway. E-mail: torbjorn.ekrem@ntnu.no

86

87 **Conflict of interest**

88 The authors declare no conflict of interest.

89

90 **Author Contributions**

91 All authors contributed to conceptualization and review of the manuscript.
92 HW, AJB, FC, FOC, ZC, SD, MFG, MG, FR, BRu, MS, NS, AMW, EW, SAW, ZC-Z, SF, UE,
93 JF, PG, WG, AH, BBH, BJ, JM, TM, GP, MAP, BWP, BRI, MALT, GV and TE provided and
94 validated data.

95 HW analysed data.

96 TE, HW, FOC, MFG, AMW, SD, MG, FR, FC, MS, SAW, ZC, AJB and BRu drafted the
97 manuscript.

98 TE and FC administered the project and coordinated workshops.

99

100 Abstract

101 Effective identification of species using short DNA fragments (DNA barcoding and DNA
102 metabarcoding) requires reliable sequence reference libraries of known taxa. Both
103 taxonomically comprehensive coverage and content quality are important for sufficient
104 accuracy. For aquatic ecosystems in Europe, reliable barcode reference libraries are
105 particularly important if molecular identification tools are to be implemented in biomonitoring
106 and reports in the context of the EU Water Framework Directive (WFD) and the Marine
107 Strategy Framework Directive (MSFD). We analysed gaps in the two most important
108 reference databases, Barcode of Life Data Systems (BOLD) and NCBI GenBank, with a
109 focus on the taxa most frequently used in WFD and MSFD. Our analyses show that
110 coverage varies strongly among taxonomic groups, and among geographic regions. In
111 general, groups that were actively targeted in barcode projects (e.g. fish, true bugs,
112 caddisflies and vascular plants) are well represented in the barcode libraries, while others
113 have fewer records (e.g. marine molluscs, ascidians, and freshwater diatoms). We also
114 found that species monitored in several countries often are represented by barcodes in
115 reference libraries, while species monitored in a single country frequently lack sequence
116 records. A large proportion of species (up to 50%) in several taxonomic groups are only
117 represented by private data in BOLD. Our results have implications for the future strategy to
118 fill existing gaps in barcode libraries, especially if DNA metabarcoding is to be used in the
119 monitoring of European aquatic biota under the WFD and MSFD. For example, missing
120 species relevant to monitoring in multiple countries should be prioritized. We also discuss
121 why a strategy for quality control and quality assurance of barcode reference libraries is
122 needed and recommend future steps to ensure full utilization of metabarcoding in aquatic
123 biomonitoring.

124

125 Keywords: DNA barcoding; reference library; biological monitoring; freshwater; marine;
126 quality assurance

127

128 1. Introduction

129 1.1 DNA barcoding for monitoring aquatic life

130 Aquatic life is of central importance to human well-being and essential for our understanding
131 of natural history, evolution and ecology. From the deepest oceans to the highest peaks, life
132 in water characterizes environmental conditions, and constitutes invaluable ecosystem
133 functions with services for a wide array of communities (Borgwardt et al., 2019; Rouillard et
134 al., 2018). For these reasons, our ability to assess aquatic biodiversity and monitor its
135 change over time is of great significance, not only to prevent biodiversity loss, but to ensure
136 our own welfare.

137 The world's oceans cover 70% of the Earth's surface and are home to approximately
138 242,000 described species (Horton et al., 2018). It is estimated, however, that 91% of
139 eukaryotic marine life is undescribed, and that the total number of marine species is around
140 2.2 million (Mora et al., 2011). More than one third of the world's human population lives in
141 the coastal zone, and ecosystem services provided by the marine environment are both
142 crucial to human well-being and affected by our activities (Barbier, 2012; Barbier, 2017). In
143 Europe, the Marine Strategy Framework Directive (MSFD, Directive 2008/56/EC) aims to
144 achieve "good environmental status" of marine waters by 2020 and to protect marine
145 environments in the European Union (European Commission, 2008). The MSFD includes a
146 wide array of requirements in its ecosystem-based approach for assessment and monitoring,
147 including information on animal and plant communities (Borja et al., 2013). A large
148 percentage of undescribed biota certainly hampers community comparisons among sites and
149 regions, and likely restrains the explanatory power of marine water quality indices (Aylagas
150 et al., 2014).

151 Although representing only 0.01% of the Earth's water, freshwater ecosystems hold about
152 6% of all described species (Balian et al., 2008; Dudgeon et al., 2006; Reid et al., 2018).
153 Freshwater represents a valuable and irreplaceable natural resource, and scarcity as well as
154 quality are likely to continue to affect the stability of human communities (Kreamer, 2012).
155 Four-fifths of the world's population now lives in areas where there is a threat to water
156 security (UN World Water Assessment Programme, 2018), and it is estimated that demand
157 for freshwater will increase by 20-30% by 2050 (Burek et al., 2016). Water quality as well as
158 access to water is of global concern, and nature-based solutions have received increased
159 attention as ways of improving water quality (UN World Water Assessment Programme,
160 2018). In Europe, assessments of water quality have been a hot topic for decades (Birk et
161 al., 2012b; Hering et al., 2010; Leese et al., 2018; Metcalfe, 1989), and the use of
162 biodiversity estimates for this purpose is central in the EU Water Framework Directive (WFD)

163 (European Commission, 2000). Moreover, the highest proportion of species extinctions to
164 date has been recorded in freshwater (Young et al., 2016), highlighting the importance of
165 monitoring and protecting these ecosystems.

166

167 Thus, together with the Groundwater Directive (GWD, Directive 2006/118/EC) and the
168 Habitats Directive (Directive 92/43/EEC), the WFD and MSFD make water quality monitoring
169 of Europe's aquatic environments legally binding in all EU member states, Norway, Iceland
170 and Switzerland.

171 However, among countries there are large differences in the way biodiversity data are used
172 to assess aquatic ecosystem quality status (Birk et al., 2012a; Kelly et al., 2015): different
173 indices, different taxonomic groups, and different taxonomic levels are applied. Despite
174 differences in methodology, the goals are similar and focus on the quantification of
175 environmental states in comparison with reference conditions. Protocols and assessment
176 metrics applied have undergone a sophisticated intercalibration procedure to harmonise data
177 among countries and make ecological status assessments comparable.

178 To assess the ecological status, identification of aquatic organisms to family, genus or
179 species-level by morphology is necessary, but it is not a straightforward process. For
180 instance, individual differences in expertise, experience and opinion of the identifiers can
181 result in different taxonomic groups being documented from the same waterbody, potentially
182 leading to contrasting ecological assessments (Carstensen and Lindegarth, 2016; Clarke,
183 2013). An extensive audit of 414 macroinvertebrate samples taken as part of the monitoring
184 programs of German rivers and streams (Haase et al., 2010) documented that 29% of the
185 specimens had been overlooked by the primary analyst in the sorting stage, and that the
186 identification of >30% of the taxa differed between the primary analyst and the auditors.

187 Importantly, these results lead to divergent ecological assessments in 16% of the samples
188 (Haase et al., 2010). Similar studies have been performed in Norway and Finland (Meissner
189 et al., 2012; Meissner et al., 2017; Petrin et al., 2016) with comparable results. Despite the
190 general challenges in using short, standardized molecular markers for identification (Hebert
191 et al., 2016), DNA barcoding and metabarcoding offer a less subjective approach than
192 morphology for the identification of organisms in aquatic assessments (Leese et al., 2018).

193 Some issues still need to be solved and standard protocols to be developed before DNA
194 metabarcoding becomes the method of choice in aquatic biomonitoring. The use of both
195 organismal and environmental DNA (eDNA) in nature management decisions is already
196 being tested in some European countries (Hering et al., 2018), and the genetic water quality
197 index recently developed for marine waters (gAMBI) is performing well (Aylagas et al., 2018;
198 Aylagas et al., 2014). The EU COST Action DNAqua-Net (CA15219) was initiated with the

199 purpose of developing genetic tools for bioassessments of aquatic ecosystems in Europe
200 (Leese et al., 2016). The network aims to evaluate existing methods and reference libraries,
201 as well as to develop protocols and good practices in the use of DNA-based monitoring and
202 assessments of aquatic habitats. By connecting scientists and stakeholders, DNAqua-Net so
203 far has been a successful platform for this purpose.

204 Comprehensive DNA barcode reference libraries, such as the Barcode of Life Data Systems
205 (BOLD (Ratnasingham and Hebert, 2007)) or GenBank (Benson et al., 2013), are essential
206 for biodiversity monitoring if one wishes to utilise species' autecological and biogeographic
207 information gathered during the last century and to compare results with previous
208 assessments. But also smaller, more taxon specific reference libraries, such as Diat.barcode
209 library, formerly called R-Syst::diatom database (Rimet et al., 2016) are important as these
210 might be easier to curate. Particularly in the current 'big biodiversity data' era, in which
211 hundreds of millions of sequences can be generated during a single high-throughput
212 sequencing run, we are no longer able to individually check sequence by sequence. It is thus
213 imperative that effective quality filtering processes are embedded, including that reference
214 libraries hold high standards and are well populated in order to trust (semi-)automated
215 taxonomic assignments (Brodin et al., 2012; Carew et al., 2017; Ekrem et al., 2007; Hebert et
216 al., 2003a; Mioduchowska et al., 2018; Porter and Hajibabaei, 2018). An elaborated quality
217 assurance / quality control (QA/QC) system can serve both purposes. Building barcode
218 libraries and associated voucher collections have therefore been major goals in individual
219 projects as well as national barcode campaigns over the last decade. In Europe, some
220 nations have been successful in obtaining funding to coordinate this work on a national level.
221 Others have contributed to reference libraries on a project-by-project basis. The way the
222 work on reference libraries has been organized is different between nations, and in some
223 cases decisive for which taxonomic groups and regions were covered. We therefore find it
224 informative and useful to briefly recapture the most important aspects of these initiatives in
225 Europe.

226 1.2 Barcode Campaigns in Europe

227 The Austrian Barcode of Life ([ABOL](#)) is an initiative with the main aim to generate and
228 provide DNA barcodes for all species of animals, plants and fungi recorded from Austria. The
229 main purpose of the pilot phase (2014-2017) was to build up a network of biodiversity experts
230 and conduct four pilot studies. Currently DNA barcodes are generated in a number of
231 independently funded projects. The pilot phase and the continued coordination of ABOL is
232 funded by the Ministry of Education, Science and Research and lies at the Natural History
233 Museum Vienna. Apart from building up the reference library, necessary for genetically

234 determining organisms, ABOL aims to stimulate biodiversity research by acquiring funds,
235 fostering diverse applications of DNA barcoding, building up and exchanging skills within the
236 network, and increasing public awareness for biodiversity.

237 The Finnish Barcode of Life ([FinBOL](#)) is a national project and a network of species experts
238 with the goal of creating DNA barcodes for all species of animals, plants and fungi occurring
239 in Finland. FinBOL has acted as a national node in the International Barcode of Life (iBOL)
240 project. FinBOL has been funded almost continuously from 2011 by several national funding
241 agencies. At the moment, FinBOL acts within the framework of the Finnish Biodiversity
242 Information Facility (FinBIF) and is coordinated by the University of Oulu. DNA barcoding
243 details for all Finnish species are provided in the [Laji.fi](#) portal, where progress is continuously
244 updated. At present, over 100,000 specimens stored in Finnish collections have been
245 subjected for barcoding, and DNA barcodes are available for about 20,000 species (~50%)
246 reported from Finland. In the near future, FinBOL aims at broadening the nationwide DNA
247 barcode reference library by adopting efficient high-throughput sequencing tools to recover
248 sequence information from older museum specimens.

249 Since November 2011, the German Federal Ministry of Education and Research (BMBF) is
250 funding a consortium of natural history museums and research institutions to set up the
251 'German Barcode of Life' initiative ([GBOL](#)). The main aim was to establish a network of
252 professionals and non-professionals to start with the construction of a DNA barcode
253 reference library for the fauna, flora and fungi of Germany. After the first phase (2011-2015)
254 a national web portal for DNA barcodes and specimen data was developed and is
255 continuously improved. It serves mainly the coordination of the collecting activities of over
256 250 scientists (amateurs and professionals) who provide their taxonomic expertise. In
257 addition, more than 50 institution-based taxonomists contribute to GBOL. Of the 48,000
258 animal and 10,000 plant species (excluding algae and fungi) present in Germany, over
259 23,000 different species have been processed and DNA barcodes for them generated. In
260 total, 295,000 specimens were submitted to GBOL institutes, and after choosing up to 10
261 individuals per species from throughout their distribution range in Germany, over 145,000 of
262 them delivered a DNA barcode. The second phase of GBOL (2016-2019) has focussed on
263 applications of DNA barcoding with dedicated PhD students working on specific aspects from
264 metabarcoding for water quality assessments to developing a diagnostic microarray chip for
265 the detection of phytopathogenic fungi. As a prerequisite for the successful implementation of
266 the new techniques a core team and network of taxonomists is further expanding the
267 reference library with DNA barcodes for another 13,800 species. With this target the
268 database will be filled with about half of the known metazoan species of German animals and
269 plants and be operable to identify the vast majority in terrestrial and aquatic environmental

270 samples. Substantial contributions to the reference library for German taxa came from the
271 project 'Barcode Fauna Bavarica (BFB)', which started in 2009 and is supported by grants
272 from the Bavarian State Government. The project focuses on animal biodiversity in Southern
273 Germany and is coordinated by the Bavarian State Collection of Zoology (ZSM). Research
274 activities involve close cooperation with the Biodiversity Institute of Ontario, which performs
275 the sequence analyses under the framework of the International Barcode of Life Project
276 ([iBOL](#)).

277 The Norwegian Barcode of Life Network ([NorBOL](#)) started in 2008 as a consortium of
278 biodiversity institutions in formal agreement of advancing DNA barcoding in Norway. The four
279 university museums in Bergen, Oslo, Tromsø and Trondheim have been hubs in the network
280 since then, and together with the Biodiversity Institute of Ontario, Canada, the main partners
281 in a national research infrastructure project that received funding from the Research Council
282 of Norway and the Norwegian Biodiversity Information Centre (NBIC) in 2014. The major
283 goal of the NorBOL-project was to database DNA barcodes of 20,000 Norwegian,
284 Scandinavian or Polar species in BOLD by the end of 2018. However, also knowledge
285 transfer, building expertise, and curation of specimen reference collections have been
286 important tasks of the network. Close collaboration with the Norwegian Taxonomy Initiative,
287 run by NBIC, has been crucial in this process as it has provided identified specimens of
288 many organism groups available for DNA analysis. Several applied research and
289 management projects have originated through collaboration in NorBOL.

290 The Swiss Barcode of Life ([SwissBOL](#)) is the national initiative for the creation of a genetic
291 catalogue for all species occurring in Switzerland. SwissBOL officially started in 2012
292 supported by the Federal Office for the Environment, with the goal of establishing a network
293 of scientists and institutions involved in the genetic inventory of Swiss biodiversity. During the
294 pilot phase (2012-2015), 24 targeted projects were developed on different taxonomic groups:
295 animals, plants, fungi, lichens and microorganisms. Ever since (transitory phase; 2016-
296 2018), the coordination of SwissBOL has been funded almost continuously, and data has
297 been acquired within only a few independently funded projects. In order to elaborate a
298 national strategy for the development of projects generating novel genetic data, a non-profit
299 association of experts was founded. Most recently, SwissBOL has been mostly working in
300 the development of the concepts for the genetic database with the major goal of ensuring
301 that the information related to the genetic data are accessible and linked together. The close
302 collaboration with the GBIF Swiss Node (<http://www.gbif.ch>) has been fundamental to ensure
303 the coherence of all the information provided with the standards defined at the national and
304 international levels.

305 The Netherlands started their barcoding initiative NBOL for plants and animals in 2008, led
306 by Naturalis Biodiversity Center in collaboration with a large number of Dutch NGOs and
307 over 50 amateur naturalists. A considerable starting grant from the national government in
308 2010 gave a tremendous boost to the DNA barcoding infrastructure at Naturalis and hence to
309 the national barcoding activities. So far, over 80,000 DNA barcodes have been generated.
310 More than half of the barcodes have been uploaded to BOLD. However, most of these
311 barcodes are still private because they are part of active research projects. Current
312 barcoding efforts focus on the completion of reference libraries of freshwater and marine
313 species (North Sea) for DNA-based biodiversity assessments, and are financed by private
314 funding organisations.

315 Among various DNA barcoding initiatives in Portugal, one of the most prominent
316 contributions has been provided by the network for barcoding marine life. This network was
317 activated in 2008 through a research grant (LusoMarBoL- Lusitanian Marine Barcode of Life)
318 from the national science funding body (Fundação para a Ciência e a Tecnologia - FCT), and
319 has been active ever since through subsequent research grants. Core reference libraries for
320 Portuguese marine life have been created, published and made available in BOLD, with
321 particular focus on marine fish (Costa et al., 2012; Oliveira et al., 2016), annelids (Lobo et al.,
322 2016), crustaceans (e.g. (Lobo et al., 2017) and molluscs (Borges et al., 2016).

323 While national DNA barcode initiatives often start opportunistically and register any species
324 available for sampling, focus shifts to fill the gaps of the databases as soon as a critical
325 number of species is registered. Which taxonomic groups have priority is typically connected
326 to funded projects, available taxonomic expertise and scientific collections, and is not
327 necessarily the same in each campaign. Among aquatic taxa, species-rich groups such as
328 arthropods and polychaetes, or economically important groups such as fish, have seen some
329 priority. However, when building barcode reference libraries, there has usually not been a
330 general focus on species or organisms that are particularly relevant for water quality
331 assessments towards WFD or MSFD from the start.

332 In addition to large national barcoding campaigns, smaller activities intended to generate
333 reference barcodes of selected taxonomic groups (e.g. Trichoptera Barcode of Life), or
334 regional biota (e.g. "Barcode Aquatic Biota of Slovakia - AquaBOL.sk" and "Israel marine
335 barcoding database") exist. These initiatives, even if lacking substantial funding, can provide
336 important data and in many cases be better targeted towards filling the gaps of barcode
337 libraries than more general campaigns.

338 1.3 Biological Quality Elements

339 Different organism groups are used as Biological Quality Elements (BQEs) to assess the
340 Ecological Quality Status (EQS) of aquatic ecosystems under the WFD. In the MSFD,
341 biodiversity data in general, along with other related descriptors, are used to define
342 Environmental Status (Borja et al., 2013; Zampoukas et al., 2014).

343 The MSFD is the first EU legislative instrument related to the protection of marine
344 biodiversity. The directive lists four European marine regions: 1) the Baltic Sea, 2) the North-
345 east Atlantic Ocean, 3) the Mediterranean Sea, and 4) the Black Sea. Member States of one
346 marine region and with neighbouring countries sharing the same marine waters, collaborate
347 in four Regional Sea Conventions (OSPAR¹, HELCOM², UNEP-MAP³ and the Bucharest
348 Convention⁴). These different regions naturally share, or aim to share, taxa/species lists for
349 biodiversity assessments and reporting status. The status is defined by eleven descriptors in
350 the MSFD (e.g. biological diversity, non-indigenous species, fishing, eutrophication, seafloor
351 integrity, etc.). For some descriptors, species ID is critical. National marine environmental
352 monitoring often focuses on regular sampling sites and observations of specific habitats and
353 its inhabitants, i.e. groups of organisms such as benthic macroinvertebrates, phytoplankton,
354 or fish. As already mentioned, there exist large differences between countries in how
355 biodiversity data are used to evaluate the quality status of aquatic ecosystems. This is
356 indeed true for the marine environment, and only few countries were able to support this
357 study with national taxalists directly associated to the MSFD. MSFD overlaps with WFD, and
358 in coastal waters MSFD is intended to apply to the aspects of *Good Environmental Status*
359 that are not covered by WFD (e.g. noise, litter, other aspects of biodiversity) (European
360 Commission, 2017). In order to perform barcode gap-analyses for taxa of relevance to the
361 directives and with a European marine perspective, we identified the possibilities of two
362 existing taxalists: AZTI's Marine Biotic Index (AMBI; (Borja et al., 2000)) and the European
363 Register of Marine Species ([ERMS](#)).

364 The AMBI is used as a component of the benthic invertebrates' assessment by several
365 Member States in the four regional seas (Borja et al., 2009; European Commission, 2018), in
366 the context of describing the sensitivity of macrobenthic species to both anthropogenic and

¹ Oslo/Paris Convention on the Protection of the Marine Environment of the North-East Atlantic
<https://www.ospar.org/convention>

² Helsinki Convention on the Protection of the Marine Environment of the Baltic Sea Area
<http://www.helcom.fi/>

³ United Nations Environment Programme - Mediterranean Action Plan to the Barcelona Convention
<http://web.unep.org/uneppmap/>

⁴ The Convention on the Protection of the Black Sea Against Pollution
http://www.blacksea-commission.org/_convention.asp

367 natural pressures (see e.g. (Borja et al., 2000)). The index uses the abundance weighted
368 average disturbance sensitivity of macroinvertebrate species in a sample (Borja et al., 2000),
369 each species being assigned to one of five ecological groups (EG I-V; (Grall and Glémarec,
370 1997). The AMBI list includes approximately 8,000 taxa (only macroinvertebrates) from all
371 seas, with representatives of the most important soft-bottom communities present at
372 estuarine and coastal systems, from the North Sea to the Mediterranean, North and South
373 America, Asia, etc. The second list used for the work is ERMS (Costello, 2000). This is a
374 taxonomic list of species occurring in the European marine environment, which includes the
375 continental shelf seas of Europe as well as the Mediterranean shelf, Baltic Seas and deep-
376 sea areas (<http://www.marbef.org/data/ermsmap.php>) up to the shoreline or splash zone
377 above the high tide mark and down to 0.5 psu salinity in estuaries. The register was founded
378 in 1998 by a grant from the EU's Marine Science and Technology Programme and contains
379 tens of thousands of marine species, so for this study we used a relevant selection of
380 organism groups within the register (see methods). In contrast to freshwater
381 microphytobenthos, where ecological indices are calculated on the base of country specific
382 index values attached to species names, marine microphytobenthos is not used for the
383 calculation of ecological indices. And while all four regional sea conventions recognize the
384 importance of marine microphytoplankton monitoring, no ecological index based on species-
385 specific values is implemented. Monitoring of marine microphytoplankton is therefore carried
386 out by monitoring the presence or abundance of all observable species as a biodiversity
387 measure with an additional focus on the search for invasive species. This approach
388 effectively extends the range of species monitored to the range of all known
389 microphytoplankton species as there is no restriction to a list of species with ecological index
390 values.

391 In freshwater, diatoms, with their huge species diversity, are particularly interesting
392 ecological indicators (Stevenson, 2014). They have been routinely used for monitoring of
393 surface waters for several decades (Rimet, 2012), and are required BQEs in assessments of
394 surface waters in Europe and the United States (Barbour and United States. Environmental
395 Protection Agency. Office of Water., 1999; European Commission, 2000). Until recently, the
396 standardized methodology for biological monitoring using diatoms was uniquely based on
397 microscopic determinations and counts (European Standard EN 14407:2014). This is quite
398 time-consuming and requires expertise in diatom taxonomy; skills that can only be acquired
399 after several months or years of practice. The development of high-throughput sequencing
400 (HTS) technologies and DNA barcoding provides an alternative to the tedious work of
401 morphological identification. The first proofs of concept, carried out on a few tens of samples,
402 showed interesting and encouraging results (Kermarrec et al., 2013; Zimmermann et al.,

403 2015). Recent studies confirmed that diatom indices obtained from DNA metabarcoding
404 provide very similar results to diatom indices calculated by microscopic counts, both on a
405 regional and national scale (Keck et al., 2017; Lefrancois et al., 2018; Rimet et al., 2018b;
406 Rivera et al., 2018a; Rivera et al., 2018b; Vasselon et al., 2018; Vasselon et al., 2017).
407 However, all these studies underlined the necessity of well-curated reference libraries. In
408 Europe, efforts to develop such a resource are made by a group of diatom experts, which
409 curate the Diat.barcode library (Rimet et al., 2016). They also proposed innovative
410 methodologies based on HTS to fill the gaps of this database (Rimet et al., 2018a).

411 Aquatic macrophytes are recognized as a valid taxonomic group for assessing water quality
412 according to the WFD. They reflect the morphological conditions of the water bodies
413 (diversity and dynamics of the substratum, degree of rigid management of the banks) and
414 are particularly interesting to assess nutrient pressure. Moreover, they react to anthropogenic
415 interventions in the hydrological regime (potamalisation and water retention). Being plant
416 organisms, macrophytes also present properties, such as longevity and immobility, that make
417 them bad bioindicators in the short-term: they are able to integrate disturbed conditions over
418 a considerably long period of time; it is impossible to accurately locate the source of
419 pressures and the area of impact (Pall and Mayerhofer, 2015). According to the traditional
420 definition, macrophytes are aquatic plants whose vegetative structure develops either in the
421 water on a permanent basis or at least for a few months, or on the surface of water (Cook et
422 al., 1974). These include species of the Charophyta (charales), the Bryophyta (mosses), the
423 Pteridophyta (ferns) and the Spermatophyta (seed plants). In the present study we decided
424 to focus our analyses on vascular plants only, which therefore regroups species from the
425 divisions Pteridophyta and Spermatophyta. Concerning the choice of markers, DNA
426 barcoding in plants is not as straightforward as in animals. The Consortium for the Barcode
427 of Life (CBOL) Plant Working Group ended up by recommending the combination of two
428 plastid loci for the standard plant barcode — rbcL+ matK (Hollingsworth et al., 2009).

429 Several groups of macroinvertebrates are frequently used to report EQS in the WFD.
430 Species-level information on crustaceans, molluscs and the insect orders Ephemeroptera,
431 Plecoptera and Trichoptera (EPT) are widely used. However higher taxa, e.g. genus- or
432 family-level, are also used as BQEs and while some countries only use family-level
433 identifications others use a mixed taxon approach, e.g. the River Invertebrate Classification
434 Tool ([RICT](#)) (Davy-Bowker et al., 2008), used in the UK. There is a great variation between
435 countries in which taxa are used to report to the WFD. For instance, freshwater assessments
436 in the Netherlands utilize 224 species of the dipteran family Chironomidae when reporting
437 water quality status, while Norway does not include species level information on any Diptera.

438 This national-level taxonomic variation in part reflects the natural difference in species
439 occurrences, but is necessary to consider when analysing gaps in the barcode libraries.

440 Freshwater fish are among the most commonly used organisms for assessing EQS
441 according to the WFD, and their community composition and structure is the base for a high
442 number of different metrics in Europe (Birk et al., 2012a). Sampling is conducted using a
443 variety of methods, including electro-fishing or netting and should deliver data on abundance,
444 species composition and age structure of fish present in a water body. However, large
445 differences between countries exist in the percentage of occurring species considered for an
446 assessment, and whether non-native species influence the overall score or not. In Ireland for
447 example, all freshwater fishes are considered for WFD monitoring, while in Austria or
448 Germany only about 60% of the complete fauna is routinely used. While according to
449 practitioners, additional species encountered during sampling are often listed as an
450 amendment to the official sampling protocols and reports, but they often have no impact on
451 the BQE score because the species are not considered in the reference condition. Individual
452 barcoding of sampled freshwater fish is of little use in biomonitoring of natural habitats.
453 However, assessing and monitoring of freshwater fish diversity using eDNA from water
454 followed by metabarcoding can be both more effective and more accurate than traditional
455 specimen sampling (Hänfling et al., 2016; Valentini et al., 2015). Studies have indicated that
456 the standard DNA barcode marker (COI) might not be optimal for this use (Kat Bruce & Emre
457 Keskin pers. obs.), likely since non-target organisms are co-amplified with the available
458 primers and mask the DNA signal from fish. Thus, a much higher sequencing depth is
459 needed to reliably detect all fish species occurring in the studied waterbody, and constitutes
460 suboptimal usage of available resources. Studies have shown that a hypervariable region of
461 the rRNA 12S marker is a suitable target to amplify fish eDNA (Civade et al., 2016; Miya et
462 al., 2015). As also discussed and successfully tested in DNAqua-Net WG3 (Field & Lab
463 Protocols) this marker has a high potential to become the gold standard for regular eDNA-
464 based fish monitoring in the future. We therefore also evaluate the completeness of the
465 reference library for European freshwater fish species for 12S sequence data.

466 Aim of this study

467 The purpose of this paper is to identify gaps in DNA barcode reference libraries that are
468 relevant for European countries when reporting water quality status to the EU in the context
469 of the WFD and MSFD. The gaps for freshwater taxa are reported by country and taxonomic
470 group, and compared across Europe, while gaps for marine organisms are evaluated by
471 taxonomic group. We also discuss the necessity of both quality assurance and quality control
472 (QA/QC) when building and curating a barcode reference library, and provide
473 recommendations for filling the gaps in the barcode library of European aquatic taxa.

474 2. Material and methods

475 2.1 Checklists and datasets

476 Checklists of taxa used for freshwater EQS assessments according to the WFD were
477 obtained from 30 nations (Supplement 1) through national contact points that were in direct
478 contact with their countries' environment agencies, water authorities, or water research
479 institutes (see acknowledgements). National lists were sorted by taxon and assigned
480 taxonomic coordinators among the authors who concatenated lists and unified the taxonomy
481 (e.g. removing synonyms, checking validity of names, etc.) while keeping the country
482 information for each taxon.

483 For marine species we used two generally accepted checklists to perform the gap-analysis of
484 species relevant to the MSFD and WFD: AMBI - an index designed to establish ecological
485 quality of European coasts, and ERMS (Costello, 2000). With the European focus of this
486 analysis we delimited the AMBI list to a geographical selection by compiling only the species
487 with European occurrence that include the following regions: Barents Sea, Norwegian Shelf,
488 British Isles, Baltic Sea, North Sea, Celtic-Biscay Shelf, Iberian Coast, Mediterranean Sea,
489 and Black Sea. The geographic distribution of each species on the original AMBI list was
490 assessed through the World Register of Marine Species ([WoRMS](#)), as well as by the Ocean
491 Biogeographic Information System ([OBIS](#)). The ERMS checklist on BOLD created by Dirk
492 Steinke, titled 'Marine Animals Europe' (BOLD checklist code: CL-MARAE; last updated on
493 20th March 2017), was used in this analysis. It contains records of 27,634 marine animals. A
494 selection consisting of 21,828 species was used for further analysis, including taxonomic
495 entities: Annelida, Arthropoda: Decapoda and Peracarida, Brachiopoda, Chordata:
496 Euchordata - Pisces, Cnidaria, Echinodermata, Mollusca: Bivalvia and Gastropoda,
497 Nemertea, Priapulida, and Sipuncula. We focused on benthic macroinvertebrates and fish
498 and did not look specifically into meiofauna or pelagic animals (except fish), although many
499 of the included species may have life-stages occurring in both environments.

500 Vascular plant checklists were checked for synonyms using three public databases: The
501 International Plant Names Index (<http://www.ipni.org>), The Plant List
502 (<http://www.theplantlist.org>) and Tropicos® (<http://www.tropicos.org>).

503 For freshwater fish, we treated Europe as geographic entity, not by its political borders, but
504 follow its definition as a "continent" with Turkey, Russia and Kazakhstan being only partly
505 included and only with faunistic elements occurring in watersheds that lie within Europe (see
506 also (Kottelat and Freyhof, 2007)). All lists were made available to taxonomic coordinators of
507 selected taxonomic groups (specialists among the authors) to assure conformity of taxonomy

508 and correct spelling. In this process, the taxonomic validation tool available from the Global
509 Biodiversity Information Facility ([GBIF](#)), and WoRMS were used. For fish, the applied
510 taxonomy mostly follows the international Catalog of Fishes (Fricke et al., 2018), which is
511 also the backbone for the BOLD taxonomy.

512 Finalized species-level checklists were concatenated and uploaded to BOLD, and initial gap-
513 analysis reports were retrieved. The reports were examined by taxonomic specialists to see if
514 any reported gaps were due to taxonomic incongruence between the checklist and the BOLD
515 taxonomic backbone. These were corrected in the uploaded checklists before final analysis
516 (Supplement 2). Separate spreadsheets retaining the country information for each taxonomic
517 group were kept for downstream analyses.

518 2.2 Gap-report analyses

519 Two sources of data were retained from BOLD for the majority of the taxonomic groups.
520 Firstly, the checklist progress report option implemented in BOLD was used. Secondly, the
521 checklists were compared to all publicly available sequence information in BOLD by using
522 datasets for each taxonomic group. Progress reports and datasets were generated on the 6th
523 July 2018 for all groups except freshwater fish (1st February 2018), freshwater Annelida (17th
524 September 2018) and Odonata (29th November 2018). The dataset for Diptera used for the
525 reverse taxonomy analysis was generated on the 18th December 2018. The analyses were
526 based on one or two barcode markers, depending on the taxonomic group (see Table 2).

527 Based on the BOLD gap reports, gap-analyses and summarizing statistics were calculated
528 for all taxonomic groups using an analytical pipeline of custom made python scripts
529 [deposited in GitHub <https://github.com/dnaquanet/gap-analysis.git>]. This pipeline was largely
530 the same for all groups, except where specified under specific taxon treatment sections.

531 The data from taxonomic checklists with country information (i.e. nations in which the
532 respective species are monitored) were combined with the information from BOLD. Species-
533 based summaries were generated containing the number of countries in which a species is
534 monitored by extracting the information from the taxonomic checklists. In addition, the total
535 number of reference sequences stored in BOLD (i.e. sequences \geq 500 bp), hereafter referred
536 to as DNA barcodes, were taken from the progress report of each checklist. Additional BOLD
537 quality criteria for barcodes, such as the availability of a trace sequence, were not
538 considered. Using information from the publicly available data from the dataset output, it was
539 possible to calculate the number of barcodes publicly stored in BOLD (BOLD public) or
540 mined from GenBank (GenBank) as well as the number of privately stored barcodes in BOLD
541 (BOLD private). Sequences flagged due to potential contamination, misidentification, or

542 presence of stop-codons, were excluded from the analyses. For some species, DNA
543 barcodes were deposited under the valid species name as well as under synonyms. In these
544 cases, synonyms were part of the BOLD checklists and the barcode hits were merged to the
545 valid species names.

546 In a further step, the proportion of species represented by a minimum number of DNA
547 barcodes (threshold of 1 or 5) was calculated for each checklist. Additionally, country-based
548 summaries were generated, providing an overview of the number of monitored species
549 together with the percentage of barcode coverage for each taxonomic group in the reference
550 libraries (threshold of 1 or 5). For both summary overviews, the available barcode information
551 was sorted into three classes: BOLD public, BOLD total (including BOLD public and BOLD
552 private) and total (including BOLD public, BOLD private, and GenBank). The data were
553 visualized using the python-module matplotlib (Hunter, 2007) and cartopy
554 (scitools.org.uk/cartopy) together with geographical information from naturalearthdata.com.

555 In contrast to all other gap-analyses, no geographical data were included for the marine taxa.
556 Hence, the country-based analysis steps of the pipeline were omitted. Due to the large size
557 of the ERMS checklists, no datasets could be produced in BOLD. Thus, only the results of
558 the progress report were analysed for the availability of reference sequences. In the analysis
559 of species used to calculate the AMBI, datasets could be produced in BOLD, and our
560 analyses could distinguish between BOLD public, BOLD private, and GenBank sequence
561 data.

562 To identify if species belonging to different ecological groups of the AMBI are equally well
563 represented by reference sequences, a further gap-analysis was performed with species
564 classified based on their ecological value.

565 For diatoms, the Diat.barcode library version 7 (Rimet et al., 2016) rather than BOLD was
566 used, as this database is curated by diatom experts to ensure high-quality barcodes. Two
567 genetic markers (rbcL and 18S) are used for barcoding diatoms (e.g. (Vasselon et al., 2018;
568 Zimmermann et al., 2014), and the taxonomic checklists were compared to all available rbcL
569 and 18S data in the database. Both, valid species names and synonyms were considered;
570 subspecies were also accepted as valid. An overall gap-analysis and country-based
571 summaries were generated. However, only a threshold of 1 was used. As all barcodes in
572 Diat.barcode are publicly available at https://www6.inra.fr/carrtel-collection_eng/Barcode-database, the differentiation between public and private data did not apply. Due to the high
574 species diversity in diatoms, estimated at 100,000 (Mann and Vanormelingen, 2013), many
575 low-frequency species could potentially negatively impact the barcode coverage, while the
576 high-frequency (abundant) species could be sufficient for monitoring (Lavoie et al., 2009).

577 Hence, we re-analysed the barcode coverage for two checklists (France freshwater
578 phytobenthos and Croatia marine diatoms) using only high-frequency species.

579 Two standard barcode markers (rbcL or matK) are accepted for vascular plants in BOLD.
580 However, the checklist progress report does not include information on which of the two
581 barcode markers were covered for each taxon. Hence, the first part of the analyses
582 described above was conducted for vascular plants regardless of which of the two markers
583 was present (rbcL OR matK). In contrast, the BOLD dataset includes information on which
584 marker is sequenced for a certain record. Hence, for the public data (BOLD public and
585 GenBank) gap-analyses were performed for each marker as well as for the combination of
586 both markers (rbcL AND matK).

587 For gap-analysis of freshwater fish we also included the 12S marker. Since there are no 12S
588 sequence data available in BOLD (as of February 1st 2018) for European freshwater fishes,
589 we manually compared our target species list with the available mitochondrial genomes from
590 MitoFish (<http://mitofish.aori.u-tokyo.ac.jp>), and NCBI's RefSeq and Nucleotide databases.
591 All available sequence data for Actinopterygii (whole mitochondrial genomes and full or
592 partial 12S sequences) were imported into the software Geneious version 7.1.9 (Biomatters
593 Ltd, New Zealand) and after aligning with the MAFFT-plugin (Katoh and Standley, 2013)
594 trimmed to the hypervariable region of the 12S rRNA gene using the published primer pair
595 MiFish-U/E (Miya et al., 2015) as correctly given in Ushio et al. (Ushio et al., 2018). In the
596 final alignment only species present with sequence information for this locus (ca. 175 bp)
597 were retained and used for the gap list evaluation. Due to the completeness of the barcoding
598 databases for species used in country-based monitoring lists, in general, no geographical
599 information was used for the gap-analysis. However, a map was generated for species of the
600 European-wide fish list where barcodes are still missing.

601 Finally, we refrained from providing any particular DNA barcode gap-analysis for
602 groundwater ecosystems and their species pools. This is because the biological component
603 is currently not considered for subterranean freshwater monitoring and reporting under the
604 umbrella of the WFD, which relies on the chemical status and water quantity in aquifers
605 instead.

606 2.3 Reverse taxonomy

607 As a case study, we analysed the proportion of public barcodes originating from reverse
608 taxonomy for freshwater macroinvertebrates, i.e. specimen identification via its DNA barcode
609 and not by morphology. In the datasets obtained from BOLD, the entry "Identification
610 Method" was screened for the presence of several keywords e.g. "BOLD ID Engine", "BIN

611 Taxonomy Match", "Tree based identification" or "DNA Barcoding". A full list is deposited in
612 Supplement 3. For each species, the number of public barcodes originating from reverse
613 taxonomy was compared to the total number of available public barcodes in BOLD. Four
614 cases were considered, in which reverse taxonomy can have a strong influence: i) all public
615 data originates from reverse taxonomy, ii) more than half of the public data originates from
616 reverse taxonomy, iii) only when including barcodes based on reverse taxonomy, at least five
617 public barcodes are present and iv) when less than five public barcodes are present, at least
618 one originates from reverse taxonomy.

619 3. Results

620 Our results revealed considerable variation in barcode coverage for selected major groups in
621 the queried databases (Table 1). Freshwater vascular plants and freshwater fish had the
622 largest coverage, though still less than 70% of the species had five or more barcodes
623 available. The lowest barcode coverage is found in the marine invertebrates of the ERMS list
624 9.9% (five or more barcodes) to 22.1% (one or more barcodes) and diatoms (14.6%), while
625 more than 60% of the 4502 freshwater invertebrate species used in ecological quality
626 assessments of freshwater ecosystems had one or more barcodes (Table 1).

627

628 Table 1. Overall barcode coverage for selected major groups.

Taxonomic group	Barcode marker	Species in checklist	Barcode coverage [%]		Database source
			≥ 1 Barcode	≥ 5 Barcodes	
Marine invertebrates - ERMS	COI	16,962	22.1	9.9	BOLD
Marine invertebrates - AMBI	COI	3,012	47.6	25.0	BOLD
Marine fish ^a	COI	1,489	82.1	64.3	BOLD
Diatoms (marine and freshwater)	rbcL / 18S	3,716	14.6	N/A	Diat.barcode v7
Freshwater vascular plants	rbcL / matK	683	83.0	69.4	BOLD

Freshwater	COI	4,502	64.5	41.8	BOLD
invertebrates					
Freshwater fish	COI	627	87.9	66.2	BOLD/NCBI

629 ^a Actinopterygii, Elasmobranchii and Holocephali

630

631 3.3 Marine macroinvertebrates & fish


632 3.3.1 Gap-analysis for the European AMBI-list

633 A total of 3,012 marine species were compiled in the AMBI checklist for Europe. Forty-eight
634 percent of them have at least one representative DNA barcode sequence in either BOLD or
635 GenBank, but as much as 23% of those species only have private records (Fig. 1,
636 Supplement 2), and 22% of those with barcodes are single specimen records.

637

638

639



640

641 Figure 1. Cumulative barcode coverage for marine invertebrates in the AMBI list. Barcode
642 coverage of at least 1 reference sequence (A) or a minimum of five reference sequences (B).
643 If barcodes of a species were not recorded in the BOLD public library, the BOLD private
644 library was queried, and subsequently GenBank. Numbers on bars refer to total number of
645 species in checklist. Thick bars represent phyla, thin bars represent taxa of lower taxonomic
646 rank. Taxonomic groups with less than ten species are not indicated.


647 Among the 10 largest taxonomic groups included in this particular analysis, the Chordata
648 (excluding Vertebrata) displayed the lowest proportion of species with DNA barcodes (38%),
649 though only 26 species (within Ascidiacea) were listed for this taxon. In comparison, the best
650 represented taxon was the Nemertea, which has DNA barcodes for 81% of the 27 species
651 considered, while the second most complete group has 67% (Echinodermata). Most of the
652 remaining taxa have completion levels between 40 and 50%, including the three most

653 species-rich taxa (Annelida, Mollusca and Arthropoda), that comprise 85% of the species in
654 the European AMBI checklist (Fig. 1).

655 A narrower analysis of Mollusca shows that Bivalvia and Gastropoda have only moderate
656 levels of completion (50 and 47%, respectively), whereas within malacostracan crustaceans,
657 Decapoda (Arthropoda) is far more complete (84%) than Peracarida (45%). However, the
658 number of species considered is highly disparate for these two groups (25 Decapoda vs. 649
659 Peracarida) (Fig. 1). The proportion of singletons (i.e. only one barcode sequence available)
660 per taxonomic group ranges from 10% to 25%, although for some taxa the observed
661 proportion of singletons was considerably higher (e.g. 50% in Brachiopoda and 38% in
662 Sipuncula).

663 Most of the species from the AMBI checklist have public DNA barcodes available either from
664 BOLD or GenBank, with only 11% represented exclusively by private records. Two groups
665 have slightly higher values, Echinodermata (15%) and Arthropoda (12%). The levels of
666 completion by AMBI's Ecological groups (I to IV) are similar, ranging from 43% in group IV to
667 56% in group III (Supp. Fig. 1). However, 215 species were not assigned to ecological
668 groups, and among these the completion is low (ca. 38%). Species barcodes found
669 exclusively in BOLD private range from 10% (IV) to 13% (V) in each of AMBI's ecological
670 groups.

671 3.1.2 Gap-analysis for the ERMS checklist



672

673 Figure 2. Barcode coverage for marine invertebrates of the ERMS checklist. Barcode
674 coverage of at least 1 reference sequence (A) or five reference sequences (B). Thick bars
675 represent phyla, thin bars represent taxa of lower taxonomic rank. Numbers on bars refer to
676 total number of species in checklist. Taxonomic groups with less than ten species are not
677 indicated.


678

679 The selection from the ERMS list on BOLD contains 16,962 species. Twenty-two percent of
680 these species have at least one DNA barcode in BOLD (Fig. 2). Of these species, 26% have

681 singletons and nearly 10% have five or more DNA barcodes. These figures include DNA
682 barcodes from GenBank that are present in BOLD. The highest coverage is found in
683 Decapoda (50%), followed by Sipuncula (42%), a phylum with 45 species only found in the
684 ERMS list (Fig. 2). At the other end, the lowest coverage (11%) is observed in Brachiopoda
685 (37 species). Nemertea also have a low coverage, 15% for the 380 listed species. The
686 coverage of most other taxonomic groups ranges from 20 to 30%.

687 Within phyla, there are clear differences in the proportion of DNA barcodes between
688 taxonomic subgroups. Arthropods have a coverage of 20% as a whole, but the Decapoda
689 reach 50%, while the Peracarida reach only 23%. Within Mollusca, with an overall coverage
690 of 20%, Bivalvia reach 24% and Gastropoda 18%. The proportion of singletons roughly
691 follows the inverse pattern as the proportion of total DNA barcodes: the lowest proportion of
692 8% is found in marine fish, while the highest proportion of 57% is found in Brachiopoda.

693 A detailed analysis of cnidarians in the ERMS checklist reveals that while 353 of the 1,201
694 species (29.4%) are listed with sequence information in BOLD, only 97 species (8.1%) have
695 sequences that meet the formal barcode requirements. We observed that many of the
696 sequences were mined from GenBank, containing limited information and are a potential
697 source of errors. A similar situation was observed for ascidians where 84 out of 402 species in
698 the ERMS checklist (20.9%) have sequence information while, only 5.7% of the species had
699 references to vouchers and sufficient metadata to be barcode compliant.



701 Figure 3. Barcode coverage for marine fish of the ERMS checklist. Barcode coverage by at
702 least 1 reference sequence (A) or five reference sequences (B). Thick bars represent all fish,
703 thin bars represent lower taxonomic rank. Numbers on bars refer to total number of species
704 in checklist.


706
707 The marine fish checklist obtained from ERMS includes 1,489 species partitioned among the
708 three most prominent classes examined as follows: Actinopterygii (1,339), Elasmobranchii
709 (143) and Holocephali (7). Overall, 82% of the species are barcoded (64% \geq 5 barcodes),
710 ranging from 100% (71% \geq 5 barcodes) for the Holocephali to 81% (63% \geq 5 barcodes) for
711 the Actinopterygii, with the Elasmobranchii coverage is in between (92% \geq 1 barcodes, 80%
712 \geq 5 barcodes) (Fig. 3).

713 3.2 Diatoms

714 Taxonomic checklists for diatoms were obtained from 16 countries and contained a total of
715 3,716 species ranging from 6 (Albania) to 2,236 species (France). This list covers very
716 different habitats, freshwater phytobenthos, freshwater phytoplankton and marine
717 phytoplankton. Some national checklists did not mention which habitat was covered.

718 The general coverage of diatoms was very low, with 15% of all species having at least one
719 sequence of rbcL or 18S (Fig. 4). The coverage of rbcL (13%) is slightly better than the
720 coverage of 18S (11%). However, in most cases both markers are present if any sequence is
721 available (9%). Per country, the coverage ranged from 10% (France) to 37% (Italy), when
722 both markers are present and 15% (France) to 55% (Italy), when at least one of the markers
723 is present (Suppl. Fig. 1).

724



725


726 Figure 4. Cumulative barcode coverage of diatoms.

727

728 A gap-analysis of diatoms ranked by the number of countries that monitor those species,
729 revealed that the most frequently monitored species have a moderate to high representation
730 for both markers (Fig. 5A). For the 16 species used in 14 countries, 81% have rbcL and 18S
731 data and additionally 13% have rbcL data only. For species monitored by few countries, the
732 barcode coverage is comparatively poor (below 20% for species monitored in ≤ 7 countries).

733

734



735

736 Figure 5. Cumulative barcode coverage of diatoms by the number of countries monitoring
737 them. (A) All diatom species, (B) freshwater phytobenthos species. Numbers on bars refer to
738 the number of species per country category.

739


740 For freshwater phytobenthos, the habitat where diatoms are used most often as ecological
741 indicators, the most frequently monitored species have a moderate to high representation of
742 both markers (Fig. 5B). Similar to all diatom datasets, most of the species monitored in
743 eleven countries are represented by both markers (70%), with additional species barcodes
744 for rbcL (20%). For species monitored by fewer countries, the coverage is considerably
745 smaller (below 20%, for species in ≤ 4 countries).

746 For the most common species of freshwater phytobenthos monitored in France, 553 of the
747 2,236 species were scored as abundant. In this subset, the barcode coverage was 33%,
748 considerably higher than the 15% of all species. The proportion of species with both rbcL and
749 18S sequenced was 20% compared to 10% for all species (Fig. 4). A similar picture was
750 evident for the marine diatoms from Croatia. Of the 100 most frequently observed marine
751 phytoplankton species (including Diatoms, Dinoflagellates, Silicoflagellates and
752 Coccolithophorids), 32 were diatoms. Of these 32 species, 50% had at least one barcode
753 available compared to 36% in the total dataset of 729 species. The proportion of species with
754 both barcodes was 34%, compared to 25% for all species (Fig. 4).

755 3.3 Vascular plants

756 General taxonomic checklists for freshwater macrophytes were obtained for 16 nations. The
757 compiled list of 1,242 species names was filtered for vascular plants, resulting in 683
758 species. In general, vascular plants are well covered by one or the other standard barcode
759 marker, with more than 83% of the species having at least one sequence, and 69% having at
760 least 5 sequences (Tab. 1).

761




762

763 Figure 6. Barcode coverage for freshwater vascular plants. (A) ≥ 1 DNA barcode available in
764 BOLD, (B) ≥ 1 DNA barcode available in BOLD or GenBank, (C) ≥ 5 DNA barcodes available
765 in BOLD or (D) ≥ 5 DNA barcodes available in BOLD or GenBank.

766

767 Compared with public records, however, these results seem slightly overemphasized as 22%
768 (153) of the species have no *rbcL* nor *matK* sequences publicly available on BOLD (or mined
769 from GenBank, Fig. 6A). Moreover, only 46% (316) of the species have barcodes for both
770 loci publicly deposited in BOLD. The remaining 214 species have incomplete data: *i*) *rbcL*
771 publicly deposited in BOLD, but *matK* sequences absent (53), or mined from GenBank (80);
772 *ii*) sequences for both loci coming from GenBank (38); *iii*) sequences for only one locus
773 issued from GenBank (*rbcL* - 28; *matK* - 15).

774 In sum, *rbcL* is the best represented DNA barcode marker for vascular plants with 75% of
775 the species having publicly deposited sequences, and 66% of the species having BOLD
776 public data (Fig. 6). Sixty-six percent of the species have publicly deposited barcodes for
777 *matK*, with only 46% of the species having sequences deposited in BOLD public.



778

779 Figure 7. Barcode coverage maps for freshwater vascular plants (lim 1 = minimum one
780 record). (A) Number of monitored species per country, (B) barcode coverage per country with
781 all available data (total), (C) rbcL-specific coverage per country publicly available in BOLD,
782 (D= matK-specific coverage per country publicly available in BOLD, (E) cumulative barcode
783 coverage of vascular plants available in BOLD by number of countries monitoring them, (F)
784 cumulative barcode coverage of vascular plants available in BOLD or GenBank by number of
785 countries monitoring them.


786

787 The number of monitored species varied strongly, ranging from six (Poland) to 394 (Hungary,
788 Fig. 7A). The average barcode coverage (BOLD and GenBank data) was relatively evenly
789 distributed with a minimum of 76% (Lithuania), reaching 100% in three countries (Austria,

790 Poland and Switzerland, Fig. 7B). A higher and more homogeneous coverage was found for
791 rbcL (67 - 90%; Fig.7C) than matK (0 - 74%; Fig. 7D), both for BOLD public and GenBank
792 data (rbcL: 71% - 100%; matK: 50% - 87%; Supp. Fig. 2). Two species were monitored in
793 twelve countries (*Alisma lanceolatum* and *A. plantago-aquatica*) and approximately one fifth
794 of the species in more than 4 countries (Fig. 7E, F). The barcode coverage of these species
795 was 100% when public and private data were taken into account. It decreased slightly for
796 species monitored in four or fewer countries. Nevertheless, more than 40% of the 330
797 species monitored in one country only had rbcL and matK data deposited publicly in BOLD
798 and 73 % had associated sequences when private BOLD and GenBank data were included.


799 3.4 Freshwater macroinvertebrates

800 The analysed national monitoring checklists comprise 4,504 species of freshwater
801 macroinvertebrates, including insects (ca. 80% of the listed species), annelids (ca. 6%),
802 arachnids (ca. 5%), crustaceans (ca. 4%), molluscs (ca. 4%), flatworms (ca. 1%) and
803 nematodes (< 0.05%). When considering all species with at least one barcode in BOLD,
804 64.5% of the species are covered (Fig. 8). Most barcodes are publicly available. For the
805 more strict criterion of ≥ 5 barcodes per species, only 41.9% of the species are covered.
806 Among all taxonomic groups considered in the analysis, the three insect orders Odonata,
807 Trichoptera and Hemiptera along with crustaceans are best covered with $\geq 80\%$ of species
808 barcoded from each taxonomic group. The groups with the least coverage are flatworms
809 (less than 5%), followed by annelids, molluscs and certain insect orders, such as Diptera and
810 Ephemeroptera, in which less than 60% of listed species are represented by at least one
811 barcode (Fig. 8). Only in the case of Hemiptera, more than 80% of the species are
812 represented by at least five barcodes while, except for Odonata, Trichoptera, Coleoptera and
813 Crustacea, less than 50% of the species are covered in the other macroinvertebrate groups.
814 For some groups, such as molluscs, annelids and crustaceans, a substantial share of the
815 available reference sequences are not deposited in BOLD, but present in GenBank (Fig. 8).
816 The most monitoring-relevant insect taxon with lowest coverage on BOLD is Diptera (ca.
817 60% of the 2,108 species in the list). Hemiptera, with 76 species listed and ca. 92% already
818 barcoded will probably be the first group to have full coverage in the near future.



819

820 Figure 8. Cumulative barcode coverage for freshwater invertebrates. Barcode coverage by at
821 least one reference sequence (A) or five reference sequences (B). If barcodes of a species
822 were not recorded in the BOLD public library, the BOLD private library was queried, and
823 subsequently GenBank. Thick bars represent higher taxonomic ranks, thin bars represent
824 insect orders. Numbers on bars refer to total number of species in checklist. Taxonomic
825 groups with less than ten species are not indicated.




826

827

828 Figure 9. Barcode coverage per species. The upper bars show the barcode coverage (up to
829 a maximum of 10 barcodes). The lower bars show the number of countries in which a
830 species is monitored. (A) Annelida, (B) Arachnida, (C) Crustacea, (D) Insecta, (E) Mollusca.

831 3.4.1 Insects
832 Insects are used for monitoring ecological status in 29 out of the 30 surveyed countries. All
833 national monitoring checklists combined comprised 3,619 insect species (Supplement 2, Fig.
834 9D). However, taxonomic resolution used between countries differed substantially. Seven
835 countries exclusively assess taxonomic groups above species level, two countries only
836 above genus level, and five countries only above family level (Supplement 1). Assessed taxa
837 per country range from 10 (Albania) to 2,903 (Czech Republic, Fig. 10). In total, eleven insect
838 orders are monitored, ranging from orders with only one relevant species (Hymenoptera) to
839 orders with 2,108 species (Diptera, Fig. 8). The top ten species monitored in most countries
840 all belong to Ephemeroptera with *Ephemera danica* and *Serratella ignita* being the most
841 frequently listed species (20 countries each).
842 On average, 65.7% of all monitored insect species are barcoded, ranging from orders with
843 only 52.7% and 58.8% barcoded species (Ephemeroptera and Diptera) to highly covered
844 orders (Trichoptera, 87.0%; Odonata, 90.5%; Hemiptera 92.1%; Fig. 8). A high proportion of
845 barcodes for these species is deposited in BOLD (95.3%; 91,066 barcodes) of which 70.9%
846 have publicly available metadata. However, for 513 barcoded species (14.2%) there is no
847 BOLD public data. For the most frequently monitored species, *Ephemera danica*, there are
848 only 4 public (and 11 private) COI barcodes in BOLD. In contrast to the top monitored
849 species, 9 of the 10 species with the most barcodes (BOLD and Genbank combined) belong
850 to Diptera with the two Chironomidae species *Paraphaenocladius impensus* (5,981
851 barcodes) and *Paratanytarsus laccophilus* (4,058) being the most often barcoded species. Of
852 the 1,240 missing insect species that are monitored in at least one country, 917 are
853 monitored in a single country (Czech Republic), and 674 of those species are exclusively
854 monitored in that country. The coverage of barcoded species per country is on average
855 87.6%, ranging from 51% (Georgia, only mayflies) and 68% (Czech Republic) to 98%
856 (Estonia) and 100% (Ireland, just three taxa monitored at the species level).
857



858

859 Figure 10. Barcode coverage maps of Insecta. (A) Number of monitored species per country.
860 (B) - (F) Barcode coverage per country for different datasets (BOLD public, BOLD total and
861 total) and thresholds (lim 1 = minimum one record; lim 5 = minimum five records).

862


863 3.4.2 Arachnids

864 A large proportion of the arachnid species records in BOLD is private (Fig. 8). The coverage
865 of the 211 species reported from all countries in total is moderate with 65% of the species
866 represented with at least one barcode. It is remarkable that 201 of the 211 arachnid species

867 are only monitored by one country, the Netherlands (Fig. 9B). Of these, 200 are solely
868 monitored in this country. The spider *Argyroneta aquatica*, which is monitored by the most
869 countries (7), has only private reference barcodes in BOLD, and five sequences in GenBank.

870 3.4.3 Crustaceans


871 A total of 193 crustacean species are included in the nationwide checklists; 22 of the 30
872 surveyed countries monitor one or more crustacean species (Fig. 11). They represent four
873 classes: Branchiopoda (62 spp.), Hexanauplia (25 spp.), Ichthyostraca (3 spp.) and
874 Malacostraca (110 spp.). Among them, the most frequently monitored species are the
875 malacostracans: common waterlouse, *Asellus aquaticus*, monitored in 19 countries, the
876 noble crayfish, *Astacus astacus* - in 16 countries, and the amphipod *Gammarus roeselii* - in
877 12 countries. Each of these species is covered with numerous private only reference
878 barcodes in BOLD or publicly available sequences in GenBank (Fig. 9C). Thirty six species
879 of crustaceans have no barcode coverage at all, neither in BOLD nor in GenBank, while 26
880 are covered only in GenBank. Among those covered in BOLD, 67 species are represented
881 by private reference barcodes only. Most of the species (121 spp.) are monitored only in one
882 country. For example, 53 species, predominantly branchiopods and hexanauplians, are
883 monitored in Sweden only. Eleven of these species have no barcode coverage, neither in
884 BOLD nor in GenBank, while 22 species are represented only by private barcodes.
885 In general, the barcode coverage (including GenBank data) per country is good and relatively
886 evenly distributed, from 70% to 100% of species barcoded in each country (Fig. 11D). These
887 values drop down immensely when only the public BOLD data are taken into account (Fig.
888 11B). In the countries such as Italy and Ireland not even 10% species is covered, while only
889 in Germany, UK, the Netherlands and Norway the coverage approaches 50% of the species
890 monitored in each of these countries.



891
892
893
894
895
896
897

Figure 11. Barcode coverage maps of Crustacea. (A) Number of monitored species per country. (B) - (F) Barcode coverage per country for different datasets (BOLD public, BOLD total and total) and thresholds (lim 1 = minimum one record; lim 5 = minimum five records).


898 3.4.4 Annelids
899 In total, 257 species of annelids are used in freshwater biomonitoring in the 21 countries that
900 supplied lists (Fig. 12). They represent two classes, Clitellata with the subclasses of
901 Oligochaeta, Hirudinea (leeches) and Branchiobdellida and Polychaeta with the subclass
902 Sedentaria. Among them, three species of leeches, *Erpobdella octoculata*, *Glossiphonia*
903 *complanata* and *Helobdella stagnalis* are monitored in 20 countries (Fig. 9A). Further 21
904 species of both leeches and oligochaetes are monitored in 11 to 19 countries. The most
905 commonly monitored polychaete is the freshwater alien *Hypania invalida* included in lists of
906 five countries. The other alien species, *Marenzelleria neglecta*, is generally brackish water
907 and is monitored only in Germany. A couple of other brackish water native species are
908 generally monitored in single countries only. Altogether almost 50% of the listed species are
909 represented by DNA sequences. However, they are generally poorly represented in BOLD,
910 where barcodes for ca. 40% of the species are deposited and only some 20% are publicly
911 available. Most of the species with no barcodes at all are monitored in few countries only
912 (predominantly in Czech Republic and Slovakia) with some notable exceptions, such as the
913 leech *Alboglossiphonia heteroclitia* present on the lists of 15 countries, and the oligochaete
914 *Haplotaxis gordioides* monitored in 12 countries.
915 Country wise, the barcode coverage (including GenBank data) extends from ca. 50% of
916 species barcoded in Czech Republic and Slovakia to 100% in Norway (Fig. 12D). When only
917 public BOLD records are considered, the barcode coverage per country drops down to 20%-
918 40% (Fig. 12B).



919
920
921
922
923
924
925

Figure 12. Barcode coverage maps Annelida (A) Number of monitored species per country. (B) - (F) Barcode coverage per country for different datasets (BOLD public, BOLD total and total) and thresholds (lim 1 = minimum one record; lim 5 = minimum five records).

926 3.4.5 Molluscs
927 The national checklists of freshwater molluscs contain a total of 161 species, ranging from
928 one (Cyprus) to 77 (Czech Republic) species per country (Fig. 13). *Ancylus fluviatilis*, the
929 most commonly surveyed species, is included in 20 national checklists, while a total of 67
930 species are considered by a single checklist only (22 of them in Georgia) (Fig. 9D). The total
931 barcode coverage of freshwater molluscs (about 60%) was in the range of most freshwater
932 invertebrate groups (Fig. 8). While the proportion of species with public barcodes deposited
933 in BOLD was relatively low (only 15%), the proportion of species with sequences derived
934 only from GenBank was considerably high (24%). A similar pattern was evident when a
935 minimum coverage of five barcodes was used (Fig. 8B). Here, 41% of the species met the
936 criteria when all public and private data were considered, 10% of the species were covered
937 in the BOLD public database, while 21% of the species only had sufficient barcodes if
938 GenBank data were considered together with data from BOLD.



939

940

941 Figure 13. Barcode coverage maps of Mollusca. (A) Number of monitored species per
942 country. (B) - (F) Barcode coverage per country for different datasets (BOLD public, BOLD
943 total and total) and thresholds (lim 1 = minimum one record; lim 5 = minimum five records).

944

945 A high proportion of the missing barcodes was found for species that are used in freshwater
946 monitoring in a single country (41 species, Fig. 9D). Only five of the 35 species surveyed in
947 at least ten countries had no barcodes available. However, a comparatively high number of


948 widely distributed and as such often listed freshwater molluscs (at least on 5 lists) do not yet
949 have barcode data (23%). The barcode coverage per country was relatively evenly
950 distributed, with an average coverage of 23% (min: 0% - Cyprus, max: 38% - Italy) when
951 public barcodes in BOLD were considered, 56% (min: 0% - Cyprus, max: 76% - Finland)
952 when public and private data on BOLD were used and 76% (min: 0% - Cyprus, max: 94% -
953 Finland) for the full BOLD and GenBank datasets (Fig. 13).

954 3.4.6 Platyhelminthes
955 Overall, 61 freshwater flatworm species are used for monitoring in 16 countries. The number
956 of species monitored per country ranged from one (Estonia) to 39 (Czech Republic). While
957 most species are observed in only a few countries, there are nine species on at least ten
958 national checklists, with *Dendrocoelum lacteum* being the most common (14 countries;
959 Supplement 2). The barcode coverage of freshwater Platyhelminthes was very low (4.9%,
960 Fig. 8) as only three species had sequences deposited in examined databases. Of these, two
961 species (*Dugesia cretica* and *Girardia tigrina*) had only one COI-sequence mined from
962 GenBank, while 51 private barcodes were available for *Dugesia gonocephala*.


963 3.4.7 Nematodes
964 Nematodes can be ascribed to both meio- and macroinvertebrate fauna depending on size of
965 the respective freshwater forms. Only ten of the national checklists include (assumed) free-
966 living and semi-parasitic forms, mostly on a coarse taxonomic level. The lists contained one
967 taxonomically wrong classification (*Gordius aquaticus* as Nematoda instead Nematomorpha),
968 one fish parasite, semi-parasitic forms of the family Mermithidae, and one higher order which
969 is taxonomically no longer in use (Secernentea). Only the Romanian list contain two relevant
970 nematode species (*Dorylaimus stagnalis* and *Tobrilus gracilis*). These are common in
971 freshwater, and both are represented with barcodes in BOLD.

972 3.5 Freshwater fish
973 As of 1st February 2018, the target list for European freshwater fishes contained 627 species
974 including 18 extinct and 3 'extinct in the wild' species. After the first BOLD checklist query
975 against all available data, 110 of the 627 species were listed as in need of specimens, i. e.
976 completely lacking DNA barcode references in BOLD (coverage: 82.5%, Fig. 14A). When
977 setting the threshold for minimum number of DNA barcodes available to five, 212 species did
978 not have any or fewer than five barcodes deposited in the database (Fig. 14B). After manually
979 checking the resulting gap list and taking into account real synonyms and different taxonomic
980 concepts such as generic assignments (e.g., *Iberocypris* vs. *Squalius*, *Orsinigobius* vs.
981 *Knipowitschia*, only 60 extant species (plus 16 extinct) were not represented with DNA

982 barcodes (coverage: 90.4% or 87.9% including extinct species). Only three species listed in
983 BOLD had records that did not fulfil the formal requirements for DNA barcode status.
984



985
986 Figure 14. Barcode coverage for freshwater fish. A) A minimum of one DNA barcode, B) \geq 5
987 DNA barcodes or C) one 12S sequence per species. Numbers on bars refer to the number of
988 species in checklist. Eighteen extinct and 3 extinct in the wild species are included.
989



990

991 Figure 15. Missing barcodes for freshwater fish species. (A) Number of all species with
992 missing barcodes per country. (B) Number of extinct species with missing barcodes per
993 country. (C) Number of extant species with missing barcodes per country.

994


995 In general, the DNA barcode (COI) coverage for extant species is very good in most countries
996 (100% coverage in 16 countries) and only a few species are missing reference records from
997 certain regions (Fig. 15A, C). In the Scandinavia and the UK, a number of chars (*Salvelinus*
998 spp.), trouts (*Salmo* spp.) and whitefishes (*Coregonus* spp.) are not yet represented in the
999 databases. While for Austria, Germany and Switzerland, a smaller number of whitefishes (<10)
1000 are still missing in the DNA barcode reference libraries. Concerning extinct and extinct in the

1001 wild species (Fig. 15B), only a few species are missing, the highest number of them (6)
1002 reported from Switzerland.

1003 3.6 Reverse taxonomy

1004 Documented use of reverse taxonomy was observed in all groups of freshwater
1005 macroinvertebrates where public data was available, except for Neuroptera (Fig. 16,
1006 Supplement 3). The proportion of identified sequences originating from reverse taxonomy
1007 compared to all available barcodes ranged from 1% (Crustacea, Ephemeroptera, Hemiptera,
1008 Lepidoptera and Odonata) to 20% (Coleoptera) and 59% (Diptera). Since these values rely on
1009 the cumulative number of BOLD-public, BOLD-private and GenBank data, and since the use
1010 of reverse taxonomy is known only from public sequences in BOLD, the calculated proportions
1011 can be underestimations. For instance, when only public data in BOLD is considered, reverse
1012 taxonomy can be found in up to 61% (Annelida) and 82% (Diptera) of the deposited sequences.
1013 The fraction of species with barcodes originating from reverse taxonomy ranged from 3%
1014 (Arachnida, Coleoptera and Ephemeroptera) to 16% (Diptera) and 20% (Megaloptera).
1015 Although the proportion of species having reverse taxonomy of potentially strong influence was
1016 low for most taxonomic groups, it was comparatively high for Diptera (12%) and Megaloptera
1017 (20%).

1018



1019
1020 Figure 16. Overview of reverse taxonomy in freshwater macroinvertebrates. (A) The proportion
1021 of species with reverse taxonomy barcodes for the different taxonomic groups. (B) The
1022 proportion of barcodes originating from reverse taxonomy for the different taxonomic groups.
1023 Thick bars represent higher taxonomic ranks, thin bars represent insect orders. Numbers on
1024 bars refer to total number of (A) species in checklist or (B) total barcodes. Taxonomic groups
1025 with less than ten species or without public data are not indicated.

1026 4. Discussion

1027 3.1 National checklists

1028 We collected lists of species that are used in the national implementations of the WFD and
1029 MSFD monitoring programs and water quality assessments in each target country. However,
1030 since countries have different strategies on how they report and comply with the regulations,
1031 the lists were very different in terms of taxonomic coverage and level of classification among
1032 nations. WFD monitoring requires the use of stressor-specific Multimetric Indices (MMIs,
1033 (Hering et al., 2006)), intercalibrated and validated at river basin level (Hering et al., 2010).
1034 Each country has developed their own set of MMIs, each consisting of biotic indices which
1035 are best suited to describe water quality status in their region (Birk et al., 2012a; European
1036 Commission, 2018). The taxonomic depth of data required for calculation of these indices is
1037 highly variable between countries. In cases where indices are dependent on species-specific
1038 traits, all species counts and complete species-level identification is required. Thus, the
1039 checklists of species from each country that we received and have used as basis for our gap-
1040 analysis can be grouped into four major types.

1041 The first group contains 'full national lists of species'. Such lists are typically generated from
1042 the Pan-European species lists, or compiled individually from literature. Some countries (e.g.
1043 Czech Republic) use these complete lists as basis for their WFD monitoring, even if many
1044 taxa are not regularly encountered. The second group includes lists from countries that use
1045 the national taxa lists as a basis, but narrow down the selection based on experience or
1046 challenges with identification to species level. In Hungary, for example, only species that
1047 were previously recorded during WFD monitoring are used. Other countries would limit the
1048 identification of selected groups to family- or genus-level, or completely discard semi-aquatic
1049 taxa or taxa that are non-aquatic but closely connected to aquatic environments (e.g.
1050 Carabidae, Chrysomelidae or Curculionidae beetles). These restrictions have been taken
1051 into account in index development. In the third group, it is common to regularly monitor the
1052 frequency/occurrence of certain 'highly indicative' species/taxa, and use only these species
1053 in the calculation of MMIs. Thus, a highly restricted 'operational taxon lists' for WFD
1054 monitoring is compiled. Such a list can be extensive or quite short, dependent on country.
1055 The fourth group includes lists that are exclusively based on family or genus-level
1056 identifications. The differences in the submitted taxa lists influence how the geographical
1057 coverage of DNA barcodes should be interpreted. It might be most obvious for the first group
1058 of lists, where a considerable number of taxa are not regularly encountered but still included
1059 in the barcode gap-analysis through the checklists from Sweden, Czech Republic and
1060 Slovakia (Figs 10-13).

1061 4.2 Marine macroinvertebrates & fish
1062 Based on the ERMS checklist, the gap in the reference library for marine species is relatively
1063 large (> 70%) for all analysed taxa, with the exception of fish (18%). The gap is much smaller
1064 when the AMBI checklist is used (little over 50%), but still comparatively high in global terms,
1065 particularly compared to libraries of freshwater taxa. The lower coverage of the ERMS list is
1066 somehow expected since this list contains six times more taxa than the AMBI list (18,451
1067 versus 3,012, respectively). For the purpose of bioassessment under the WFD and MSFD,
1068 AMBI is a more relevant checklist since it describes the sensitivity of macrozoobenthic
1069 species to both anthropogenic and natural pressures, and it is currently used as a
1070 component of benthic invertebrates' assessment in many EU member states within the four
1071 regional seas (Borja et al., 2007; Borja et al., 2009; European Commission, 2018).
1072 The percentage of barcoded species greatly differed between lists and targeted taxonomic
1073 groups. Marine fish (only included in the ERMS list), were by far the best represented
1074 taxonomic group, with barcodes available for 82% of the nearly 1,500 species in the list.
1075 Partially due to the commercial importance of this group, marine fish have been the target of
1076 comprehensive DNA barcoding campaigns along multiple marine regions in Europe (e.g.
1077 (Costa et al., 2012; Keskin and Atar, 2013; Knebelsberger et al., 2014b; Landi et al., 2014;
1078 Oliveira et al., 2016)). However, a fair proportion of the barcode records for marine fish may
1079 not have originated from specimens collected in European seas (Oliveira et al., 2016), since
1080 many species have considerably large distributions (e.g. (Ward et al., 2008)) including, for
1081 example, those occurring also in the northwest and south Atlantic Ocean.
1082 For marine benthic macroinvertebrates in the AMBI list, the three most species-rich phyla;
1083 Annelida, Mollusca and Arthropoda (ca. 85% of the total species in the list), have moderate
1084 levels of completion (40% to 50%), while less represented groups such as Nemertea,
1085 Sipuncula and Echinodermata have completion levels of at least 65%. Within the ERMS list,
1086 the levels of completion were lower than those of the AMBI list, but followed similar trends of
1087 those reported for the AMBI list, with the exception of the nemerteans. The Annelida,
1088 Mollusca and Arthropoda, that accounted for ca. 77% of the species in the ERMS list, have
1089 fair levels of completion (20% to 30%) and lower than less diverse groups in the list, such as
1090 Echinodermata (35%) and Sipuncula (42%).
1091 Our results suggest that many of the barcode studies focused on Annelida, Mollusca and
1092 Arthropoda, may have targeted particular species or groups at the order or family level (e.g.
1093 Crustacea (Costa et al., 2007; Raupach et al., 2015); Decapoda (Matzen da Silva et al.,
1094 2011a; Matzen da Silva et al., 2011b; Matzen da Silva et al., 2013); Amphipoda (Lobo et al.,
1095 2017); Gastropoda (Barco et al., 2013; Barco et al., 2016; Borges et al., 2016); Polychaeta

1096 (Lobo et al., 2016); Bivalvia (Barco et al., 2016). A closer look into particular taxonomic
1097 groups in our analysis supports this: for the order Decapoda, which comprises only 25
1098 species in the AMBI list and 693 species in the ERMS list, 84% of the species are barcoded
1099 in the former, and ca. 50% in the latter. For a larger group such as the superorder
1100 Peracarida, which comprises 649 species in the AMBI list and 2,643 species in the ERMS
1101 list, the total number of barcoded species is much far from completion (45% and 24%,
1102 respectively).

1103 In addition to the globally modest levels of completion for marine macroinvertebrates, the
1104 gap-analyses based on the AMBI checklist also reveals some insufficiencies of the available
1105 data, namely the presence of a sizeable proportion of private records, which are unavailable
1106 for full access in bioassessment studies employing DNA-based tools. For some groups,
1107 private records on BOLD were even higher than the public, such as for Sipuncula (25%
1108 *versus* 10%) and Annelida (20% *versus* 18%). An ISI Web of Science search, at the time of
1109 writing (30th November 2018), with the search terms “barcoding” AND “marine” AND “the
1110 taxonomic group of interest” also supports the absence of published reference libraries for
1111 Sipuncula, or the low number of studies found for Annelida, compared to other above-
1112 mentioned groups (e.g. fish and Crustacea). Another aspect worth of consideration is the
1113 number of singletons in the reference libraries. Although the percentage of singletons is
1114 generally low, some taxa have a considerable proportion of single representatives per
1115 species. Whereas relatively low levels of barcode coverage for some of these groups clearly
1116 reflect fewer efforts to barcode those taxa, a considerable proportion of the gap must also be
1117 ascribed to failed DNA sequencing, due to either primer mismatch, sample contamination or
1118 PCR inhibitors. This is particularly obvious for the marine Annelida, for which COI
1119 sequencing success rates may be down to 40-50 % on average (Kongsrud et al., 2017).
1120 Barcoding of annelids has also revealed unexpected high levels of genetic diversity,
1121 prompting traditional species taxa to be torn apart (Nygren, 2014; Nygren et al., 2018). A
1122 relatively high proportion of private data may reflect that some species taxonomies are
1123 currently in a certain state of flux. [ref. to this insert:

1124 By increasing the threshold of at least one barcode per species to five barcodes, the level of
1125 completion of both lists (i.e. ERMS and AMBI) fell to about half. For instance, the levels of
1126 completion remained acceptable only for fish and Decapoda, but for most groups these are
1127 greatly distant from what would be recommendable, in particular for Sipuncula, Nemertea,
1128 Cnidaria, Brachiopoda and Annelida. Ideally, reference libraries should have a fair and
1129 balanced representation of specimens across the geographic distribution for each species, to
1130 capture the range of intraspecific variation in the DNA barcodes in the best possible way.

1131 Such representation is also key for efficient quality assurance, quality control and validation
1132 of reference libraries, as discussed below.

1133 Within the AMBI list, almost half of the species fall into the ecological group I, which are the
1134 “sensitive” species, and the remaining half is distributed among the other 5 ecological
1135 groups. However, the completion levels were higher for species from ecological groups III
1136 (56%) and V (52%) and lower for species that do not have any ecological group assigned
1137 (38%). Similar results were encountered when the first attempt of using a genetics based
1138 marine biotic index (gAMBI), with available GenBank sequences for AMBI species, has been
1139 performed (Aylagas et al., 2014). At the time, the authors concluded that the available
1140 genetic data was not sufficient or did not fulfil the requirements for a reliable AMBI
1141 calculation, that needs an even distribution of taxa across the disturbance gradient. On the
1142 other hand, when gAMBI values were calculated by using the most frequent species within
1143 each ecological group, the reliability of AMBI values increased significantly (Aylagas et al.,
1144 2014). Nevertheless, in the current study we have found a much higher completion level (e.g.
1145 48% versus 14%), since numerous new records have been generated in the meantime and
1146 our gap-analyses also included BOLD data.

1147 4.3 Diatoms

1148 Several diatom studies have pointed out the barcode reference library as the Achilles heel of
1149 using metabarcoding of diatoms for environmental monitoring (Kermarrec et al., 2013;
1150 Rivera et al., 2018a; Rivera et al., 2018b; Vasselon et al., 2017). The barcode reference
1151 library must be as comprehensive as possible in order to assign a high proportion of
1152 environmental sequences to known taxa, and it requires regular expert curation in order to
1153 maintain quality. This is why experts from several countries joined efforts to curate a single
1154 reference library, Diat.barcode (formerly called R-Syst::diatom). Our results show that a large
1155 majority of the most common species (registered in the checklists of all countries) are
1156 present in this library, but that many rare species lack representation.

1157 A comprehensive barcode reference library for diatoms is difficult to achieve for two reasons.
1158 Firstly, because more than 100,000 species are estimated to exist globally (Mann and
1159 Vanormelingen, 2013), many of which are undescribed. Registration of barcodes and
1160 metadata of all these species in the reference library will require an overweening effort. Thus,
1161 an effort should be focused on the most common, not yet barcoded species. Secondly,
1162 diatoms need to be isolated and cultured in order to obtain high quality, vouchered, barcode
1163 records. This work is tedious and often unsuccessful because many species are difficult or
1164 impossible to cultivate. As a remedy to this, an alternative method using high throughput
1165 sequencing of environmental samples was proposed by (Rimet et al., 2018b). By using this

1166 method routinely, we will be able to quickly complete the barcode reference library of the
1167 most common diatoms in the near future.

1168 4.4 Vascular plants

1169 For vascular plants the standard DNA barcode is the combination of two plastid loci, *rbcL*
1170 and *matK*. Logically, this simple fact doubles the effort needed when barcoding plants.
1171 Fortunately, some national campaigns of flora barcoding have been developed in the last
1172 decade (e.g. <http://botany.si.edu/projects/dnabarcoding/index.htm>;
1173 <https://botanicgarden.wales/science/saving-plants-and-fungi/dna-barcoding/>;
1174 <https://www.rbge.org.uk/science-and-conservation/scientific-and-technical-services/dna-barcoding/dna-barcoding-britains-liverworts/>) and the vascular plant species used for water
1176 quality assessments are well represented in the public databases (BOLD, GenBank), with
1177 barcodes registered for more than 83% of the species. The gap-analysis tool on BOLD that
1178 was used here does not require both loci to be barcoded for plants, as it reports the
1179 percentage of barcoded species regardless of whether sequences exist for both loci or only
1180 one. Only a manual check on the public data in BOLD could overcome this problem, whereas
1181 no information can be obtained for private data about the barcoding marker used. With a total
1182 of 515 barcoded species, the locus *rbcL* is better represented than *matK* (449 species).
1183 Amplification and sequencing of the *matK* barcoding region is widely known to be difficult due
1184 to high sequence variability in the primer binding sites (Hollingsworth et al., 2011).
1185 Considerable efforts have been made for developing efficient primers across multiple
1186 angiosperm families, as reflected in the recent study published by Heckenhauer and
1187 colleagues (Heckenhauer et al., 2016).

1188 In order to have a complete evaluation of the state of DNA barcode data for the
1189 macrophytes, analyses should also be performed for charophytes and bryophytes. One
1190 should, however, be aware that the situation is far from simple. A universal DNA barcode has
1191 yet to be identified for bryophytes, for which commonly used markers have low amplification-
1192 sequencing success or lack of resolving power at the species level (Hassel et al., 2013). As
1193 for the charophytes, species morphological delineation might be complicated given the
1194 plasticity of the discriminatory characters. Recent studies based on DNA barcode analyses
1195 showed that differentiation of closely related *Chara* species is not always possible and
1196 questioned the relevance of certain morphological traits in the species differentiation
1197 (Schneider et al., 2015), by highlighting an incomplete process of speciation (Nowak et al.,
1198 2016).

1199

1200 4.5 Freshwater macroinvertebrates
1201 Macroinvertebrates are central BQEs in freshwater biomonitoring programs. Our barcode
1202 gap-analyses of the BOLD reference library, including data mined from GenBank, show that
1203 while there is comparatively few species missing sequences in some insect orders (e.g.
1204 Hemiptera, Odonata and Trichoptera), other taxonomic groups lack barcodes for a majority
1205 of the regularly monitored species (e.g. Platyhelminthes). Diptera, the most species rich
1206 group used in biomonitoring in Europe, had a fairly low coverage with only about 60% of the
1207 species represented in the reference libraries. This result is similar to what was recorded in a
1208 gap-analysis of the North American freshwater invertebrates (Curry et al., 2018), although
1209 their analysis was done on genus-level. In a barcode-gap analysis of the Great Lakes fauna
1210 Trebitz et al. (2015) found that rotifers, annelids and mites had particularly low coverage,
1211 while about 70% of all insect species were represented by barcodes in BOLD. While these
1212 numbers might have changed by now, it is interesting to see that the coverage of mites and
1213 annelids appears better in Europe, while insects are slightly better covered for the Great
1214 Lakes. Generally, in our results, there is a pronounced increase in taxonomic coverage when
1215 private data in BOLD and GenBank data are included. This is particularly obvious for
1216 Annelida, Arachnida, Crustacea and Mollusca (Fig. 8). It should also be noted that while
1217 species-level coverage is low for some groups, coverage often increases at higher taxonomic
1218 ranks. This is of relevance, as some taxonomic groups are only reported at the genus, family
1219 or even order level by several countries. Below we discuss some of the characteristics
1220 observed for each major taxonomic group.

1221 4.5.1 Insects
1222 Insects are among the or even the most important and most often monitored organisms in
1223 freshwater assessments. This is reflected by both a high number of countries monitoring
1224 insects and a high number of monitored species in national monitoring checklists. However,
1225 the taxonomic level applied as well as the number of monitored taxa differs vastly among
1226 countries. The differences typically reflect the national monitoring programs (Birk et al.,
1227 2012a; Kelly et al., 2015) and hinder a direct comparison of countries in many cases
1228 (requiring sophisticated intercalibrations) and also affect the overall gap-analysis result.
1229 Almost two-thirds (65.7%) of the monitored insect species are barcoded. When looking at the
1230 taxa with the lowest barcode coverage, it becomes apparent that most of the missing species
1231 (70%) belong to Diptera, of which 72.9% are exclusively monitored in a single country
1232 (Czech Republic). Excluding only these missing Diptera species from the gap-analysis
1233 increases the overall coverage from 65.7% to 79.7% of the species, rendering the observed
1234 gap in the barcode coverage partly a problem resulting from one excessively long national
1235 checklist. This is further supported by the fact that otherwise, on average, 88.5% of the

1236 monitored species across all other surveyed countries have sequences in the reference
1237 libraries. However, similar to the observations made by Trebitz et al. (2015) for the Great
1238 Lakes fauna, the barcode coverage is significantly reduced when considering species that
1239 are represented by at least five barcodes. Moreover, since regional coverage in barcode
1240 reference libraries is important to account for the genetic diversity that is correlated with
1241 geographic distance (Bergsten et al., 2012), geographic coverage maps (Fig. 10, Suppl. Figs
1242 3-9) can be useful to identify priority areas when filling gaps in the barcode library. For some
1243 countries (e.g. Georgia), the low coverage of barcoded species can be explained by many
1244 unique species in their national checklist. In such cases, regional representation in the
1245 barcode library is crucial for implementation of DNA barcoding in freshwater biomonitoring.

1246 One obvious discrepancy was observed for the common mayfly *Ephemera danica*. While this
1247 species is one of the two most monitored species, there are only 15 available barcodes in
1248 BOLD despite there being 151 registered records. The low sequencing success of this
1249 species can be explained by suboptimal lab protocols (e.g. primer cocktails), and better
1250 representation of this commonly monitored species on BOLD could probably be achieved
1251 through protocol optimization. In conclusion, even if gaps still need to be closed, the
1252 reference databases for insects in Europe are well developed making this group already
1253 qualified for monitoring through DNA metabarcoding in several countries (e.g. (Morinière et
1254 al., 2017)).

1255 4.5.2 Arachnids

1256 Aquatic arachnids are not commonly monitored in Europe, at least not for the WFD. The
1257 most species-rich group, water mites, are well suited for monitoring environmental change of
1258 many habitats (Cantonati et al., 2006; Gerecke and Lehmann, 2005). Species-level
1259 identification using molecular tools will make information from this group more readily
1260 available in the near future. Currently, most of the barcode data on water mites in BOLD are
1261 private, but the coverage is relatively high (Fig. 9B) thanks to efforts in the Netherlands and
1262 Norway (pers. obs.). Barcode data has revealed taxonomic challenges in water mites, as
1263 revealed by the 18 specimens of *Lebertia porosa* from Norway that comprise 7 BINs (Stur,
1264 2017), and show a mean intraspecific p-distance of 11.7% (max 18.5%). Knowing that this
1265 species has currently 27 taxonomic synonyms, it will need some efforts to disentangle the
1266 names potentially associated with each genetic cluster. For the species *Hygrobates fluviatilis*
1267 a similar situation was solved with the help of DNA barcodes (Pešić et al., 2017). It is notable
1268 that the divergence of lineages with potentially different environmental preferences within the
1269 *H. fluviatilis* complex would not have been easily discovered without the comparison of
1270 sequence data in a barcode library.

1271 4.5.3 Crustaceans
1272 Crustaceans, predominantly malacostracans, are quite commonly monitored in European
1273 countries. However, the level of their taxonomic identification varies a lot from country to
1274 country, and depends on the crustacean group considered. The species are generally well
1275 covered in BOLD, however, almost half of them are represented by private barcodes only. As
1276 such, they potentially form parts of large datasets deposited in BOLD in result of ongoing
1277 studies, which eventually will be published soon. Still a comprehensive DNA barcode library
1278 for European freshwater crustaceans, such as the one published for marine crustaceans
1279 (Raupach et al., 2015), is far from completion. Yet, there are numerous recent publications
1280 providing a wealth of DNA barcode sequences as a side effect of phylogeographic or
1281 taxonomic studies, revealing the presence of high cryptic species diversity in numerous
1282 morphospecies (e.g. (Christodoulou et al., 2012; Mamos et al., 2016). For groups such as
1283 amphipods, publication of barcodes along with descriptions of new species and cryptic
1284 lineages has become almost a rule (e.g. (Rudolph et al., 2018). Thus the prognosis for
1285 further extending the reference libraries in a foreseeable future is positive.

1286 4.5.4 Annelids
1287 Despite the fact that numerous species of annelids are monitored in European countries,
1288 they are poorly covered in BOLD and most of the barcodes are kept private. A substantial
1289 share of barcode sequences mined from GenBank only. It seems that so far, there is no
1290 general habit of using BOLD as a repository for sequence data, even though COI barcodes
1291 were proven useful for identification of pseudo-cryptic and cryptic species of medicinal
1292 leeches almost a decade ago (Phillips and Siddall, 2009; Siddall Mark et al., 2007). Soon
1293 thereafter, an incongruence between morphological and molecular species boundaries was
1294 proven for *Erpobdella* leeches (Koperski et al., 2011). More recent studies revealed
1295 substantial cryptic diversity within several genera and species of freshwater oligochaetes
1296 (e.g. (Liu et al., 2017; Martin et al., 2018; Martinsson et al., 2013; Martinsson and Erseus,
1297 2018). Thus, it appears that DNA barcoding would be immensely beneficial for identification
1298 of annelids in biomonitoring.

1299 4.5.5 Molluscs
1300 A remarkable finding for freshwater molluscs was their comparatively high number of DNA
1301 barcodes deposited in GenBank, and not in BOLD. This pattern can be interpreted in terms
1302 of early initiated molecular taxonomic endeavours in the pre-BOLD era, or by a community-
1303 behaviour of submitting sequences to GenBank rather than to BOLD (e.g. (Benke et al.,
1304 2011; Prie et al., 2012). When doing so, relevant metadata might be omitted or not
1305 immediately linked to the barcode. Thus, direct BOLD submissions are highly encouraged.
1306 Furthermore, a considerable proportion of frequently listed, and presumably widely

1307 distributed species do not yet have any barcode data available. This lack of data might be
1308 even more pronounced, as several integrative taxonomic studies on freshwater molluscs
1309 indicate that widely distributed morphospecies often comprise complexes of distinctly defined
1310 genetic lineages (cryptic species). A good example is *Ancylus fluviatilis*, the most often listed
1311 freshwater mollusc in our dataset, which actually constitutes a complex of at least six cryptic
1312 species (Albrecht et al., 2006; Pfenninger et al., 2003; Weiss et al., 2018).

1313 4.5.6 Platyhelminthes and Nematoda

1314 Both flatworms and nematodes are diverse and of indicative value. While some countries do
1315 register Platyhelminthes in existing surveys, Nematoda are generally neglected. For
1316 nematodes in the Palaearctic, 1580 species should be relevant for the WFD (Eisendle et al.
1317 2017). Thus, a barcode library of freshwater nematodes can have a potentially large impact
1318 on the use of this organism group in future biomonitoring.

1319 4.6 Freshwater fish

1320 With about 88% coverage with at least one DNA barcode in BOLD, European freshwater
1321 fishes are well represented and the species being reliably identifiable based on COI in real
1322 world applications. While 47 species have only one specimen with DNA barcode deposited in
1323 BOLD, a large proportion (two thirds) is available with at least five individuals. The coverage
1324 with 12S data is much lower and only a third of the species was found to be available in
1325 public databases.

1326 About three-fourth of all European freshwater fish species fall into the three higher taxa
1327 presented in more detail (Perciformes, Cyprinidae and Salmoniformes), which contain
1328 commercially important game and food species (perch, pike-perch, carp, bream, roach, trout,
1329 whitefishes and chars). The largest and most widespread family is Cyprinidae, which has its
1330 (extant) species completely covered by DNA barcodes in BOLD, with only five species
1331 missing - all of which being regarded or listed by IUCN as extinct (*Alburnus danubicus*,
1332 *Chondrostoma scodrense*, *Iberocypris palaciosi*, *Pelasgus epiroticus*, *Romanogobio antipai*).
1333 Especially given the potential of molecular identification and detection tools for non-invasive
1334 and highly sensitive approaches to assess a species' presence or even abundance (e.g.
1335 (Ushio et al., 2018), we argue that it is also important to cover those species in the
1336 databases, which are thought to be extinct. Among the ten completely missing perciform
1337 species are four tad-pole gobies (*Benthophilus* spp.), two sculpins (*Cottus* spp.) and two
1338 dwarf gobies (*Knipowitschia* spp.) with predominantly Eastern European and putative
1339 Caspian basin distributions, areas which are generally less well studied and explored from an
1340 ichthyologist's perspective. An exemption is the elusive *Zingel balcanicus* from Macedonia
1341 and Greece (protected through Annex II of the European Union Habitats Directive

1342 92/43/EEC), which has been re-discovered and anatomically analysed recently (Arsovská et
1343 al., 2014), but as no DNA-material has been secured cannot be assessed via molecular tools
1344 at the moment.


1345 The largest gaps in the reference database pertain to the salmoniform group with chars
1346 (*Salvelinus* spp. - 19 species), trouts (*Salmo* spp. - 9 species) and whitefishes (*Coregonus*
1347 spp. - 16 species). While these groups contain many commercially exploited species, they
1348 are known to be notoriously difficult to identify based on general morphology (Kottelat and
1349 Freyhof, 2007), but also applying standard DNA barcoding routines (Geiger et al., 2014;
1350 Knebelsberger et al., 2014a). This is most likely due to the presence of post-glacially evolved
1351 species flocks, which are only little differentiated genetically - at least judging from the groups
1352 that have been studied so far (Dierking et al., 2014; Hudson et al., 2011; Vonlanthen et al.,
1353 2012). From a geographic point of view, most missing species occur in Scandinavia and UK
1354 (chars), the Alp region (whitefishes), and the Eastern Mediterranean (trout).

1355 4.7 Quality measures for DNA barcode reference libraries

1356 In a barcoder's perfect world, all species on Earth would be identifiable based on their DNA
1357 barcodes. However, this ideal conception is hampered by several biological and human-
1358 made phenomena. For example, time since speciation might be rather short, and the
1359 universal marker considered not diverse (i.e. not informative) enough to resolve this
1360 speciation event (e.g. (Weiss et al., 2018)). Additionally, gene flow might be still possible,
1361 even between less closely related species, leading to the (unidirectional) introgression of
1362 genomes, and hence to the (partial) intermixture of barcodes (e.g. (Weigand et al., 2017)).
1363 Besides these natural processes complicating the diagnostic utility of DNA barcodes, human-
1364 made artefacts during reference library development directly affect the reliability of DNA
1365 barcoding to correctly identify specimens to species. This includes identification errors,
1366 sequence contamination, incomplete reference data or inadequate data management. It was
1367 thus not surprising that subsequent to the proposal of the term "DNA barcode" (Floyd et al.,
1368 2002; Hebert et al., 2003a; Hebert et al., 2003b), special emphasis was laid on formal
1369 standardisation guidelines for DNA barcodes in the context of reference library development
1370 (see e.g. (Ratnasingham and Hebert, 2007; Walters and Hanner, 2006)). Those include the
1371 criteria that any 'formal' barcode sequence: a) derives from an accepted gene region, b)
1372 meets certain sequence quality standards (e.g. demonstrating at least 75% of contiguous
1373 high quality bases or less than 1% Ns and being associated with trace files and primer
1374 information), c) is linked to a voucher specimen in a major collection, and d) ideally but not
1375 always mandatorily possesses further collection and identification details (i.e. georeference
1376 data, name of collector and identifier). Since then several biomonitoring and assessment

1377 applications have moved from classical single specimen identifications to highly parallelized
1378 characterisations of communities via DNA metabarcoding (Leese et al., 2018). Given the
1379 often overwhelming quantity of 'big biodiversity data' and automated pipelines in those HTS
1380 approaches, data quality aspects of DNA barcode references gain an even higher relevance.
1381 Thus, some research communities, such as European diatom experts have worked with the
1382 European Standardization Committee to publish a methodology as a first step towards
1383 standardization of reference barcode libraries for diatoms (CEN, 2018).

1384 In principle, two quality components can be distinguished: Quality assurance (QA) is
1385 process-orientated, providing and maintaining quality standards for DNA barcodes and
1386 reference libraries. Quality control (QC), on the other hand, is user-orientated, enabling the
1387 cross-validation of taxonomic assignments or flagging of doubtful barcodes. More generally
1388 speaking, QA and QC measures can be seen as internal (or preventive) and external (or
1389 reactive) curation of reference libraries, respectively (Fig. 17). The implementation of QA
1390 measures during reference library development is the first important step for a sustainable
1391 data quality management. Linked to a valid taxonomy, formally-correct barcode sequences
1392 are deposited in line with (digital) voucher specimens and extensive metadata information.
1393 The taxonomic backbone should be regularly updated with modifications being visible to the
1394 users. An open access and fully transparent reference library allowing for versioning of
1395 barcode collections and the possibility to track taxonomic changes can be seen as the gold
1396 standard here. Simultaneously, this will allow a more sophisticated QC by the barcoding
1397 community. Library entries can be flagged for contamination and the most recent taxonomic
1398 changes (i.e. newly described species, integrative revisions) incorporated into the reference
1399 library taxonomic backbone more easily. A library which communicates with other ecological
1400 or geographic datasets and which provides access to the full data lifecycle from deposition to
1401 publication of data will further smoothen the integrative utilisation of barcode datasets. The
1402 generation of custom reference libraries and their annotation with digital object identifiers
1403 (DOI) finally can account for transparency and the specific demands of the users.



1404

1405 Figure 17. An overview of the reference library steps 'development', 'maintenance' and
1406 'utilisation', their quality assurance (QA) and quality control (QC) measures.

1407

1408 Although a variety of QA/QC measures are implemented at the stages of reference library
1409 development and maintenance, improvements are possible for the QA/QC components
1410 during reference library utilisation. This holds especially true for complex metabarcoding
1411 studies based on multiplexed HTS data. In most of those cases, taxonomic identifications are
1412 achieved by semi-automatically comparing the clustered or individual metabarcodes with a
1413 reference library and applying flexible similarity thresholds. The sequence is thus linked to a
1414 Linnaean name, e.g. by a 2% similarity threshold for species-level identification of a
1415 molecular operational taxonomic unit (MOTU). By doing so, only the availability and match of
1416 barcodes are considered, neglecting any additional metadata. Yet, knowledge about the
1417 number of barcoded specimens per species, their morphological identifiers and the
1418 distribution area covered, are likewise valuable information and should be available for direct
1419 QC. Special cases of mito-nuclear discordance, the number of already known MOTUs for a
1420 given Linnaean species name and 'extraordinary' barcodes such as those originating from
1421 type specimens should be additionally highlighted in the output results. All this combined
1422 information could be used to establish an evaluation system for metabarcode identifications,

1423 sorting taxonomic results by their plausibility and hence establishing further QA for reference
1424 library identification performance.

1425 The ultimate reference library goal is to link a DNA barcode to a voucher specimen,
1426 accompanying metadata and its Linnaean name. However, more and more frequently, a
1427 reverse taxonomic approach is applied for the generation and deposition of reference
1428 barcodes, e.g. the 'reverse BIN taxonomy' in BOLD. During this process, a sequence with a
1429 taxonomic annotation above the species-level (e.g. family or genus) is included in the
1430 database and identified by the already available barcodes. For instance, a chironomid
1431 specimen (BOLD sequence page: GMGMC1513-14) of a vouchered collection in BOLD
1432 bears the species name *Polypedilum convictum* including a specimen identifier in its
1433 metadata. Only when accessing the internal specimen page (BOLD: BIOUG16529-F11) the
1434 identification method information "BIN Taxonomy Match" is given, however, without
1435 presenting the original morphological identification level. Strictly speaking, species
1436 identification through DNA barcoding has generated this 'reference', and not expert
1437 identification by morphology. Subsequently, this sequence is considered on species-level in
1438 the database and a more precise initial morphological identification is pretended. At present,
1439 reverse BIN taxonomy sequences (see Supplement 3) a) can be found in up to 16% and
1440 20% of the monitored species of a taxonomic group (Diptera and Megaloptera, respectively),
1441 b) represent up to 61% and 82% of a higher taxon's public barcodes (Annelida and Diptera)
1442 and up to 20% to 59% of all barcodes (Coleoptera and Diptera), c) can be found in species
1443 with only few public barcodes (e.g. three out of five *Crangonyx pseudogracilis* sequences are
1444 reverse BIN taxonomy sequences) and d) represent more than half (e.g. 38/75 for *Mytilus*
1445 *edulis*) or all (e.g. 35/35 for *Lumbricillus rivalis*) public sequences of a species. As such,
1446 wrong species-level DNA barcodes are potentially introduced, with often incorrect metadata
1447 for 'specimen identification' going along with them. They must be seen as a geographic
1448 reference for a MOTU rather than as a reliable taxonomic reference. The 'reverse BIN
1449 taxonomy' practice will also bias the evaluation system for the interpretation of metabarcode
1450 identification results.

1451 An auditing and annotation system for reference libraries of DNA barcodes has been
1452 originally proposed by Costa et al. (Costa et al., 2012), and later updated by Oliveira et al.
1453 (2016) to accommodate the BIN system. The application of this QC system was particularly
1454 adequate for reference libraries of marine fishes (Cariani et al., 2017; Oliveira et al., 2016),
1455 but it has been equally applied to other taxa, such as Gastropoda (Borges et al., 2016) and
1456 Polychaeta (Lobo et al., 2016). Essentially, this system lies in the verification of the
1457 concordance between morphology-based identifications and BIN-based sequence clusters –
1458 within a given reference library (e.g. fishes of Europe) – and the subsequent annotation of

1459 each species with one to five available grades, i.e. ranging from maximum concordance
1460 (grade A) to complete discordance (grade E). Annotated grades are ought to be regularly
1461 reviewed and updated as required. Rather than requiring decisions about the taxonomic
1462 status and validity of a given species, this procedure simply considers the annotation of the
1463 level of congruency between morphological and molecular data. The auditor only needs to
1464 make decisions on the grade of congruency to apply.

1465 The auditing system of Costa et al. (Costa et al., 2012) differs in a number of ways from the
1466 "BIN discordance report" tool implemented in BOLD, which only flags BINs that include
1467 records with more than one taxon name, but does not point out cases of the same species
1468 occurring in multiple BINs (note that BIN discordance reports of all data on BOLD only is
1469 available in BOLD v3). Also, because the BIN discordance report is an automated computer-
1470 based procedure, it does not distinguish true discordance from misspelled species names,
1471 synonyms, or patent cases of discordance resulting from cross-contamination or mislabelling
1472 of samples (e.g. (Knebelsberger et al., 2014b)). Hence, as a result of the auditing and
1473 annotation framework, end-users will have an indication of the reliability and accuracy of a
1474 given species match, and will be immediately alerted for records with insufficient data, or
1475 uncertain or misleading matches.

1476 5. Conclusions and Recommendations

1477 For marine macroinvertebrates, future efforts should focus initially on filling the gaps of the
1478 AMBI checklist, especially those more dominant in the datasets, which greatly influence the
1479 AMBI result (Aylagas et al., 2014), while keeping the long-term goal of completing the ERMS
1480 checklist. For freshwater macroinvertebrates, species-groups that are widely used in WFD
1481 monitoring such as Annelida, Crustacea, Insecta and Mollusca should be prioritized. For
1482 marine groups, gaps should be filled first to maximize phylogenetic representativeness,
1483 thereby yielding to the collection of reference barcodes of representative species from
1484 missing orders, then missing families, and so forth down to genera. This strategy aims to
1485 provide, at the very least, an interim proximate taxonomic assignment for metabarcoding
1486 reads lacking species level matches. However, most of the work has still to be done at the
1487 species level, because within the same genus, there are species belonging to different
1488 ecological groups, and thus the identification at species level is mandatory for reliable EQS
1489 and environmental status assessment. Hence, subsequent efforts should address species
1490 level completion, focusing on the taxonomic groups with greater gaps, as well as on the taxa
1491 used in AMBI's ecological categories. The increase in the number of DNA barcodes for less
1492 barcoded species must also be pursued, since most of the taxonomic groups have less than
1493 5 barcodes/species in the reference libraries. Attempts to include representative specimens

1494 across the geographic distribution range shall be made for missing species in the reference
1495 libraries. Particular care must be taken regarding the QA/QC of the reference barcode
1496 records to be produced, as failure to do so will limit their application, render them useless, or
1497 even introduce wrong outcomes. Moreover, as new HTS techniques are developed to obtain
1498 full-length reference barcodes from old type material (Prosser et al., 2016), this strategy
1499 should be used to resolve the taxonomy and names of key taxa used in biomonitoring.

1500 6. Acknowledgements

1501 This paper is a deliverable of the European Cooperation in Science and Technology (COST)
1502 Action DNAqua-Net (CA15219) Working Group 1, lead by Torbjørn Ekrem and Fedor
1503 Čiampor. Thanks to the University of Minho and University of Pécs for hosting workshops
1504 and working group meetings. We also thank staff at National Environment Agencies and
1505 others that provided national checklists of taxa used in biomonitoring, and otherwise assisted
1506 with checklist proof-reading: Jarmila Makovinská and Emília Mišíková Elexová (Slovakia);
1507 Steinar Sandøy and Dag Rosland (Norway); Mišel Jelić (Croatia); Marlen Vasquez (Cyprus);
1508 Adam Petrusk (Czech Republic); Kristel Panksep (Estonia); Panagiotis Kaspiditis (Greece);
1509 Matteo Montagna (Italy); Marija Katarzyte (Lithuania); Ana Rotter (Slovenia); Rosa Trabajo
1510 (Spain); Florian Altermatt (Switzerland); Kristian Meissner (Finland), Rigers Bakiu (Albania),
1511 Valentina Stamenkovic and Jelena Hinic (Macedonia); Patricia Mergen (Belgium); Gael
1512 Denys & the French Biodiversity Agency (France); Mary Kelly-Quinn (Ireland); Cesare Mario
1513 Puzzi (Italy); Carole Fitzpatrick (United Kingdom); Simon Vitecek (Austria); Ana Filipa Filipe
1514 (Portugal); Peter Anton Stæhr & Anne Winding (Denmark); Michael Monaghan (Germany);
1515 Alain Dohet, Lionel L'Hoste, Nora Welschbillig & Luc Ector (Luxembourg), Lujza Keresztes,
1516 (Romania). The authors also want to thank Dirk Steinke for providing the original European
1517 ERMS list for marine taxa and Florian Malard for comments on the manuscript. The
1518 preparation of the AMBI checklist was carried out in the scope of a Short-term Scientific
1519 Mission (ECOST-STSM-CA15219-150217-082111) granted to Sofia Duarte visiting AZTI
1520 Tecnalia, Spain.

1521 7. Supplementary files

1522 Supplement 1. Overview of obtained checklists of taxa used in national biomonitoring of
1523 aquatic ecosystems.
1524 Supplement 2. Raw data from gap-analysis of all taxonomic groups showing: Species,
1525 Countries, Barcodes BOLD (public, private), GenBank records.
1526 Supplement 3. Reverse BIN taxonomy statistics in BOLD.

1527 8. References

1528 Albrecht C, Trajanovski S, Kuhn K, Streit B, Wilke T. Rapid evolution of an ancient lake
1529 species flock: Freshwater limpets (Gastropoda : Aculyidae) in the Balkan Lake Ohrid.
1530 Organisms Diversity & Evolution 2006; 6: 294-307.

1531 Arsovská J, Ristovska M, Kostov V, Prelic D, Slavevska-Stamenkovic V. Osteological
1532 description of Zingel balcanicus (Teleostei: Percidae). Biologia 2014; 69: 1742-1756.

1533 Aylagas E, Borja Á, Muxika I, Rodríguez-Ezpeleta N. Adapting metabarcoding-based benthic
1534 biomonitoring into routine marine ecological status assessment networks. Ecological
1535 Indicators 2018; 95: 194-202.

1536 Aylagas E, Borja Á, Rodríguez-Ezpeleta N. Environmental Status Assessment Using DNA
1537 Metabarcoding: Towards a Genetics Based Marine Biotic Index (gAMBI). PLoS ONE
1538 2014; 9: e90529.

1539 Balian EV, Segers H, Léveque C, Martens K. The Freshwater Animal Diversity Assessment:
1540 an overview of the results. Hydrobiologia 2008; 595: 627-637.

1541 Barbier EB. Progress and Challenges in Valuing Coastal and Marine Ecosystem Services.
1542 Review of Environmental Economics and Policy 2012; 6: 1-19.

1543 Barbier EB. Marine ecosystem services. Current Biology 2017; 27: 507-510.

1544 Barbour MT, United States. Environmental Protection Agency. Office of Water. Rapid
1545 bioassessment protocols for use in streams and wadeable rivers periphyton, benthic
1546 macroinvertebrates, and fish. Washington, DC: U.S. Environmental Protection
1547 Agency, Office of Water, 1999.

1548 Barco A, Evans J, Schembri PJ, Taviani M, Oliverio M. Testing the applicability of DNA
1549 barcoding for Mediterranean species of top-shells (Gastropoda, Trochidae, Gibbula
1550 s.l.). Marine Biology Research 2013; 9: 785-793.

1551 Barco A, Raupach MJ, Laakmann S, Neumann H, Knebelsberger T. Identification of North
1552 Sea molluscs with DNA barcoding. Molecular Ecology Resources 2016; 16: 288-297.

1553 Benke M, Brandle M, Albrecht C, Wilke T. Patterns of freshwater biodiversity in Europe:
1554 lessons from the spring snail genus *Bythinella*. Journal of Biogeography 2011; 38:
1555 2021-2032.

1556 Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, et al. GenBank.
1557 Nucleic Acids Research 2013; 41: 36-42.

1558 Bergsten J, Bilton DT, Fujisawa T, Elliott M, Monaghan MT, Balke M, et al. The Effect of
1559 Geographical Scale of Sampling on DNA Barcoding. Systematic Biology 2012; 61:
1560 851-869.

1561 Birk S, Bonne W, Borja A, Brucet S, Courrat A, Poikane S, et al. Three hundred ways to
1562 assess Europe's surface waters: An almost complete overview of biological methods
1563 to implement the Water Framework Directive. Ecological Indicators 2012a; 18: 31-41.

1564 Birk S, Bonne W, Borja A, Brucet S, Courrat A, Poikane S, et al. Three hundred ways to
1565 assess Europe's surface waters: An almost complete overview of biological methods
1566 to implement the Water Framework Directive. Ecological Indicators 2012b; 18: 31-41.

1567 Borges LM, Hollatz C, Lobo J, Cunha AM, Vilela AP, Calado G, et al. With a little help from
1568 DNA barcoding: investigating the diversity of Gastropoda from the Portuguese coast.
1569 Scientific Reports 2016; 6: 20226.

1570 Borgwardt F, Robinson L, Trauner D, Teixeira H, Nogueira AJA, Lillebo AI, et al. Exploring
1571 variability in environmental impact risk from human activities across aquatic
1572 ecosystems. Science of the Total Environment 2019; 652: 1396-1408.

1573 Borja A, Elliott M, Andersen JH, Cardoso AC, Carstensen J, Ferreira JG, et al. Good
1574 Environmental Status of marine ecosystems: What is it and how do we know when
1575 we have attained it? Marine Pollution Bulletin 2013; 76: 16-27.

1576 Borja A, Franco J, Pérez V. A Marine Biotic Index to Establish the Ecological Quality of Soft-
1577 Bottom Benthos Within European Estuarine and Coastal Environments. Marine
1578 Pollution Bulletin 2000; 40: 1100-1114.

1579 Borja A, Josefson AB, Miles A, Muxika I, Olsgard F, Phillips G, et al. An approach to the
1580 intercalibration of benthic ecological status assessment in the North Atlantic
1581 ecoregion, according to the European Water Framework Directive. *Marine Pollution*
1582 *Bulletin* 2007; 55: 42-52.

1583 Borja A, Miles A, Occhipinti-Ambrogi A, Berg T. Current status of macroinvertebrate methods
1584 used for assessing the quality of European marine waters: implementing the Water
1585 Framework Directive. *Hydrobiologia* 2009; 633: 181-196.

1586 Brodin Y, Ejdung G, Strandberg J, Lyrholm T. Improving environmental and biodiversity
1587 monitoring in the Baltic Sea using DNA barcoding of Chironomidae (Diptera).
1588 *Molecular Ecology Resources* 2012; 13: 996-1004.

1589 Burek P, Satoh Y, Fischer G, Kahil MT, Scherzer A, Tramberend S, et al. Water Futures and
1590 Solution - Fast Track Initiative (Final Report), IIASA, Laxenburg, Austria, 2016.

1591 Cantonati M, Gerecke R, Bertuzzi E. Springs of the Alps - Sensitive ecosystems to
1592 environmental change: From biodiversity assessments to long-term studies.
1593 *Hydrobiologia* 2006; 562: 59-96.

1594 Carew ME, Nichols SJ, Batovska J, St Clair R, Murphy NP, Blacket MJ, et al. A DNA barcode
1595 database of Australia's freshwater macroinvertebrate fauna. *Marine and Freshwater*
1596 *Research* 2017; 68: 1788-1801.

1597 Cariani A, Messinetti S, Ferrari A, Arculeo M, Bonello JJ, Bonnici L, et al. Improving the
1598 Conservation of Mediterranean Chondrichthyans: The ELASMOMED DNA Barcode
1599 Reference Library. *PLoS One* 2017; 12: e0170244.

1600 Carstensen J, Lindegarth M. Confidence in ecological indicators: A framework for quantifying
1601 uncertainty components from monitoring data. *Ecological Indicators* 2016; 67: 306-
1602 317.

1603 CEN. Water quality - Technical report for the management of diatom barcodes, Brussels,
1604 Belgium, 2018, pp. 1-11.

1605 Christodoulou M, Antoniou A, Magoulas A, Koukouras A. Revision of the freshwater genus
1606 *Atyaephyra* (Crustacea, Decapoda, Atyidae) based on morphological and molecular
1607 data. *Zookeys* 2012: 53-110.

1608 Civade R, Dejean T, Valentini A, Roset N, Raymond J-C, Bonin A, et al. Spatial
1609 Representativeness of Environmental DNA Metabarcoding Signal for Fish Biodiversity
1610 Assessment in a Natural Freshwater System. *Plos One* 2016; 11: e0157366.

1611 Clarke RT. Estimating confidence of European WFD ecological status class and WISER
1612 Bioassessment Uncertainty Guidance Software (WISERBUGS). *Hydrobiologia* 2013;
1613 704: 39-56.

1614 Cook CDK, Gut BJ, Rix EM. Water plants of the world : a manual for the identification of the
1615 genera of freshwater macrophytes: The Hague : Junk, 1974.

1616 Costa FO, deWaard JR, Boutilier J, Ratnasingham S, Dooh RT, Hajibabaei M, et al.
1617 Biological identifications through DNA barcodes: the case of the Crustacea. *Canadian*
1618 *Journal of Fisheries and Aquatic Sciences* 2007; 64: 272-295.

1619 Costa FO, Landi M, Martins R, Costa MH, Costa ME, Carneiro M, et al. A ranking system for
1620 reference libraries of DNA barcodes: application to marine fish species from Portugal.
1621 *PLoS One* 2012; 7: e35858.

1622 Costello M. Developing Species Information Systems: The European Register of Marine
1623 Species (ERMS). Vol 13, 2000.

1624 Curry CJ, Gibson JF, Shokralla S, Hajibabaei M, Baird DJ. Identifying North American
1625 freshwater invertebrates using DNA barcodes: are existing COI sequence libraries fit
1626 for purpose? *Freshwater Science* 2018; 37: 178-189.

1627 Davy-Bowker J, Clarke R, Corbin T, Vincent H, Pretty J, Hawczak A, et al. River Invertebrate
1628 Classification Tool. Final report. Scotland & Northern Ireland Forum for Environmental
1629 Research, Edinburgh, UK, 2008, pp. 276.

1630 Dierking J, Phelps L, Praebel K, Ramm G, Prigge E, Borcherding J, et al. Anthropogenic
1631 hybridization between endangered migratory and commercially harvested stationary
1632 whitefish taxa (*Coregonus* spp.). *Evolutionary Applications* 2014; 7: 1068-1083.

1633 Dudgeon D, Arthington AH, Gessner MO, Kawabata ZI, Knowler DJ, Lévêque C, et al.
1634 Freshwater biodiversity: importance, threats, status and conservation challenges.
1635 Biological Reviews 2006; 81: 163-182.

1636 Ekrem T, Willassen E, Stur E. A comprehensive DNA sequence library is essential for
1637 identification with DNA barcodes. Molecular Phylogenetics and Evolution 2007; 43:
1638 530-542.

1639 Eisendle, U, Decraemer W, Abebe E, De Ley P (2017): A global checklist of freshwater
1640 nematodes - <http://fada.biodiversity.be/group/show/6> (accessed March 2019).

1641 European Commission. Directive 2000/60/EC of the European Parliament and of the Council
1642 of 23rd October 2000 establishing a framework for Community action in the field of
1643 water policy. Official Journal of the European Communities 2000; 327: 1-72.

1644 European Commission. Directive 2008/56/EC of the European Parliament and of the Council
1645 establishing a framework for community action in the field of marine environmental
1646 policy (Marine Strategy Framework Directive). Official Journal of the European
1647 Communities 2008; L164: 19-40.

1648 European Commission. Commission Decision (EU) 2017/848 of 17 May 2017 laying down
1649 criteria and methodological standards on good environmental status of marine waters
1650 and specifications and standardised methods for monitoring and assessment, and
1651 repealing Decision 2010/477/EU. Official Journal of the European Communities 2017;
1652 L125: 43-74.

1653 European Commission. Commission Decision (EU) 2018/229 of 12 February 2018
1654 establishing, pursuant to Directive 2000/60/EC of the European Parliament and of the
1655 Council, the values of the Member State monitoring system classifications as a result
1656 of the intercalibration exercise and repealing Commission Decision 2013/480/EU.
1657 Official Journal of the European Communities 2018; L47: 1-91.

1658 Floyd R, Abebe E, Papert A, Blaxter M. Molecular barcodes for soil nematode identification.
1659 Molecular Ecology 2002; 11: 839-850.

1660 Fricke R, Eschmeyer WN, van der Laan R. Catalog of fishes: genera, species, references.
1661 California Academy of Sciences, 2018.

1662 Geiger M, Herder F, Monaghan M, Almada V, Barbieri R, Bariche M, et al. Spatial
1663 Heterogeneity in the Mediterranean Biodiversity Hotspot Affects Barcoding Accuracy
1664 of its Freshwater Fishes, 2014.

1665 Gerecke R, Lehmann EO. Towards a long-term monitoring of Central European water mite
1666 faunas (Acari: Hydrachnidia and Halacaridae) – considerations on the background of
1667 data from 1900 to 2000. Limnologica 2005; 35: 45-51.

1668 Grall J, Glémarec M. Using biotic indices to estimate macrobenthic community perturbations
1669 in the Bay of Brest. Estuarine, Coastal and Shelf Science 1997; 44: 43-53.

1670 Haase P, Pauls SU, Schindelhütte K, Sundermann A. First audit of macroinvertebrate
1671 samples from an EU Water Framework Directive monitoring program: human error
1672 greatly lowers precision of assessment results. Journal of the North American
1673 Benthological Society 2010; 29: 1279-1291.

1674 Hassel K, Segreto R, Ekrem T. Restricted variation in plant barcoding markers limits
1675 identification in closely related bryophyte species. Mol Ecol Resour 2013; 13: 1047-
1676 57.

1677 Hebert P, Cywinski A, Ball S, deWaard J. Biological identifications through DNA barcodes.
1678 Proceedings of the Royal Society London, B 2003a; 270: 313 - 321.

1679 Hebert P, Ratnasingham S, deWaard J. Barcoding animal life: cytochrome c oxidase subunit
1680 1 divergences among closely related species. Proc Roy Soc Lond B 2003b; 270: 96 -
1681 99.

1682 Hebert PDN, Hollingsworth PM, Hajibabaei M. From writing to reading the encyclopedia of
1683 life. Philosophical Transactions of the Royal Society B: Biological Sciences 2016;
1684 371.

1685 Heckenhauer J, Barfuss MH, Samuel R. Universal multiplexable matK primers for DNA
1686 barcoding of angiosperms. Appl Plant Sci 2016; 4: 1500137.

1687 Hering D, Borja A, Carstensen J, Carvalho L, Elliott M, Feld CK, et al. The European Water
1688 Framework Directive at the age of 10: A critical review of the achievements with
1689 recommendations for the future. *Science of the Total Environment* 2010; 408: 4007-
1690 4019.

1691 Hering D, Borja A, Jones JI, Pont D, Boets P, Bouchez A, et al. Implementation options for
1692 DNA-based identification into ecological status assessment under the European
1693 Water Framework Directive. *Water Research* 2018; 138: 192-205.

1694 Hering D, Feld CK, Moog O, Ofenböck T. Cook book for the development of a Multimetric
1695 Index for biological condition of aquatic ecosystems: Experiences from the European
1696 AQEM and STAR projects and related initiatives. *Hydrobiologia* 2006; 566: 311-324.

1697 Hollingsworth PM, Forrest LL, Spouge JL, Hajibabaei M, Ratnasingham S, van der Bank M,
1698 et al. A DNA barcode for land plants. *Proceedings of the National Academy of
1699 Sciences of the United States of America* 2009; 106: 12794-12797.

1700 Hollingsworth PM, Graham SW, Little DP. Choosing and Using a Plant DNA Barcode. *Plos
1701 One* 2011; 6: e19254.

1702 Horton T, Kroh A, Ahyong S, Bailly N, Boury-Esnault N, Brandão SN, et al. World Register of
1703 Marine Species (WoRMS). WoRMS Editorial Board, 2018.

1704 Hudson AG, Vonlanthen P, Seehausen O. Rapid parallel adaptive radiations from a single
1705 hybridogenic ancestral population. *Proceedings of the Royal Society B-Biological
1706 Sciences* 2011; 278: 58-66.

1707 Hunter JD. Matplotlib: A 2D graphics environment. *Computing in Science & Engineering*
1708 2007; 9: 90-95.

1709 Häneling B, Lawson Handley L, Read DS, Hahn C, Li J, Nichols P, et al. Environmental DNA
1710 metabarcoding of lake fish communities reflects long-term data from established
1711 survey methods. *Molecular Ecology* 2016; 25: 3101-3119.

1712 Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7:
1713 Improvements in Performance and Usability. *Molecular Biology and Evolution* 2013;
1714 30: 772-780.

1715 Keck F, Vasselon V, Tapolczai K, Rimet F, Bouchez A. Freshwater biomonitoring in the
1716 Information Age. *Frontiers in Ecology and the Environment* 2017; 15: 266-274.

1717 Kelly MG, Schneider SC, King L. Customs, habits, and traditions: the role of nonscientific
1718 factors in the development of ecological assessment methods. *Wiley Interdisciplinary
1719 Reviews: Water* 2015; 2: 159-165.

1720 Kermarrec L, Franc A, Rimet F, Chaumeil P, Humbert JF, Bouchez A. Next-generation
1721 sequencing to inventory taxonomic diversity in eukaryotic communities: a test for
1722 freshwater diatoms. *Molecular Ecology Resources* 2013; 13: 607-619.

1723 Keskin E, Atar HH. DNA barcoding commercially important fish species of Turkey. *Molecular
1724 Ecology Resources* 2013; 13: 788-797.

1725 Knebelsberger T, Dunz AR, Neumann D, Geiger MF. Molecular Diversity of Germany's
1726 Freshwater Fishes and Lampreys assessed by DNA barcoding. *Molecular Ecology
1727 Resources* 2014a; 15: 562-572.

1728 Knebelsberger T, Landi M, Neumann H, Kloppmann M, Sell AF, Campbell PD, et al. A
1729 reliable DNA barcode reference library for the identification of the North European
1730 shelf fish fauna. *Mol Ecol Resour* 2014b; 14: 1060-1071.

1731 Kongsrød JA, Bakken T, Oug E, Alvestad T, Nygren A, Kongshavn K, et al. Assessing
1732 species diversity in marine bristle worms (Annelida, Polychaeta): integrating
1733 barcoding with traditional morphology-based taxonomy. *Genome* 2017; 60: 956-956.

1734 Koperski P, Milanowski R, Krzyk A. Searching for cryptic species in *Erpobdella octoculata*
1735 (L.) (Hirudinea: Clitellata): discordance between the results of genetic analysis and
1736 cross-breeding experiments. *Contributions to Zoology* 2011; 80: 85-94.

1737 Kottelat M, Freyhof J. *Handbook of European freshwater fishes*. Berlin: Cornol and Freyhof,
1738 2007.

1739 Kreamer DK. The Past, Present, and Future of Water Conflict and International Security.
1740 *Journal of Contemporary Water Research & Education* 2012; 149: 87-95.

1741 Landi M, Dimech M, Arculeo M, Biondo G, Martins R, Carneiro M, et al. DNA Barcoding for
1742 Species Assignment: The Case of Mediterranean Marine Fishes. PLOS ONE 2014; 9:
1743 e106135.

1744 Lavoie I, Dillon PJ, Campeau S. The effect of excluding diatom taxa and reducing taxonomic
1745 resolution on multivariate analyses and stream bioassessment. Ecological Indicators
1746 2009; 9: 213-225.

1747 Leese F, Altermatt F, Bouchez A, Ekrem T, Hering D, Meissner K, et al. DNAqua-Net:
1748 Developing new genetic tools for bioassessment and monitoring of aquatic
1749 ecosystems in Europe. Research Ideas and Outcomes 2016; 2: e11321.

1750 Leese F, Bouchez A, Abarenkov K, Altermatt F, Borja Á, Bruce K, et al. Why We Need
1751 Sustainable Networks Bridging Countries, Disciplines, Cultures and Generations for
1752 Aquatic Biomonitoring 2.0: A Perspective Derived From the DNAqua-Net COST
1753 Action. In: Bohan DA, Dumbrell AJ, Woodward G, Jackson M, editors. Next
1754 Generation Biomonitoring. Part 1. 58. Academic Press, 2018, pp. 63-99.

1755 Lefrancois E, Apothéloz-Perret-Gentil L, Blancher P, Botreau S, Chardon C, Crepin L, et al.
1756 Development and implementation of eco-genomic tools for aquatic ecosystem
1757 biomonitoring: the SYNAQUA French-Swiss program. Environmental Science and
1758 Pollution Research 2018; 25: 33858-33866.

1759 Liu Y, Fend SV, Martinsson S, Erséus C. Extensive cryptic diversity in the cosmopolitan
1760 sludge worm *Limnodrilus hoffmeisteri* (Clitellata, Naididae). Organisms, Diversity &
1761 Evolution 2017; 17: 477-495.

1762 Lobo J, Ferreira MS, Antunes IC, Teixeira MAL, Borges LMS, Sousa R, et al. Contrasting
1763 morphological and DNA barcode-suggested species boundaries among shallow-
1764 water amphipod fauna from the southern European Atlantic coast. Genome 2017; 60:
1765 147-157.

1766 Lobo J, Teixeira MA, Borges LM, Ferreira MS, Hollatz C, Gomes PT, et al. Starting a DNA
1767 barcode reference library for shallow water polychaetes from the southern European
1768 Atlantic coast. Mol Ecol Resour 2016; 16: 298-313.

1769 Mamos T, Wattier R, Burzynski A, Grabowski M. The legacy of a vanished sea: a high level
1770 of diversification within a European freshwater amphipod species complex driven by
1771 15 My of Paratethys regression. Molecular Ecology 2016; 25: 795-810.

1772 Mann DG, Vanormelingen P. An inordinate fondness? The number, distributions, and origins
1773 of diatom species. Journal of Eukaryote Microbiology 2013; 60: 414-20.

1774 Martin P, Martinsson S, Wuillot J, Erseus C. Integrative species delimitation and phylogeny
1775 of the branchiate worm *Branchiodrilus* (Clitellata, Naididae). Zoologica Scripta 2018;
1776 47: 727-742.

1777 Martinsson S, Achurra A, Svensson M, Erseus C. Integrative taxonomy of the freshwater
1778 worm *Rhyacodrilus falciformis* s.l. (Clitellata: Naididae), with the description of a new
1779 species. Zoologica Scripta 2013; 42: 612-622.

1780 Martinsson S, Erseus C. Cryptic diversity in supposedly species-poor genera of
1781 Enchytraeidae (Annelida: Clitellata). Zoological Journal of the Linnean Society 2018;
1782 183: 749-762.

1783 Matzen da Silva J, Creer S, Dos Santos A, Costa A, Cunha M, Costa F, et al. Systematic and
1784 Evolutionary Insights Derived from mtDNA COI Barcode Diversity in the Decapoda
1785 (Crustacea: Malacostraca). Plos One 2011a; 6: e19449.

1786 Matzen da Silva J, Dos Santos A, Cunha M, Costa F, Creer S, Carvalho G. Multigene
1787 Molecular Systematics Confirm Species Status of Morphologically Convergent
1788 *Pagurus* Hermit Crabs. Plos One 2011b; 6: e28233.

1789 Matzen da Silva J, Dos Santos A, Cunha M, Costa F, Creer S, Carvalho G. Investigating the
1790 molecular systematic relationships amongst selected Plesionika (Decapoda:
1791 Pandalidae) from the Northeast Atlantic and Mediterranean Sea. Marine Ecology
1792 2013; 34: 157-170.

1793 Meissner K, Björklöf K, Jaale M, Könönen K, Rissanen J, Leivuori M. Proficiency Test SYKE
1794 7/2011. Boreal lake littoral and NE Baltic benthic macroinvertebrate taxonomic

1795 identification. Reports of Finnish Environment Institute. 16. Finnish Environment
1796 Institute, Helsinki, 2012, pp. 14.

1797 Meissner K, Nygård H, Björklöf K, Jaale M, Hasari M, Laitila L, et al. Proficiency Test
1798 04/2016. Taxonomic identification of boreal freshwater lotic, lentic, profundal and
1799 North-Eastern Baltic benthic macroinvertebrates. Reports of the Finnish Environment
1800 Institute. 2. Finnish Environment Institute, Helsinki, 2017, pp. 20.

1801 Metcalfe JL. Biological Water-Quality Assessment of Running Waters Based on
1802 Macroinvertebrate Communities - History and Present Status in Europe.
1803 Environmental Pollution 1989; 60: 101-139.

1804 Mioduchowska M, Czyż MJ, Gołdyn B, Kur J, Sell J. Instances of erroneous DNA barcoding
1805 of metazoan invertebrates: Are universal cox1 gene primers too “universal”? Plos
1806 One 2018; 13: e0199609.

1807 Miya M, Sato Y, Fukunaga T, Sado T, Poulsen JY, Sato K, et al. MiFish, a set of universal
1808 PCR primers for metabarcoding environmental DNA from fishes: detection of more
1809 than 230 subtropical marine species. Royal Society Open Science 2015; 2.

1810 Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B. How Many Species Are There on Earth
1811 and in the Ocean? PLOS Biology 2011; 9: e1001127.

1812 Morinière J, Hendrich L, Balke M, Beermann AJ, König T, Hess M, et al. A DNA barcode
1813 library for Germany's mayflies, stoneflies and caddisflies (Ephemeroptera, Plecoptera
1814 and Trichoptera). Molecular Ecology Resources 2017; 17: 1293-1307.

1815 Nowak P, Schubert H, Schaible R. Molecular evaluation of the validity of the morphological
1816 characters of three Swedish Chara sections: *Chara*, *Grovesia*, and *Desvauxia*
1817 (Charales, Charophyceae). Aquatic Botany 2016; 134: 113-119.

1818 Nygren A. Cryptic polychaete diversity: a review. Zoologica Scripta 2014; 43: 172-183.

1819 Nygren A, Parapar J, Pons J, Meissner K, Bakken T, Kongsrød JA, et al. A mega-cryptic
1820 species complex hidden among one of the most common annelids in the North East
1821 Atlantic. Plos One 2018; 13.

1822 Oliveira LM, Knebelsberger T, Landi M, Soares P, Raupach MJ, Costa FO. Assembling and
1823 auditing a comprehensive DNA barcode reference library for European marine fishes.
1824 Journal of Fish Biology 2016; 89: 2741-2754.

1825 Pall K, Mayerhofer V. Guidance on the monitoring of the biological quality elements Part 4 —
1826 Macrophytes. Systema Bio- und Management Consulting GmbH, Vienna, 2015, pp.
1827 68.

1828 Pešić V, Asadi M, Cimpean M, Dabert M, Esen Y, Gerecke R, et al. Six species in one:
1829 evidence of cryptic speciation in the *Hygrobates fluviatilis* complex (Acariformes,
1830 Hydrachnidia, Hygrobatidae). Systematic and Applied Acarology 2017; 22: 1327-
1831 1377.

1832 Petrin Z, Bækkelie KAE, Bongard T, Bremnes T, Eriksen TE, Kjærstad G, et al. Innsamling
1833 og bearbeiding av bunndyrprøver – hva vi kan enes om. NINA Report. 1276. NINA,
1834 Trondheim, 2016, pp. 41.

1835 Pfenniger M, Staubach S, Albrecht C, Streit B, Schwenk K. Ecological and morphological
1836 differentiation among cryptic evolutionary lineages in freshwater limpets of the
1837 nominal form-group *Ancylus fluviatilis* (O.F. Müller, 1774). Molecular Ecology 2003;
1838 12: 2731-2745.

1839 Phillips AJ, Siddall ME. Poly-paraphyly of Hirudinidae: many lineages of medicinal leeches.
1840 Bmc Evolutionary Biology 2009; 9: 246.

1841 Porter TM, Hajibabaei M. Over 2.5 million COI sequences in GenBank and growing. Plos
1842 One 2018; 13: e0200177.

1843 Prie V, Puillandre N, Bouchet P. Bad taxonomy can kill: molecular reevaluation of *Unio*
1844 *mancus* Lamarck, 1819 (Bivalvia: Unionidae) and its accepted subspecies.
1845 Knowledge and Management of Aquatic Ecosystems 2012: 08.

1846 Prosser SWJ, deWaard JR, Miller SE, Hebert PDN. DNA barcodes from century-old type
1847 specimens using next-generation sequencing. Molecular Ecology Resources 2016;
1848 16: 487-497.

1849 Ratnasingham S, Hebert PDN. BOLD: The Barcode of Life Data System
1850 (www.barcodinglife.org). Molecular Ecology Notes 2007; 7: 355-364.

1851 Raupach MJ, Barco A, Steinke D, Beermann J, Laakmann S, Mohrbeck I, et al. The
1852 Application of DNA Barcodes for the Identification of Marine Crustaceans from the
1853 North Sea and Adjacent Regions. Plos One 2015; 10: e0139421.

1854 Reid AJ, Carlson AK, Creed IF, Eliason EJ, Gell PA, Johnson PTJ, et al. Emerging threats
1855 and persistent conservation challenges for freshwater biodiversity. Biological Reviews
1856 2018.

1857 Rimet F. Recent views on river pollution and diatoms. Hydrobiologia 2012; 683: 1-24.

1858 Rimet F, Abarca N, Bouchez A, Kusber WH, Jahn R, Kahlert M, et al. The potential of High-
1859 Throughput Sequencing (HTS) of natural samples as a source of primary taxonomic
1860 information for reference libraries of diatom barcodes. Fottea 2018a; 18: 37-54.

1861 Rimet F, Chaumeil P, Keck F, Kermarrec L, Vasselon V, Kahlert M, et al. R-Syst::diatom: an
1862 open-access and curated barcode database for diatoms and freshwater monitoring.
1863 Database (Oxford) 2016; 2016.

1864 Rimet F, Vasselon V, A.-Keszte B, Bouchez A. Do we similarly assess diversity with
1865 microscopy and high-throughput sequencing? Case of microalgae in lakes.
1866 Organisms Diversity & Evolution 2018b; 18: 51-62.

1867 Rivera SF, Vasselon V, Ballorain K, Carpentier A, Wetzel CE, Ector L, et al. DNA
1868 metabarcoding and microscopic analyses of sea turtles biofilms: Complementary to
1869 understand turtle behavior. Plos One 2018a; 13: e0195770.

1870 Rivera SF, Vasselon V, Jacquet S, Bouchez A, Ariztegui D, Rimet F. Metabarcoding of lake
1871 benthic diatoms: from structure assemblages to ecological assessment.
1872 Hydrobiologia 2018b; 807: 37-51.

1873 Rouillard J, Lago M, Abhold K, Roschel L, Kafyeke T, Mattheiss V, et al. Protecting aquatic
1874 biodiversity in Europe: How much do EU environmental policies support ecosystem-
1875 based management? Ambio 2018; 47: 15-24.

1876 Rudolph K, Coleman CO, Mamos T, Grabowski M. Description and post-glacial demography
1877 of *Gammarus jazdzewskii* sp nov (Crustacea: Amphipoda) from Central Europe.
1878 Systematics and Biodiversity 2018; 16: 587-603.

1879 Schneider SC, Rodrigues A, Moe TF, Ballot A. DNA Barcoding the Genus Chara: Molecular
1880 Evidence Recovers Fewer Taxa Than the Classical Morphological Approach. Journal
1881 of Phycology 2015; 51: 367-380.

1882 Siddall Mark E, Trontelj P, Utevsky Serge Y, Nkamany M, Macdonald Kenneth S. Diverse
1883 molecular data demonstrate that commercially available medicinal leeches are not
1884 *Hirudo medicinalis*. Proceedings of the Royal Society B: Biological Sciences 2007;
1885 274: 1481-1487.

1886 Stevenson J. Ecological assessments with algae: a review and synthesis. Journal of
1887 Phycology 2014; 50: 437-61.

1888 Stur E. DNA barcoding of Norwegian water mites. 2017. NTNU University Museum,
1889 www.norbol.org, 2017.

1890 Trebitz AS, Hoffman JC, Grant GW, Billehus TM, Pilgrim EM. Potential for DNA-based
1891 identification of Great Lakes fauna: match and mismatch between taxa inventories
1892 and DNA barcode libraries. Scientific Reports 2015; 5: 12162.

1893 UN World Water Assessment Programme. The United Nations World Water Development
1894 Report 2018: Nature-Based Solutions for Water. UNESCO, Paris, 2018, pp. 139.

1895 Ushio M, Murakami H, Masuda R, Sado T, Miya M, Sakurai S, et al. Quantitative monitoring
1896 of multispecies fish environmental DNA using high-throughput sequencing.
1897 Metabarcoding and Metagenomics 2018; 2: e23297.

1898 Valentini A, Taberlet P, Miaud C, Civade R, Herder J, Thomsen PF, et al. Next-generation
1899 monitoring of aquatic biodiversity using environmental DNA metabarcoding. Molecular
1900 Ecology 2015; 25: 929-942.

1901 Vasselon V, Bouchez A, Rimet F, Jacquet S, Trobajo R, Corniquel M, et al. Avoiding
1902 quantification bias in metabarcoding: Application of a cell biovolume correction factor

1903 in diatom molecular biomonitoring. *Methods in Ecology and Evolution* 2018; 9: 1060-
1904 1069.

1905 Vasselon V, Rimet F, Tapolczai K, Bouchez A. Assessing ecological status with diatoms
1906 DNA metabarcoding: Scaling-up on a WFD monitoring network (Mayotte island,
1907 France). *Ecological Indicators* 2017; 82: 1-12.

1908 Vonlanthen P, Bittner D, Hudson AG, Young KA, Muller R, Lundsgaard-Hansen B, et al.
1909 Eutrophication causes speciation reversal in whitefish adaptive radiations. *Nature*
1910 2012; 482: 357-U1500.

1911 Walters C, Hanner R. Platforms for DNA banking. In: De Vicente MC, Andersson MS,
1912 editors. *DNA Banks - Providing Novel Options for Gene Banks?* International Plant
1913 Genetic Resources Institute, Rome, 2006, pp. 22-35.

1914 Ward RD, Costa FO, Holmes BH, Steinke D. DNA barcoding of shared fish species from the
1915 North Atlantic and Australasia: minimal divergence for most taxa, but *Zeus faber* and
1916 *Lepidopus caudatus* each probably constitute two species. *Aquatic Biology* 2008; 3:
1917 71-78.

1918 Weigand H, Weiss M, Cai HM, Li YP, Yu LL, Zhang C, et al. Deciphering the origin of mito-
1919 nuclear discordance in two sibling caddisfly species. *Molecular Ecology* 2017; 26:
1920 5705-5715.

1921 Weiss M, Weigand H, Weigand AM, Leese F. Genome-wide single-nucleotide polymorphism
1922 data reveal cryptic species within cryptic freshwater snail species-The case of the
1923 *Ancylus fluvialis* species complex. *Ecology and Evolution* 2018; 8: 1063-1072.

1924 Young HS, McCauley DJ, Galetti M, Dirzo R. Patterns, Causes, and Consequences of
1925 Anthropocene Defaunation. *Annual Review of Ecology, Evolution, and Systematics*
1926 2016; 47: 333-358.

1927 Zampoukas N, Palialexis A, Duffek A, Graveland J, Giorgi G, Hagebro C, et al. Technical
1928 guidance on monitoring for the Marine Strategy Framework Directive. EUR - Scientific
1929 and Technical Research Reports, Luxembourg, 2014.

1930 Zimmermann J, Abarca N, Enk N, Skibbe O, Kusber WH, Jahn R. Taxonomic Reference
1931 Libraries for Environmental Barcoding: A Best Practice Example from Diatom
1932 Research. *Plos One* 2014; 9.

1933 Zimmermann J, Glockner G, Jahn R, Enke N, Gemeinholzer B. Metabarcoding vs.
1934 morphological identification to assess diatom diversity in environmental studies.
1935 *Molecular Ecology Resources* 2015; 15: 526-542.