

1 **Root microbiota assembly and adaptive differentiation among European**
2 ***Arabidopsis* populations**

3
4 Thorsten Thiergart^{1,7}, Paloma Durán^{1,7}, Thomas Ellis², Ruben Garrido-Oter^{1,3}, Eric Kemen⁴, Fabrice
5 Roux⁵, Carlos Alonso-Blanco⁶, Jon Ågren^{2,*}, Paul Schulze-Lefert^{1,3,*}, Stéphane Hacquard^{1,*}.

6
7 ¹Max Planck Institute for Plant Breeding Research, 50829 Cologne, Germany
8 ²Department of Ecology and Genetics, Evolutionary Biology Centre, Uppsala University, SE-752 36
9 Uppsala, Sweden
10 ³Cluster of Excellence on Plant Sciences (CEPLAS), Max Planck Institute for Plant Breeding Research,
11 50829 Cologne, Germany

12 ⁴Department of Microbial Interactions, IMIT/ZMBP, University of Tübingen, 72076 Tübingen,
13 Germany

14 ⁵LIPM, INRA, CNRS, Université de Toulouse, 31326 Castanet-Tolosan, France
15 ⁶Departamento de Genética Molecular de Plantas, Centro Nacional de Biotecnología (CNB), Consejo
16 Superior de Investigaciones Científicas (CSIC), 28049 Madrid, Spain

17 ⁷These authors contributed equally: Thorsten Thiergart, Paloma Durán

18 *e-mail: jon.agren@ebc.uu.se, schlef@mpipz.mpg.de, hacquard@mpipz.mpg.de

19
20 **Summary**
21 Factors that drive continental-scale variation in root microbiota and plant adaptation are poorly
22 understood. We monitored root-associated microbial communities in *Arabidopsis thaliana* and co-
23 occurring grasses at 17 European sites across three years. Analysis of 5,625 microbial community
24 profiles demonstrated strong geographic structuring of the soil biome, but not of the root microbiota.
25 Remarkable similarity in bacterial community composition in roots of *A. thaliana* and grasses was
26 explained by the presence of a few diverse and geographically widespread taxa that disproportionately
27 colonize roots across sites. In a reciprocal transplant between two *A. thaliana* populations in Sweden
28 and Italy, we uncoupled soil from location effects and tested their respective contributions to root

29 microbiota variation and plant adaptation. The composition of the root microbiota was affected by
30 location and soil origin, and to a lesser degree by host genotype. The filamentous eukaryotes were
31 particularly strongly affected by location. Strong local adaptation between the two *A. thaliana*
32 populations was observed, with difference in soil properties and microbes of little importance for the
33 observed magnitude of adaptive differentiation. Our results suggest that, across large spatial scales,
34 climate is more important than are soil conditions for plant adaptation and variation in root-associated
35 filamentous eukaryotic communities.

36

37 **Introduction**

38 Plants interact with multi-kingdom microbial communities (e.g. bacteria, fungi, oomycetes) that can
39 impact host fitness, either directly, or indirectly through microbe-microbe interactions^{1, 2, 3, 4, 5}. The
40 immune system of plants, rhizodeposits, and microbial interactions are known determinants of root-
41 associated microbial assemblages and make them distinct from the surrounding soil biota^{6, 7, 8, 9}. Large-
42 scale spatial variation in the composition of the soil biota has been associated with difference in edaphic
43 and climatic conditions⁹. Particularly, local edaphic factors such as soil pH primarily predict geographic
44 distribution of soil bacteria^{10, 11, 12}, whereas climatic variables better predict fungal distribution in soil¹³.
45 However, systematic field studies exploring and disentangling the extent to which variation in soil and
46 climatic conditions impacts root microbiota composition and adaptive differentiation in plants are
47 lacking.

48

49 Local adaptation has been documented in a large number of plant species and across both small and
50 large spatial scales¹⁴. For example, reciprocal transplants and common-garden experiments have
51 provided evidence of strong adaptive differentiation among natural populations of the model plant
52 *Arabidopsis thaliana*^{15, 16}. However, the relative importance of different abiotic and biotic factors for the
53 evolution and maintenance of local adaptation is poorly known^{17, 18}. Particularly, soil edaphic factors
54 and soil microbes are known to influence flowering phenology and modulate host fitness in natural
55 soils^{19, 20, 21, 22, 23}, even at the scale of a few meters²⁴. Yet, information about the extent to which
56 differences in soil properties contribute to divergent selection and the maintenance of adaptive

57 differentiation among plant populations is still limited beyond classical examples of adaptation to
58 extreme soil conditions²⁵.

59
60 Here, we tested whether roots of *A. thaliana* and co-occurring grasses growing in various soils and
61 climatic environments establish stable associations with bacterial and filamentous eukaryotic
62 communities across a latitudinal gradient in Europe. In a reciprocal transplant between two *A. thaliana*
63 populations in Sweden and Italy, we uncoupled soil from location effects and experimentally tested the
64 hypothesis that soil properties and climate drive root microbiota assembly and adaptive differentiation
65 between the two *A. thaliana* populations.

66
67 We found that a widespread set of bacteria, but not filamentous eukaryotes, establish stable associations
68 with roots of *A. thaliana* and grasses across 17 sites in Europe, despite strong geographical structuring
69 and variation in the surrounding soil communities. The reciprocal transplant of soil and plant genotypes
70 between two native *A. thaliana* populations in northern and southern Europe showed that the
71 composition of root microbiota was more affected by soil properties and location than by host genotype.
72 The effect of soil was stronger than that of location for root-associated bacteria, whereas the effect of
73 location was stronger for root-associated fungi and oomycetes. Transplant location, rather than origin
74 of soil, also largely accounted for strong selection against the nonlocal *A. thaliana* genotype at each site.
75 Our results suggest that climate is a primary force driving geographic variation in filamentous eukaryotic
76 communities in roots and adaptive differentiation between *A. thaliana* populations in northern and
77 southern Europe.

78

79 **Results**

80
81 **Continental-scale survey of the *A. thaliana* root microbiota.** We sampled natural *A. thaliana*
82 populations at the flowering stage at 17 sites along a latitudinal gradient in Europe in three consecutive
83 years (2015, 2016 and 2017). We harvested bulk soil (soil), rhizosphere (RS), rhizoplane (RP), and root
84 endosphere (root) compartments of *A. thaliana* and co-occurring grasses (**Supplementary Fig. 1a, b**)

85 at four sites in Sweden (SW1-4), six in Germany (GE1-6), three in France (FR1-3), one in Italy (IT1),
86 and three in Spain (SP1-3) (**Fig. 1a**), each having distinct environmental and soil characteristics
87 (**Supplementary Table 1**). DNA was isolated and microbial community composition analyzed for a
88 total of 1,125 samples. Bacterial, fungal and oomycetal communities were profiled using primer pairs
89 targeting the V2V4 and V5V7 regions of the bacterial 16S rRNA gene, the ITS1 and ITS2 segments of
90 the fungal ITS, and the ITS1 segment of the oomycetal ITS, resulting in the sequencing of 5,625
91 microbial community profiles (**Supplementary Table 2**). Given the correlation observed at the class
92 level between the independently used primer pairs that target the two regions of the bacterial 16S rRNA
93 gene (Pearson correlation, $r = 0.40$, $p < 0.001$) and the two regions of the fungal ITS (Pearson correlation,
94 $r = 0.93$, $p < 0.001$) (**Supplementary Fig. 1c**), only the bacterial V5V7 and fungal ITS1 variable
95 segments were considered for later analyses.

96

97 **Convergence in root microbiota composition across European *A. thaliana* populations.** If plant
98 roots establish stable associations with microbial communities across large geographical distances, we
99 expect a strong host filtering effect on root microbiota composition. Inspection of alpha-diversity indices
100 (i.e. Shannon index, number of observed OTUs) revealed a gradual decrease of bacterial, fungal, and
101 oomycetal diversity from the soil to the root endosphere (Kruskal-Wallis test, $p < 0.05$), with
102 significantly stronger decrease for root-associated bacteria than for filamentous eukaryotes (**Fig. 1b** and
103 **Supplementary Fig. 2a, b**). Analysis of microbial community structure based on Bray-Curtis distances
104 across sites and compartments revealed that bacterial communities in the root endosphere and RP cluster
105 by compartments, and by site in the RS and soil (**Fig. 1b**). A clustering by compartment was also visible
106 among root endosphere samples for fungi, but not for oomycetes (**Fig. 1b**). By considering soil and root
107 endosphere samples, compartment explained variation in bacterial community composition more than
108 did site (Compartment: 18.9%; site: 17.2%, PERMANOVA with Bray-Curtis distances, $p < 0.01$;
109 **Supplementary Table 3**). In contrast, site explained variation in fungal and oomycetal community
110 composition more than did compartment (Site: 20.2% for fungi, 15.5% for oomycetes; Compartment:
111 6.5% for fungi, 2.6% for oomycetes; PERMANOVA with Bray-Curtis distances, $p < 0.01$,
112 **Supplementary Table 3**). Nonetheless, the observation that endosphere-associated bacterial and fungal

113 communities show overall more similarities across sites than across compartments at a given site implies
114 structural convergence of the *A. thaliana* root microbiota at a continental scale. This is well illustrated
115 by the marked differences in soil biota observed between Swedish soils (SW1-4) and the other European
116 soils (**Fig. 1c** and **d**, **Supplementary Fig. 2c**), which is largely diminished in the corresponding root
117 endosphere samples (**Fig. 1d**). Compared to surrounding soil samples, microbial communities in plant
118 roots showed a significant enrichment of taxa belonging to the bacterial classes Beta- and Gamma-
119 Proteobacteria, the fungal classes Leotiomycetes and Dothideomycetes and the oomycetal order
120 Pythiales (Wilcoxon rank sum test, FDR < 0.05, **Fig. 1c** and **Supplementary Fig. 2d**). Our results
121 demonstrate that the root environment drives remarkable convergence in bacterial, and to a lesser extent
122 fungal and oomycetal community composition across European sites separated by up to 3,500 km,
123 despite considerable differences in soil properties (**Supplementary Table 1**) and soil microbial
124 communities.

125
126 **Root endosphere bacteria and *A. thaliana* exhibit stable associations across Europe.** Given the
127 limited variation in microbial community composition observed in plant roots across geographically
128 distant sites, we hypothesized that the presence of geographically widespread microbes might contribute
129 to convergence in root microbiota composition. To identify microbial OTUs that are widely distributed
130 in roots of *A. thaliana* across Europe, we calculated their prevalence across all 17 sites and those detected
131 in more than 80% of the sites were defined as geographically widespread (**Fig. 2a** and **Supplementary**
132 **Fig. 3a**). Remarkably, we observed a positive correlation (linear regression, $R^2 = 0.24$, $p < 0.001$)
133 between the prevalence of root-associated bacteria across sites and their relative abundance (RA) in root
134 endosphere samples, suggesting that bacterial taxa that colonize *A. thaliana* roots across Europe are also
135 the most abundant in this niche (**Fig. 2a**). This implies that a small subset of bacterial taxa have evolved
136 mechanisms to dominate the bacterial root microbiota at a continental scale, irrespective of major
137 differences in the surrounding local bacterial soil biota. In contrast, no significant correlation was
138 observed between relative abundance in plant roots and prevalence across sites for root-associated fungi
139 and oomycetes (linear regression, $p > 0.05$, **Fig. 2a**). By inspecting the abundance profiles of OTUs with
140 restricted or widespread geographic distribution across compartments (**Fig. 2b**), we observed that

141 geographically widespread bacterial OTUs are significantly more abundant in RP and Root samples than
142 in the corresponding soil samples (Kruskal-Wallis test, FDR < 0.01). In contrast, fungal OTUs that have
143 a narrow geographical distribution are significantly more abundant in RP and Root samples than in the
144 corresponding bulk soil (Kruskal-Wallis test, FDR < 0.01).

145
146 We identified 13 geographically widespread bacterial OTUs that are consistently detected in the root
147 endosphere of *A. thaliana* across European sites (Fig. 2a, c and Supplementary Table 4), accounting
148 for 38% of the total relative abundance in this niche (Fig. 2d). These taxa belong to 5 different classes,
149 and cover 10 bacterial genera, including *Bradyrhizobium*, *Pseudomonas*, *Polaromonas*, *Acidovorax*,
150 *Ralstonia*, *Massilia*, *Burkholderia*, *Kineospora* and *Flavobacterium* (Fig. 2c), which indicates
151 convergent adaptation to the root environment in phylogenetically distant bacterial lineages at a
152 continental scale. Notably, these 13 bacterial OTUs were detected across all three years (Fig. 2a), and
153 nine of them were significantly enriched in plant roots compared to soil (FDR < 0.05, highlighted with
154 a star in Fig. 2c). In contrast, the abundance of the 14 geographically widespread OTUs of root-
155 associated filamentous eukaryotes varied among years, and were dominated by fungi from only three
156 classes (Sordariomycetes, Leotiomycetes and Dothideomycetes) and oomycetes from a single genus
157 (*Pythium*), and were not particularly root-enriched (Fig. 2a, c, Supplementary Table 4). Our results
158 suggest that the few geographically widespread bacteria that abundantly colonize roots of *A. thaliana*
159 drive convergence in bacterial community composition at a continental scale.

160
161 **Geographically widespread bacterial OTUs in *A. thaliana* roots are ubiquitous in distantly related**
162 **plant species.** The conserved taxa that consistently associate with roots of *A. thaliana* might represent
163 a widespread microbial multi-kingdom community that also associate with distantly related plant
164 species. To test this hypothesis, we harvested co-occurring grasses at each of the 17 sites across Europe
165 and compared microbial community composition between roots of *A. thaliana* and grasses. The factor
166 host species weakly, but significantly explained bacterial, fungal, and oomycetal community
167 composition in root endosphere samples when considering the whole dataset (1.3%, 0.89% 1.8% of the
168 variance, respectively, PERMANOVA with Bray–Curtis distances, p < 0.01, Supplementary Table 5).

169 The overall effect of host species is likely underestimated because we could not sample the same grass
170 species at all sites and we pooled plant individuals for efficient RP and root fractionation (see **Methods**).
171 By inspecting the species effect at each site separately using PERMANOVA, we observed a stronger
172 effect of the host species on the root microbiota, although these differences were significant for only
173 few sites (average explained variance: 11% for bacteria, 8.8% for fungi, 7.7% for oomycetes,
174 **Supplementary Table 6**). Comparison of microbial OTU prevalence in roots of *A. thaliana* with those
175 of co-occurring grasses revealed overall consistency in root OTU prevalence at a continental scale
176 (Spearman rank correlation, $r_s = 0.69$ for bacteria; $r_s = 0.79$ for fungi; $r_s = 0.72$ for oomycetes; $p < 0.01$)
177 (**Supplementary Fig. 3b**). This indicates an overall conserved distribution of geographically restricted
178 and widespread OTUs in roots of phylogenetically distant plants species that evolved independently in
179 the Brassicaceae and Poaceae lineages. Inspection of the 13 geographically widespread bacterial OTUs
180 detected in roots of *A. thaliana* revealed that these are also abundantly detected in roots of co-occurring
181 grasses (13/13 detected, 36% of the total relative abundance), whereas conservation was less obvious
182 for the geographically widespread fungal OTUs (5/7 detected, 15% of the total relative abundance).
183 Further inspection of their abundance in roots of *Lotus japonicus* (Fabaceae) grown in a completely
184 different soil type (i.e. Cologne Agricultural Soil²⁶) validated the ubiquitous nature of the bacterial OTUs
185 (11/13 detected, 16% of the total RA in roots), but not of the fungal OTUs (2/8 detected, 3% of the total
186 RA in roots; **Fig. 2d**). The results suggest that a small number of geographically widespread bacteria
187 can efficiently colonize roots of distantly related plant species and establish potentially stable
188 associations with plant roots over evolutionary time.

189
190 **Spatial and temporal variation in root microbiota differentiation.** Despite limited among-site
191 variation in microbial communities in root endosphere samples across European sites, we did observe
192 spatial and temporal variation in their composition (**Fig. 3a, Supplementary Fig. 4a, b**). Site explained
193 19.4%, 20.1% and 17.7% of the variance in bacterial, fungal, and oomycetal community composition in
194 roots of European *A. thaliana* populations respectively, compared with 54.5%, 51.3%, and 28.6% of the
195 variance in corresponding soil samples (PERMANOVA with Bray–Curtis distances, $p < 0.01$, **Fig. 3b**
196 and **Supplementary Table 5**). Inspection of correlations between environmental or soil variables and

197 microbial community composition revealed a gradual decrease in explanatory power from soil towards
198 root compartments, as well as stronger correlations for bacterial communities than for filamentous
199 eukaryotes (**Fig. 3b**, **Supplementary Fig. 5**, and **Supplementary Tables 1 and 5**). Although part of the
200 observed correlations can be due to confounding effects between variables (**Supplementary Fig. 5a, b**),
201 latitude and soil pH explained the highest proportions of variation in bacterial and fungal community
202 composition in soil (bacteria: 21.2% and 22.8%, respectively; fungi: 10.2% and 10.5%, respectively;
203 PERMANOVA, $p < 0.01$), and remained among the most significant variables explaining among-site
204 variation in the composition of the root microbiota across European *A. thaliana* populations (bacteria:
205 3.2% and 3.0%, respectively; fungi: 3.0% and 2.7%, respectively; PERMANOVA, $p < 0.01$) (**Fig. 3b**,
206 **Supplementary Fig. 5** and **Supplementary Table 5**). To assess among-year variation in root
207 microbiota composition, we sampled *A. thaliana* and grass populations at the same phenological stage
208 in three successive years in spring 2015, 2016, and 2017. Remarkably, “year” explained more variation
209 in root-associated microbial communities (bacteria: 6.9%, fungi: 13.2%, oomycetes: 15.8%,
210 PERMANOVA, $p < 0.01$), than did “host species” (< 2%, PERMANOVA, $p < 0.01$), suggesting that
211 year-to-year environmental variation affected the establishment of the root microbiota more than did
212 differences between hosts separated by > 140 million years of reproductive isolation²⁷ (**Fig. 3b** and
213 **Supplementary Fig. 4b** and **Supplementary Table 5**). Among-year variation was particularly strong
214 for soil- and root-associated fungal (soil: 4.0 %, root: 13.2%, PERMANOVA, $p < 0.01$) and oomycetal
215 (soil: 15.9%, root: 15.8%, PERMANOVA, $p < 0.01$) communities (**Fig. 3a, b**, and **Supplementary Fig.**
216 **4b** and **Supplementary Table 5**).

217
218 Consistent with earlier reports, a significant proportion of the variance remained unexplained in
219 bacterial, fungal, and oomycetal communities in both soil and root compartments (> 40%, data not
220 shown), likely arising from unmeasured environmental variables, stochastic processes, or species
221 interactions such as microbe-microbe interactions. To identify signatures of microbial interactions, we
222 quantified correlations between bacterial and fungal alpha diversity across sites. The correlation between
223 bacterial and fungal diversity was positive in soil samples (Spearman’s rank correlation, observed
224 OTUs: $r_s = 0.30$, $p < 0.001$, Shannon index: $r_s = 0.21$, $p = 0.002$), but negative in root endosphere samples

225 (Observed OTUs: $r_s = -0.15$, $p = 0.092$, Shannon index: $r_s = -0.24$, $p = 0.005$) (**Supplementary Fig. 6a, b**). The observation that bacterial diversity is negatively correlated with fungal diversity in the root 226 endosphere suggests that microbial interactions also contribute to community differentiation at the soil- 227 root interface, as recently reported³. Overall, our results suggest that among-site variation in 228 environmental conditions affected root-associated microbial communities more strongly than did 229 temporal variation within sites. In contrast, differences between host species had less impact on root 230 microbiota differentiation than had spatial and temporal variations. Therefore, differences in soil 231 properties (e.g., pH, which ranged from 5.1 to 7.9) and climate are likely causes of the variation in root 232 microbiota among European populations of *A. thaliana*.

234

235 **Site-specific differences in soil, climate, and genotype drive root microbiota differentiation**
236 **between two *A. thaliana* populations.** We hypothesized that continental-scale variation in the *A.*
237 *thaliana* root microbiota is mediated by interactions between soil, climate, and host genotype and that
238 these factors might differentially influence the establishment of bacteria and filamentous eukaryotes in
239 plant roots. To disentangle the respective contribution of these three factors on microbial community
240 composition at the root interface, we conducted a transplant experiment between two geographically
241 widely separated *A. thaliana* populations in Sweden and Italy (SW4 and IT1) (**Supplementary Fig. 7a,**
242 **b**). In this experiment, we transplanted seedlings of each plant genotype into soils from both SW4 and
243 IT1 at each site in autumn, at the time of natural seedling establishment in the local populations (**Fig.**
244 **4a**). At fruit maturation, the two *A. thaliana* genotypes were harvested and community composition was
245 defined for bacteria, fungi and oomycetes in soil (Soil with RS-associated microbes) and whole root
246 (Root- with RP-associated microbes) samples ($n = 131$, **Fig. 4b**).

247

248 Consistent with the European transect experiment, we observed a greater compartment effect for bacteria
249 than for filamentous eukaryotes, explaining 17.8%, 7.4%, and 3.1% of the variance in bacterial, fungal,
250 and oomycetal community composition, respectively (PERMANOVA, $p < 0.001$, **Fig. 4b** and
251 **Supplementary Table 7**). However, the degree of structural convergence in roots was weaker, probably
252 due to the fact that we did not partition the RP compartment from the root endosphere in this experiment.

253 Because no Italian plants survived at the Swedish site, we used a canonical analysis of principal
254 coordinates (CAP; capscale function) to assess the effect of genotype among whole root samples at the
255 Italian site only. CAP analysis constrained by genotype indicated a significant effect of host genotype
256 for all three microbial groups in plant roots, explaining 2.4% ($p = 0.001$), 2.36% ($p = 0.026$) and 2.9%
257 ($p = 0.002$) of the variance in bacterial, fungal, and oomycetal community composition, respectively
258 (**Supplementary Fig. 7d**).

259
260 The origin of soil (Italy *vs.* Sweden) explained substantially more variation in bacterial soil biota than
261 did transplant location (soil origin: 62.3%, transplant location: 15.8%; PERMANOVA, $p < 0.001$). This
262 was also the case for bacterial communities associated with whole root samples (soil origin: 47.1%,
263 transplant location: 17.4%; PERMANOVA, $p < 0.05$) (**Fig. 4b** lower panel, **Supplementary Table 7**).
264 The percentage of variation explained by the origin of soil was weaker for fungi and oomycetes in the
265 soil fraction (fungi: 29.6%, oomycetes: 18.54%; PERMANOVA, $p < 0.001$), whereas differences other
266 than soil origin, including climatic differences between sites, also largely accounted for between-site
267 variation in microbial eukaryotic assemblages in soil (fungi: 27.5%, oomycetes: 6.3%; PERMANOVA,
268 $p < 0.001$). In whole root samples, the effect of the location was as strong as the effect of the soil for
269 fungi (location: 14.6%, soil origin: 15.1%; PERMANOVA, $p < 0.001$) and stronger than the effect of
270 the soil for oomycetes (location: 10.6%, soil origin: 4.7%; PERMANOVA, $p < 0.001$) (**Fig 4b** lower
271 panel, **Supplementary Table 7**). These results indicated that soil origin and transplant location
272 differentially affect the assembly of root-associated bacteria and filamentous eukaryotes.

273
274 Inspection of root-associated microbial OTUs that responded to differences in soil origin and/or location
275 in the reciprocal transplant demonstrated the stable prevalence of geographically widespread bacterial
276 OTUs in root samples representing different soil \times location \times genotype combinations (**Supplementary**
277 **Fig. 8a, b and c**). Abundance profiles of microbial taxa in whole root samples were impacted by soil
278 origin, location, and their interaction, with notable differences depending on microbial classes. For
279 example, variation in root-associated Actinobacteria was almost exclusively explained by soil origin,
280 whereas variation in root-associated Alpha-, Beta-, and Gamma-proteobacteria was explained by soil

281 origin and location to similar degrees (**Supplementary Fig. 8d**). Similarly, the presence of root-
282 associated fungi belonging to Dothideomycetes was primarily explained by location, whereas the
283 presence of root-associated fungi belonging to Leotiomycetes was largely explained by soil origin
284 (**Supplementary Fig. 8d**). The results point to differential impact of site and soil specific factors on
285 different taxonomic groups of the root microbiota. Overall, 74.4% and 86.3% of root-associated fungal
286 and oomycetal OTUs, respectively, responded to difference in location or location x soil, whereas this
287 percentage was reduced to 44.7% for root-associated bacterial OTUs (**Supplementary Fig. 8d**). Taken
288 together, our data suggest that location-specific factors such as climatic conditions affect differentiation
289 of root-associated filamentous eukaryotic communities more than that of bacteria, and this likely
290 explains variation in community structure observed for these microbes among European sites and across
291 successive years (**Fig. 3** and **Supplementary Fig. 4b**).

292
293 **Site-specific differences in soil conditions only weakly contributes to adaptive differentiation**
294 **between *A. thaliana* populations.** If adaptation to the biotic or abiotic characteristics of the local soil
295 contributes to adaptive differentiation between *A. thaliana* populations, the advantage of the local over
296 nonlocal genotype should be greater when plants are grown on local soil than when grown on non-local
297 soils. To test this hypothesis, we scored plant survival and fecundity (number of fruits per reproducing
298 plant) and estimated overall fitness (number of fruits per seedling planted) of the plants grown in the
299 reciprocal transplant experiments at the sites of IT1 and SW4 (n = 1,008; **Supplementary Table 8**).

300
301 At the IT1 site the relative fitness of the local genotype was higher on local than on non-local soil: the
302 mean fitness of the Italian genotype was 58 times higher than that of the non-local Swedish genotype
303 on Italian soil (corresponding to a selection coefficient of $s = 0.98$), and 23 times higher ($s = 0.96$) on
304 Swedish soil (significant soil x genotype interaction, $F_{1,10} = 5.7$, $p = 0.038$; **Fig. 4c, Supplementary**
305 **Table 9**). The strong local advantage was a function of both higher survival and fecundity
306 (**Supplementary Fig. 7c**). At the SW4 site, no Italian plant survived to reproduce, meaning that
307 selection coefficients were 1 on both soil types (**Fig. 4c**). It was therefore not meaningful to compare
308 adaptive differentiation on the two soil types. However, the Swedish genotype showed 3.5-fold higher

309 survival and 4.0-fold higher overall fitness when planted in Swedish soil compared to in Italian soil
310 (survival: one-tailed test, $t = 1.81$, $p = 0.060$; overall fitness: $t = 2.14$, $p = 0.039$) (**Fig. 4c**,
311 **Supplementary Fig. 7c**). These results demonstrate a strong selective advantage to the local *A. thaliana*
312 genotype at both sites, consistent with previous studies on the same populations^{15, 28}. However, despite
313 the marked differences in the geochemical properties and microbial communities of the soil, adaptation
314 to local soil conditions explained only a small fraction of the adaptive differences between the two *A.*
315 *thaliana* populations. This suggests that adaptation to climatic variables not related to the characteristics
316 of the soil are the primary drivers of adaptive differentiation between *A. thaliana* populations in northern
317 and southern Europe.

318

319 **Discussion**

320 By monitoring the root microbiota of *A. thaliana* in 17 natural populations across three successive years,
321 we observed strong geographic structuring of soil microbial communities, but a surprising degree of
322 convergence among root-associated taxa. The convergence in microbial community composition
323 observed for root endosphere samples is remarkable given the large geographical distances between
324 sites, contrasting edaphic characteristics, and distinct microbial communities in the surrounding bulk
325 soils. Our results indicate that environmental factors and variables that explain geographic distribution
326 of microbes in soil have less predictive power in plant roots. The degree of convergence in community
327 structure varied among microbial groups and was more evident for the bacterial root microbiota than for
328 filamentous eukaryotes. Differences between sites that were independent of soil characteristics largely
329 explained differentiation in filamentous eukaryotic communities in our reciprocal transplant experiment,
330 suggesting a link between climatic conditions and variation observed for filamentous eukaryotic
331 communities across sites and years. Our results indicate that difference in climate and soil between sites
332 not only contribute to variation in root-associated microbial communities, but also to adaptive
333 divergence between two *A. thaliana* populations in northern and southern Europe.

334

335 The striking structural convergence observed for bacterial communities in roots of *A. thaliana* and
336 grasses was explained by the presence of a few, diverse and geographically widespread taxa that

337 disproportionately colonize roots across sites. The conserved enrichment of Beta- and Gamma-
338 proteobacteria in *A. thaliana* roots, together with the identification of 13 habitat-generalist OTUs that
339 were consistently and abundantly detected in the roots of *A. thaliana* and grass populations suggest
340 potential co-evolutionary histories between these microbes and evolutionary distant plant species. These
341 few OTUs belong to bacterial genera that are often detected in plant roots across various environmental
342 conditions^{29, 30, 31, 32, 33}. Notably, *Bradyrhizobium* and *Burkholderia* (OTUs 8, 4987, 13, 14, and 4486
343 in this study) were the two most dominant of 47 widespread genera in roots of phylogenetically diverse
344 flora across a 10 km transect in Australia³². The low number of widespread core bacterial taxa detected
345 in our study might be due to the large distances between sites and extensive differences in soil pH
346 (ranging from 5.1 to 7.9), contrasting with the more uniform range of soil pH reported in ref.³² (i.e. 4.1
347 to 4.6). Our results nonetheless suggest that at least part of the similarity in bacterial community
348 composition observed in roots of divergent plant species in microbiome studies³⁴ is driven by the
349 presence of geographically widespread taxa that efficiently colonize plant roots across a broad range of
350 environmental conditions.

351
352 The remarkable phylogenetic diversity among the few geographically widespread bacteria detected in
353 plant roots suggests convergent evolution and metabolic adaptation to the root habitat in
354 phylogenetically distant bacterial lineages³⁵. Similar to what has been described in leaves for wild *A.*
355 *thaliana*³⁶, a *Pseudomonas* OTU in our study was also by far the most dominant taxon in *A. thaliana*
356 roots (i.e. OTU5, Relative abundance in roots = 14.4% in average), pointing to *Pseudomonas* taxa as the
357 most successful and robust colonizers of wild *A. thaliana*. Although we sampled healthy-looking plants
358 in their natural habitats, we cannot exclude the possibility that some of these bacterial strains represent
359 widespread pathogens that negatively affect plant fitness without causing disease symptoms in nature³⁶.
360 Alternatively, these taxa might carry out widespread and important beneficial functions for *A. thaliana*
361 survival such as pathogen protection e.g., *Pseudomonas*, *Pelomonas*, *Acidovorax*, *Flavobacterium*^{3, 35},
362 ³⁷, abiotic stress tolerance, or plant growth promotion (e.g., *Bradyrhizobium*, *Burkholderia*^{38, 39}). It
363 remains to be seen whether this widespread association between plant roots and bacteria is evolutionary
364 ancient and the extent to which it has contributed to plant adaptation to terrestrial ecosystems.

365 Irrespective of this, the 13 core OTUs identified here in *A. thaliana* roots in natural environments provide
366 a rational framework for future design of low complexity synthetic bacterial communities from culture
367 collections of root commensals and co-culturing with gnotobiotic plants⁴⁰ to study their contribution to
368 plant health and growth in laboratory environments.

369
370 The results of the reciprocal transplant between two widely separated *A. thaliana* populations (one in
371 Italy and one in Sweden; IT1 and SW4) demonstrated significant effects of soil and location on
372 microbiota assembly in both soil and root compartments. Our observation that transplant location
373 affected the community composition of filamentous eukaryotes more strongly than that of bacteria
374 suggest that the response to climate varies among microbial kingdoms^{41, 42}. Our data corroborate the
375 hypothesis that climate is a key driver of among-site variation and geographic distribution of filamentous
376 eukaryotes in soil, as predicted based on association studies^{13, 43, 44}. This contrasts with the geographic
377 distribution of soil bacteria which is known to be primarily controlled by edaphic factors¹⁰. The marked
378 among-year variation in community composition of filamentous eukaryotes shows that year-to-year
379 differences in environmental conditions can significantly modify geographical patterns of filamentous
380 microbes at the soil-root interface across Europe. Our field transplant experiment also demonstrated an
381 effect of host genotype (Italian vs. Swedish accession) on the composition of root microbial
382 communities. The proportion of variation in community composition explained by host genotype was
383 limited (~ 2% at OTU resolution) compared to that explained by origin of soil and location of
384 experiment, but is still comparable to estimates for host genotype obtained in previous studies of *A.*
385 *thaliana* (ref.⁴⁵; variance: < 1%, p < 0.05), *Boechera stricta* (ref.⁴⁶; genotype effect not significant), and
386 *Populus* (ref.⁴⁷; variance: ~ 3%, p < 0.05). Taken together, our results suggest that bacterial and fungal
387 assemblages in roots are differentially controlled by edaphic and climatic conditions, and that host
388 genetic differences contribute only little to root microbiota differentiation between the two *A. thaliana*
389 populations.

390
391 Local adaptation is common among plant populations, but the extent to which divergent selection and
392 local adaptation can be attributed to local soil conditions has been examined mainly in relation to high

393 concentrations of heavy metals⁴⁸, serpentine⁴⁹, and high salinity^{25, 50}. The field experiment of the present
394 study, in which both soils and plant genotypes were reciprocally transplanted at two locations in Sweden
395 and Italy, demonstrated strong local adaptation between two geographically widely separated *A.*
396 *thaliana* populations. The relative performance of local and non-local host genotypes was primarily
397 affected by the geographical location and only weakly by soil origin, despite extensive differences in
398 microbial community composition and physical and chemical properties between Swedish and Italian
399 soils (Fig. 1d and Supplementary Table 1). Particularly, differences in available Ca, reserve K,
400 available Mg, pH, iron, and available K between the two soils accounted for 99.4%, 93.3%, 87.9%, 60.7,
401 39.4% and 30%, of the total variation, respectively, observed for these parameters across all 17 soils
402 (Supplementary Table 1). In their native habitats, the two populations are exposed to widely different
403 climates and also to differences in seasonal changes of day length. Previous work has demonstrated that
404 the relative survival of the Italian genotype in Sweden is negatively related to minimum soil temperature
405 in winter¹⁵, and also that genetic differences in phenological traits such as timing of germination and
406 flowering can explain a substantial portion of selection against the non-local genotype at the two sites⁵¹,
407 ^{52, 53}. Taken together, our data indicate that differences in climate have been more important than
408 differences in soil and endogenous microbiota for the adaptive divergence between the two study
409 populations. Although we did not uncouple soil-mediated from microbe-mediated local adaptation in
410 this study, our results suggest that the extensive differences in microbial community composition
411 between IT1 and SW4 soil contribute little to adaptive differentiation between two *A. thaliana*
412 populations in northern and southern Europe. Future studies should determine whether the
413 environmental factors that affect root microbiota assembly and adaptive differentiation at large spatial
414 scales act differentially at smaller geographical scales.

415

416 **Methods**

417 **Harvesting of *A. thaliana* and grasses across 17 European sites.** We selected seventeen sites with
418 variable soil characteristics across a climatic gradient from Sweden to Spain. At each site, *A. thaliana*
419 and sympatric grasses occur naturally^{54, 55, 3, 1, 15} (Supplementary Table 1, Fig. 1a). We harvested *A.*
420 *thaliana* from February to May at the same developmental stage (bolting/flowering stage), for one (three

421 sites), two (one site) and three (thirteen sites) consecutive years (**Supplementary Table 1**). Plants were
422 harvested with their surrounding bulk soil with a hand shovel without disturbing the plant root system,
423 transferred to 7x7 cm greenhouse pots and transported immediately to nearby laboratories for further
424 processing. Four plant individuals pooled together were considered as one pooled-plant technical
425 replicate (4 technical replicates in total). In addition, four plants were not pooled and kept individually
426 as single-plant replicates to assess difference in microbial community composition between single plant
427 and pooled plants. Note that these two sampling strategies explain only 0.8, 3 and 2.4% of the variation
428 in bacterial, fungal, and oomycetal community composition, respectively (**Supplementary Table 5**,
429 **Supplementary Fig. 4b**). At each of the 17 sites, we also harvested and processed similarly three
430 neighbouring plants growing within 50 cm of *A. thaliana* and belonging to the Poaceae family. In total,
431 we collected 285 plants across three years.

432

433 **Fractionation of soil and root samples.** To distinguish and separate four microbial niches across the
434 soil-root continuum, plants and respective roots were taken out from the pot. Samples from the bulk soil
435 exempt from root or plant debris were taken, snap-frozen in liquid nitrogen and stored for further
436 processing (Soil compartment). Individual plants were manually separated from the main soil body and
437 non-tightly adhered soil particles were removed by gently shaking the roots. These roots were separated
438 from the shoot and placed in a 15-mL falcon containing 10 mL of deionized sterile water. After 10
439 inversions, roots were transferred to another falcon and further processed, while leftover wash-off
440 (containing the RS fraction) was centrifuged at 4,000 g for 10 min. Supernatant was discarded and the
441 pellet was resuspended and transferred to a new 2-mL screw-cap tube. This tube was centrifuged at
442 20,000 rpm for 10 minutes, the supernatant was discarded and the pellet was snap-frozen in liquid
443 nitrogen and stored for further processing (RS compartment). After RS removal, cleaned roots were
444 placed in a 15-mL falcon with 6 mL of detergent (1xTE + 0.1% Triton® X-100) and manually shaken
445 for 2 minutes. This step was repeated for a total of three detergent washes, in between which, roots were
446 transferred to a new 15-mL falcon with new detergent. After these washes in detergent, roots were
447 transferred to a new 15-mL falcon. The remaining washes (approximately 18 mL) were transferred to a
448 20 mL syringe and filtered through a 0.22 μ M-pore membrane. The membrane was snap-frozen in liquid

449 nitrogen until further processing (RP compartment). Lastly, three-times washed roots were subjected to
450 a further surface sterilization step to further deplete leftover microbes from the root surface. Roots were
451 sterilized one minute in 70% ethanol, followed by one minute in 3% NaClO, and rinsed five times with
452 deionized sterile water. These roots were dried using sterile Whatman paper and snap-frozen in liquid
453 nitrogen until further processing (Root compartment) (**Supplementary Fig. 1a**).

454

455 In total, 1,125 samples were produced after fractionation. We validated the fractionation protocol by
456 printing processed roots on Tryptic Soy Agar (TSA) 50% before fractionation, after each detergent wash
457 and after surface sterilization. Washes produced after each fractionation step (without treatment, after
458 each detergent step and after surface sterilization) were also plated on TSA 50%. In brief, treated roots
459 were cut and placed on plates containing TSA 50% medium for 30 sec and then removed. After 3 days
460 at 25 °C, colony forming units (CFU) were counted. Similarly, 20 µL of remaining washes were spread
461 onto TSA 50% medium-containing plates and CFU were counted after 3 days of incubation at 25 °C
462 (**Supplementary Fig. 1b**).

463

464 **Reciprocal transplant experiment.** We reciprocally transplanted both soil and local *A. thaliana*
465 genotypes between sites in Italy (population IT4) and Sweden (population SW4; **Supplementary Fig.**
466 **7a**). See ref.¹⁵ for a description of study sites and plant genotypes. Soil was collected at the two
467 experimental sites in spring 2016 and stored at 6°C until the establishment of the experiments. Seeds
468 were planted in Petri dishes on agar, cold stratified in the dark at 6°C for one week, and then moved to
469 a growth room (22°C/16°C, 16 h day at 150 µE/m²/sec PAR, 8 h dark) where the seeds germinated.
470 Nine days after germination, seedlings were transplanted to 299-cell plug trays (cell size: 20 mm × 20
471 mm × 40 mm) with local and non-local soils in blocks of 6 x 7 cells in a checkerboard design (three
472 blocks of each soil type per tray). At each site, we transplanted 20 local and 22 non-local seedlings into
473 randomized positions in each of six blocks per soil type, with a total of 120 local and 132 non-local
474 plants for each site × soil combination. During transplantation, plug trays were kept in a greenhouse at
475 about 18°C/12°C and 16 h day/8 h night. Within six days, the trays were transported to the field sites
476 where they were sunk into the ground (on 9 September 2016 in Sweden, and on 29 October 2016 in

477 Italy). The transplanted seedlings were at approximately the same stage of development as naturally
478 germinating plants in the source population.

479
480 At fruit maturation in spring 2017, we scored survival to reproduction and number of fruits per
481 reproducing plant (an estimate of fecundity), and estimated total fitness as the number of fruits produced
482 per seedling planted (following ref.¹⁵). Statistical analyses were based on block means for each genotype.
483 No Italian plant survived at the Swedish site, and it was therefore not possible to fit a model examining
484 the site × soil × genotype interaction. Instead, we conducted analyses separately by site. For the Italian
485 site, we assessed differences in overall fitness due to soil, genotype, and the soil × genotype using a
486 mixed-model ANOVA, with block (random factor) nested within soil type. In Sweden, we used a one-
487 tailed t-test to test the prediction that survival and overall fitness of the Swedish genotype is higher when
488 planted in Swedish than when planted in Italian soil.

489
490 At the time of fitness assessment, we harvested plants and their surrounding soil by removing the whole
491 soil plug. Soil was separated from the roots manually and a soil sample was taken. After removing loose
492 soil particles by gentle shaking, roots were placed in a 15-mL falcon and washed by inverting with 10
493 mL of deionized water and surface-sterilized by washing for one minute with 70% ethanol, then washing
494 with 3% NaClO for one minute and rinsed several times with deionized sterile water. These whole root
495 samples correspond to the combination of Root and RP compartments described for the European
496 transect survey. Six to twelve soil samples were harvested for each of the eight combinations of soil,
497 location and genotype (n = 72), as well as three to twelve whole root samples per condition (n = 59).
498 Note that no Italian plants survived at the Swedish site. In total, 131 root and soil samples were
499 harvested, stored in dry ice, and processed for DNA isolation and microbial community profiling.

500
501 **Microbial community profiling.** Total DNA was extracted from the aforementioned samples using the
502 FastDNA SPIN Kit for Soil (MP Biomedicals, Solon, USA). Samples were homogenized in Lysis
503 Matrix E tubes using the Precellys 24 tissue lyzer (Bertin Technologies, Montigny-le-Bretonneux,
504 France) at 6,200 rpm for 30 s. DNA samples were eluted in 60 µL nuclease-free water and used for

505 bacterial, fungal and oomycetal community profiling^{1, 3}. The concentration of DNA samples was
506 fluorescently quantified, diluted to 3.5 ng/µL, and used in a two-step PCR amplification protocol. In the
507 first step, V5–V7 of bacterial 16S rRNA (799F - 1192R), V2-V4 of bacterial 16S rRNA (341F – 806R),
508 fungal ITS1 (ITS1F - ITS2), fungal ITS2 (fITS7 – ITS4) and oomycetal ITS1 (ITS1-O - 5.8 s-Rev-O)
509 (**Supplementary Table 2**) were amplified. Under a sterile hood, each sample was amplified in triplicate
510 in a 25 µl reaction volume containing 2 U DFS-Taq DNA polymerase, 1x incomplete buffer (both Bioron
511 GmbH, Ludwigshafen, Germany), 2 mM MgCl₂, 0.3% BSA, 0.2 mM dNTPs (Life technologies GmbH,
512 Darmstadt, Germany) and 0.3 µM forward and reverse primers. PCR was performed using the same
513 parameters for all primer pairs (94°C/2 min, 94°C/30 s, 55°C/30 s, 72°C/30 s, 72°C/10 min for 25
514 cycles). Afterward, single-stranded DNA and proteins were digested by adding 1 µl of Antarctic
515 phosphatase, 1 µl Exonuclease I and 2.44 µl Antarctic Phosphatase buffer (New England BioLabs
516 GmbH, Frankfurt, Germany) to 20 µl of the pooled PCR product. Samples were incubated at 37°C for
517 30 min and enzymes were deactivated at 85°C for 15 min. Samples were centrifuged for 10 min at 4000
518 rpm and 3 µl of this reaction were used for a second PCR, prepared in the same way as described above
519 using the same protocol but with cycles reduced to 10 and with primers including barcodes and Illumina
520 adaptors (**Supplementary Table 2**). PCR quality was controlled by loading 5 µl of each reaction on a
521 1% agarose gel. Afterwards, the replicated reactions were combined and purified: 1) bacterial amplicons
522 were loaded on a 1.5% agarose gel and run for 2 hours at 80 V; bands with the correct size of ~500 bp
523 were cut out and purified using the QIAquick gel extraction kit (QIAGEN, Hilden, Germany); 2) fungal
524 and oomycetal amplicons were purified using Agencourt AMPure XP beads. DNA concentration was
525 again fluorescently determined, and 30 ng DNA of each of the barcoded amplicons were pooled in one
526 library per microbial group. Each library was then purified and re-concentrated twice with Agencourt
527 AMPure XP beads, and 100 ng of each library were pooled together. Paired-end Illumina sequencing
528 was performed in-house using the MiSeq sequencer and custom sequencing primers (**Supplementary**
529 **Table 2**).

530

531 **16S rRNA gene and ITS read processing.** All paired rRNA amplicon sequencing reads have been
532 analysed with a pipeline described recently³. The main parts include scripts from QIIME⁵⁶ and usearch⁵⁷.

533 OTUs were clustered at a 97% threshold (usearch – cluster otus). Bacterial reads were checked against
534 the greengenes database⁵⁸ to remove non-bacterial reads. Fungal and oomycetal reads were checked
535 against an ITS sequences database (full length ITS sequences from NCBI) to remove unwanted reads.
536 Taxonomic assignment was done via QIIME (assign_taxonomy) for bacterial OTUs, using the
537 greengenes database. Taxonomic assignment for fungal OTUs was done via RDP-classifier⁵⁹ against the
538 Warcup database⁶⁰. Assignment of oomycetal OTUs was also done via RDP but against a self-
539 established database. For rRDP-based classification a cut-off of 0.5 was used to filter out uncertain
540 assignments.

541

542 **Computational analyses.** All OTU tables obtained from the before mentioned pipelines were filtered
543 for very low abundant OTUs prior any analysis and post processing. For this, only OTUs having at least
544 0.1% RA in at least one sample were kept. These filtered tables were used for all further analyses. For
545 OTU based analysis of bacterial data, OTUs assigned as chloroplast- or mitochondria- derived were
546 removed prior to analysis. Alpha diversity indices (Shannon index and observed OTUs) were calculated
547 using OTU tables rarefied to 1,000 reads. Bray Curtis distances between samples were calculated by
548 using OTU tables that were normalized using cumulative-sum scaling (CSS⁶¹). Average Bray-Curtis
549 distances across replicates and years, were calculated by using the mean across all combinations between
550 two sets of sites (e.g. between all soil samples from site x and site y). Comparisons of samples coming
551 from different years were not considered for this analysis. These average distances were hierarchically
552 clustered (hclust in R, method = “complete”). Similarly, averaged relative abundances (avg. RA) of
553 taxonomic groups was achieved by averaging across all samples from a particular site – compartment
554 combination. All principal coordinate analyses were calculated using the respective Bray-Curtis distance
555 matrices as an input for cmdscale function (standard R). Explained variance of different factors was
556 based on Bray-Curtis distances. These were used as an input for PERMANOVA analysis using the
557 adonis function in R (vegan package). All factors have been analysed independently. Dependency of
558 data was either tested via a linear regression (using lm function from R) or Spearman rank correlation
559 (cor.test function in R). Unless noted differently, a p value < 0.01 was considered significant.

560 Differential abundance of taxonomic groups was tested using the Wilcoxon rank sum test (wlicox.test
561 in R, FDR < 0.05).

562

563 For calculating site prevalence of OTUs, OTU tables were restricted to samples having > 1,000 reads.
564 Count tables were transformed to relative abundances by division of total column (sample) sums. An
565 OTU was marked as present in a given sample if its RA was > 0.1%. OTUs present in < 5 samples were
566 not considered further. Site prevalence reflects in how many sites (out of all European sites sampled in
567 a given year) an OTU was present on average across years (here only years where an OTU was detected
568 were taken into account). Site prevalence for OTUs present in neighbouring plants was calculated in the
569 same way. To calculate the average RA across sites and years for a given OTU, only those samples were
570 the OTU was actually present were considered. In this way, prevalence and mean RA are independent.
571 OTUs present in > 80% of sites on average were considered to be geographically widespread.
572 Differential abundance of OTUs in the different groups (geographically restricted, common, and
573 widespread) was calculated between compartment using Kruskal-Wallis test (kruskal.test with dunn.test
574 in R, FDR < 0.05). Root-specific enrichment for widespread OTUs, compared to soil and RS samples
575 was tested with a generalized linear model, as described in ref.²² (FDR, p < 0.05). To compare generalist
576 OTUs from this study with OTUs found in roots of *Lotus japonicus*²⁶ bacterial OTUs were directly
577 compared using the representative sequences (Blastn, 98% sequence identity). To compare fungal
578 OTUs, the representative sequences from geographically widespread OTUs were blasted against full
579 ITS sequences from the UNITE database⁶². The best hits received were then used to find similar OTUs
580 in the *L. japonicus* data (98% sequence identity).

581

582 Enrichment patterns of root associated OTUs in the transplant experiment (see heatmap in
583 **Supplementary Fig. 8**) were examined as follows. For bacterial and fungal datasets, samples with less
584 than 1,000 reads were removed, whereas for the oomycetal dataset, samples with less than 250 reads
585 were removed. OTUs having a mean relative abundance > 0.01% across all root samples were kept for
586 the analysis. Relative abundance entries of zero were replaced by 0.001%. All relative abundance values
587 were log2 transformed and these data were used as input for generating a heatmap (heatmap.2 function

588 in R, gplots library). Enrichment in one of the six tested conditions (Swedish location, Italian location,
589 Swedish soil, Italian soil, Italian location and soil, Swedish location and soil) was estimated by
590 comparing the mean relative abundance of each OTU across all samples with the mean relative
591 abundance in respective sample combinations. E.g. to be enriched at the Italian location, an OTU must
592 be more abundant at the Italian location compared to the Swedish location, irrespective of the soil (see
593 panel in **Supplementary Fig. 8**). OTUs that are not enriched in any of the conditions but are prevalent
594 across samples (mean relative abundance above 0.1%) are marked in grey in the heatmap. OTUs that
595 were not prevalent and not enriched in any condition are not shown. Geographically widespread OTUs
596 were identified by comparing representative sequences from the transect data with those of the reciprocal
597 transplant experiment (Blast at 98% sequence identity).

598

599 **Data availability**

600 Sequencing reads of samples from the European transect experiment and the reciprocal transplant
601 experiment (MiSeq 16S rRNA and ITS reads) have been deposited in the European Nucleotide Archive
602 (ENA) under accession numbers ENA: ERP115101 and ENA: ERP115102, respectively.

603

604 **Code availability**

605 All scripts for computational analysis and corresponding raw data are available at
606 <https://github.com/ththi/European-Root-Suppl>.

607

608 **Acknowledgements**

609 We thank Neysan Donnelly for scientific English editing. This work was supported by funds from the
610 Max Planck Society to S.H. and P.S.-L., a European Research Council starting grant ([MICRORULES](#))
611 to S.H., a European Research Council advanced grant ([ROOTMICROBIOTA](#)), and the “Cluster of
612 Excellence on Plant Sciences” program funded by the Deutsche Forschungsgemeinschaft to P.S.-L., and
613 grants from the Swedish Research Council to JÅ. C.A.-B. lab was funded by grant BIO2016-75754-P
614 (AEI/FEDER).

615

616 **Author contributions**

617 S.H., P.S-L. and J.Å. conceived the project. E.K., F.R., C.A-B., J.Å., and S.H. identified natural *A.*
618 *thaliana* populations. P.D. and S.H. collected the samples. P.D. prepared all samples and performed
619 microbial community profiling. P.D. and T.T. analyzed microbiota data. T.E. and J.Å. prepared the field
620 reciprocal transplant experiment. J.Å., T.E., and P.D. analyzed plant fitness data. S.H. supervised the
621 project. T.T., P.D., J.Å., P.S-L., and S.H. wrote the paper.

622

623 **Competing Interests**

624 The authors declare no competing interests.

625

626 **References**

- 627 1. Agler, M. T. et al. Microbial Hub Taxa Link Host and Abiotic Factors to Plant Microbiome
628 Variation. *PLOS Biol.* **14**, e1002352 (2016).
- 629 2. Hassani, M. A., Durán, P. & Hacquard, S. Microbial interactions within the plant holobiont.
630 *Microbiome* **6**, 58 (2018).
- 631 3. Durán, P. et al. Microbial Interkingdom Interactions in Roots Promote *Arabidopsis* Survival.
632 *Cell* **175**, 973–983.e14 (2018).
- 633 4. Berendsen, R. L., Pieterse, C. M. J. & Bakker, P. A. H. M. The rhizosphere microbiome and
634 plant health. *Trends Plant Sci.* **17**, 478–486 (2012).
- 635 5. Berendsen, R. L. et al. Disease-induced assemblage of a plant-beneficial bacterial consortium.
636 *ISME J.* **12**, 1496–1507 (2018).
- 637 6. Lebeis, S. L. et al. Salicylic acid modulates colonization of the root microbiome by specific
638 bacterial taxa. *Science* **349**, 860–4 (2015).
- 639 7. Haichar, F. el Z. et al. Plant host habitat and root exudates shape soil bacterial community
640 structure. *ISME J.* **2**, 1221–1230 (2008).
- 641 8. Stringlis, I. A. et al. MYB72-dependent coumarin exudation shapes root microbiome assembly
642 to promote plant health. *Proc. Natl. Acad. Sci.* **115**, E5213–E5222 (2018).

643 9. Bahram, M. et al. Structure and function of the global topsoil microbiome. *Nature* **560**, 233–
644 237 (2018).

645 10. Fierer, N. & Jackson, R. B. The diversity and biogeography of soil bacterial communities. *Proc.*
646 *Natl. Acad. Sci.* **103**, 626–631 (2006).

647 11. Karimi, B. et al. Biogeography of soil bacteria and archaea across France. *Sci. Adv.* **4**, eaat1808
648 (2018).

649 12. Delgado-Baquerizo, M. et al. Ecological drivers of soil microbial diversity and soil biological
650 networks in the Southern Hemisphere. *Ecology* **99**, 583–596 (2018).

651 13. Tedersoo, L. et al. Global diversity and geography of soil fungi. *Science* **346**, 1256688 (2014).

652 14. Leimu, R. & Fischer, M. A Meta-Analysis of Local Adaptation in Plants. *PLoS One* **3**, e4010
653 (2008).

654 15. Ågren, J. & Schemske, D. W. Reciprocal transplants demonstrate strong adaptive differentiation
655 of the model organism *Arabidopsis thaliana* in its native range. *New Phytol.* **194**, 1112–1122
656 (2012).

657 16. Fournier-Level, A. et al. A map of local adaptation in *Arabidopsis thaliana*. *Science* **334**, 86–9
658 (2011).

659 17. Wadgymar, S. M. et al. Identifying targets and agents of selection: innovative methods to
660 evaluate the processes that contribute to local adaptation. *Methods Ecol. Evol.* **8**, 738–749
661 (2017).

662 18. Brachi, B. et al. Investigation of the geographical scale of adaptive phenological variation and
663 its underlying genetics in *Arabidopsis thaliana*. *Mol. Ecol.* **22**, 4222–4240 (2013).

664 19. Wagner, M. R. et al. Natural soil microbes alter flowering phenology and the intensity of
665 selection on flowering time in a wild *Arabidopsis* relative. *Ecol. Lett.* **17**, 717–26 (2014).

666 20. Panke-Buisse, K., Poole, A. C., Goodrich, J. K., Ley, R. E. & Kao-Kniffin, J. Selection on soil
667 microbiomes reveals reproducible impacts on plant function. *ISME J.* **9**, 980–989 (2015).

668 21. Lu, T. et al. Rhizosphere microorganisms can influence the timing of plant flowering.
669 *Microbiome* **6**, 231 (2018).

670 22. Robbins, C. et al. Root-Associated Bacterial and Fungal Community Profiles of *Arabidopsis*
671 *thaliana* Are Robust Across Contrasting Soil P Levels. *Phytobiomes J.* **2**:1: 24-34 (2018).

672 23. Fitzpatrick, C. R., Mustafa, Z. & Viliunas, J. Soil microbes alter plant fitness under competition
673 and drought. *J. Evol. Biol.* **32**, 438-45 (2019).

674 24. Frachon, L. et al. Intermediate degrees of synergistic pleiotropy drive adaptive evolution in
675 ecological time. *Nature Ecol. Evol.* **1**: 1551-1561 (2017).

676 25. Busoms, S. et al. Salinity Is an Agent of Divergent Selection Driving Local Adaptation of
677 *Arabidopsis* to Coastal Habitats. *Plant Physiol.* **168**, 915–29 (2015).

678 26. Zgadzaj, et al. *Lotus japonicus* symbiosis signaling genes and their role in the establishment of
679 root-associated bacterial and fungal communities. *BioRxiv*. doi: <https://doi.org/10.1101/547687>
680 (2019).

681 27. Chaw SM, Chang CC, Chen HL, Li WH. Dating the monocot-dicot divergence and the origin
682 of core eudicots using whole chloroplast genomes. *J Mol Evol.* **58**, 424-41 (2004).

683 28. Ågren, J., Oakley, C. G., McKay, J. K., Lovell, J. T. & Schemske, D. W. Genetic mapping of
684 adaptation reveals fitness tradeoffs in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U. S. A.* **110**,
685 21077–82 (2013).

686 29. Schlaeppi, K., Dombrowski, N., Oter, R. G., Ver Loren van Themaat, E. & Schulze-Lefert, P.
687 Quantitative divergence of the bacterial root microbiota in *Arabidopsis thaliana* relatives. *Proc.*
688 *Natl. Acad. Sci. U. S. A.* **111**, 585–92 (2014).

689 30. Edwards, J. et al. Structure, variation, and assembly of the root-associated microbiomes of rice.
690 *Proc. Natl. Acad. Sci.* **112**, E911–E920 (2015).

691 31. Fitzpatrick, C. R. et al. Assembly and ecological function of the root microbiome across
692 angiosperm plant species. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E1157–E1165 (2018).

693 32. Yeoh, Y. K. et al. Evolutionary conservation of a core root microbiome across plant phyla along
694 a tropical soil chronosequence. *Nat. Commun.* **8**, 215 (2017).

695 33. Xu, L. et al. Drought delays development of the sorghum root microbiome and enriches for
696 monoderm bacteria. *Proc. Natl. Acad. Sci.* **115**, E4284–E4293 (2018).

697 34. Hacquard, S. et al. Microbiota and Host Nutrition across Plant and Animal Kingdoms. *Cell Host*
698 *Microbe* **17**, 603–616 (2015).

699 35. Levy, A. et al. Genomic features of bacterial adaptation to plants. *Nat. Genet.* **50**, 138–150
700 (2018).

701 36. Karasov, T. L. et al. *Arabidopsis thaliana* and *Pseudomonas* Pathogens Exhibit Stable
702 Associations over Evolutionary Timescales. *Cell Host Microbe* **24**, 168–179.e4 (2018).

703 37. Kwak, M.-J. et al. Rhizosphere microbiome structure alters to enable wilt resistance in tomato.
704 *Nat. Biotechnol.* **36**, 1100–1109 (2018).

705 38. Garrido-Oter, R. et al. Modular Traits of the Rhizobiales Root Microbiota and Their
706 Evolutionary Relationship with Symbiotic Rhizobia. *Cell Host Microbe* **24**, 155–167.e5 (2018).

707 39. Poupin, M. J., Timmermann, T., Vega, A., Zuñiga, A. & González, B. Effects of the Plant
708 Growth-Promoting Bacterium *Burkholderia phytofirmans* PsJN throughout the Life Cycle of
709 *Arabidopsis thaliana*. *PLoS One* **8**, e69435 (2013).

710 40. Bai, Y., et al. Functional overlap of the *Arabidopsis* leaf and root microbiota. *Nature* **528**, 364–
711 369 (2015).

712 41. Anderson, L.C. et al. 2014. Bacteria and Fungi Respond Differently to Multifactorial Climate
713 Change in a Temperate Heathland, Traced with ¹³C-Glycine and FACE CO₂. *PLoS One* **9**,
714 e85070 (2014)

715 42. Peay, K. G. et al. Convergence and contrast in the community structure of Bacteria, Fungi and
716 Archaea along a tropical elevation–climate gradient. *FEMS Microbiol. Ecol.* **93**, (2017).

717 43. Talbot, J. M. et al. Endemism and functional convergence across the North American soil
718 mycobiome. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 6341–6 (2014).

719 44. Coleman-Derr, D. et al. Plant compartment and biogeography affect microbiome composition
720 in cultivated and native *Agave* species. *New Phytol.* **209**, 798–811 (2016).

721 45. Lundberg, D. S. et al. Defining the core *Arabidopsis thaliana* root microbiome. *Nature* **488**, 86–
722 90 (2012).

723 46. Wagner, M. R. et al. Host genotype and age shape the leaf and root microbiomes of a wild
724 perennial plant. *Nat. Commun.* **7**, 12151 (2016).

725 47. Cregger, M. A. et al. The *Populus* holobiont: dissecting the effects of plant niches and genotype
726 on the microbiome. *Microbiome* **6**, 31 (2018).

727 48. Wright, K. M. et al. Adaptation to heavy-metal contaminated environments proceeds via
728 selection on pre-existing genetic variation. *bioRxiv* 29900 (2015)

729 49. Wright, J. W., Stanton, Maureen L. & Scherson, R. Local adaptation to serpentine and non-
730 serpentine soils in *Collinsia sparsiflora*. *Evol. Ecol. Res.* **8**, 1–21 (2006).

731 50. Busoms, S. et al. Fluctuating selection on migrant adaptive sodium transporter alleles in coastal
732 *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E12443–E124522018 (2018).

733 51. Postma, F. M. & Ågren, J. Early life stages contribute strongly to local adaptation in *Arabidopsis*
734 *thaliana*. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 7590–5 (2016).

735 52. Postma, F. M. & Ågren, J. Among-year variation in selection during early life stages and the
736 genetic basis of fitness in *Arabidopsis thaliana*. *Mol. Ecol.* **27**, 2498–2511 (2018).

737 53. Ågren, J., Oakley, C. G., Lundemo, S. & Schemske, D. W. Adaptive divergence in flowering
738 time among natural populations of *Arabidopsis thaliana*: Estimates of selection and QTL
739 mapping. *Evolution (N. Y.)* **71**, 550–564 (2017).

740 54. Méndez-Vigo B, Gomaa NH, Alonso-Blanco C, & Picó, FX. Among- and within- population
741 variation in flowering time of Iberian *Arabidopsis thaliana* estimated in field and glasshouse
742 conditions. *New Phytol.* **197**: 1332–1343 (2013)

743 55. Bartoli, C. et al. In situ relationships between microbiota and potential pathobiota in *Arabidopsis*
744 *thaliana*. *ISME J.* **12**, 2024–2038 (2018).

745 56. Caporaso, J. G. et al. QIIME allows analysis of high-throughput community sequencing data.
746 *Nat. Methods* **7**, 335–6 (2010).

747 57. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**,
748 2460–2461 (2010).

749 58. DeSantis, T. Z. et al. Greengenes, a chimera-checked 16S rRNA gene database and workbench
750 compatible with ARB. *Appl. Environ. Microbiol.* **72**, 5069–72 (2006).

751 59. Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Naive Bayesian classifier for rapid
752 assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.* **73**,
753 5261–7 (2007).

754 60. Deshpande, V. et al. Fungal identification using a Bayesian classifier and the Warcup training
755 set of internal transcribed spacer sequences. *Mycologia* **108**, 1–5 (2016).

756 61. Paulson, J. N., Stine, O. C., Bravo, H. C. & Pop, M. Differential abundance analysis for
757 microbial marker-gene surveys. *Nat. Methods* **10**, 1200–2 (2013).

758 62. Nilsson, R. H. et al. The UNITE database for molecular identification of fungi: handling dark
759 taxa and parallel taxonomic classifications. *Nucleic Acids Res.* **47**, D259–D264 (2019).

760

761

762

763

764

765

766

767

768

769

770

771

772


773

774

775

776

777



803

804

805

Fig. 2: Geographically widespread taxa in the roots of *A. thaliana* and grasses. **a**, Correlation between OTUs prevalence across sites in plant roots and averaged OTUs relative abundance (RA, \log_2) in plant roots. Bacteria: upper panel. Fungi: middle panel. Oomycetes: lower panel. For calculating averaged RA, only samples where the actual OTUs are present were considered. Blue: geographically restricted OTUs (site prevalence < 20%). Orange: geographically common OTUs (site prevalence 20–80%). Red: geographically widespread OTUs (site prevalence > 80%). The different shapes highlight root-associated OTUs detected one year, or across two or three years. RA and prevalence are averaged across the years where one OTU is present. **b**, Boxplots of the averaged RA (\log_2) of geographically restricted, common and widespread OTUs detected in each of the four compartments. Letters depict significant differences across compartments (Kruskal-Wallis test, FDR < 0.01). **c**, Phylogenetic trees of geographically widespread root-associated OTUs (red symbols in panel a) constructed based on the 16s rRNA V5V7 gene fragments (bacteria) and the ITS1 region (fungi, oomycetes). Microbial OTUs significantly enriched in root compared to soil samples are indicated with a star (FDR < 0.05) **d**, Geographically widespread OTUs detected in roots of *A. thaliana* and conserved signatures in roots of grasses and *Lotus japonicus*. The RA and proportion of widespread bacterial and fungal OTUs detected in *A. thaliana* roots are shown for *A. thaliana* (17 sites), co-occurring grasses (17 sites), as well as for *Lotus japonicus* grown in the Cologne Agricultural Soil (CAS). All shown OTUs have RA > 0.1%. The total RA of these OTUs in root samples is indicated below the circular diagrams.


824

825

826

827

828



829
830

831 **Fig. 3: Factors shaping the *A. thaliana* root microbiota at a continental scale.** **a**, Principal coordinate analysis (PCoA) based on Bray Curtis distances between samples harvested across 17 sites, four compartments and three successive years (2015, 2016, 2017). Microbial communities are presented for the whole *A. thaliana* dataset for bacteria (upper panel, 881 samples), fungi (middle panel, 893 samples), and oomycetes (lower panel, 875) and color-coded either according to the site or the harvesting year. OTUs with relative abundance < 0.1% were excluded from the dataset. **b**, Effect of site, harvesting year, host species, as well as of individual soil and environmental variables measured at each site, on bacterial, fungal, and oomycetal community composition. Explained variance (%) for each explanatory variable is shown for the different compartments and ranked according to the best explanatory variable in soil. The percentage of explained variance for each parameter was calculated based on permutational multivariate analysis of variance and only significant associations are shown (PERMANOVA, $p < 0.01$). RS: rhizosphere. RP: rhizoplane.

832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857





Fig. 4: Reciprocal transplant between two *A. thaliana* populations in Sweden and Italy. **a**, Schematic overview of the reciprocal transplant experiment. Soils and plant genotypes from IT1 and SW4 sites were reciprocally transplanted in the two locations (eight different treatment combinations). The symbols below the schematic view correspond to the symbols also used in the other panels. **b**, Community structures of bacteria, fungi, and oomycetes in the 131 samples were determined using principal coordinate analysis (PCoA). The first two dimensions of the PCoA are plotted based on Bray-Curtis distances. To facilitate visualization, the same PCoA plot was represented either according to the genotype and the compartment (upper panels) or according to the soil origin and the location (lower panels). Note that no Italian plant survived at the Swedish site. **c**, Overall fitness (number of fruits per seedling planted; mean \pm SE) of Italian and Swedish genotypes (red and blue color, respectively) when reciprocally planted in Italian and Swedish soils (circle and triangle symbols, respectively) and grown at Italian and Swedish locations (filled and open symbols, respectively).

896 **Supplementary Figures**

897

898



899

900

901


Supplementary Fig. 1: Validation of the root fractionation protocol and assessment of primer amplification bias. **a**, Protocol to fractionate four microbial niches across a distance gradient from bulk soil to roots' interior. Roots of *A. thaliana* grown in their natural environments were briefly washed (1) to separate loosely attached soil particles from the root surface (rhizosphere, RS). After a second washing step, roots were vigorously washed with detergent (three times) to capture microbes that tightly adhere to the root surface (2). The resulting washes were then filtered through a 0.22 μ M membrane (rhizoplane, RP). Finally, surface sterilization of root samples by consecutive EtOH and NaClO washes enriched the final root sample in microbial root endophytes (3). **b**, Validation of the fractionation protocol (depicted in panel a) was performed by printing root washes (left panel) and washed roots (right panel) on 50% Tryptic Soy Agar medium. Sequential detergent washes efficiently release microbes from the root surface and further root surface sterilization prevents the growth of rhizoplane-associated microbes (Wilcoxon rank sum test, $p < 0.01$). All three detergent steps were combined and filtered to prepare the RP fraction (light green dots, left panel). **c**, Comparison of bacterial (left panel) and fungal (right panel) classes profiled with the V2V4 and V5V7 regions of the bacterial 16s rRNA gene, and the ITS1 and ITS2 of the fungal ITS. The correlation between the relative abundances of each microbial class is shown (Pearson's correlation, $p < 0.001$).

918

919

920


921



922
923
924

925 **Supplementary Fig. 2: Microbial alpha diversity and enrichment signatures in plant roots.** **a**,
926 Microbial alpha diversity measured across all 17 sites in soil, rhizosphere (RS), rhizoplane (RP), and
927 datasets were rarefied to 1,000 reads. Individual data points within each box correspond to samples
928 from the 17 natural sites (Kruskal-Wallis test, $p < 0.05$). **b**, Microbial alpha diversity measured across
929 all 17 sites in soil, rhizosphere (RS), rhizoplane (RP), and root samples based on the number of observed
930 OTUs. All samples from a given site were taken into account and the datasets were rarefied to 1,000
931


932 reads. Individual data points within each box correspond to samples from the 17 natural sites (Kruskal-
933 Wallis test, $p < 0.05$). **c**, Comparison of taxa relative abundance (RA) between Swedish soils (SW1-4,
934 blue) and the other European soils (grey) for bacteria (left), fungi (middle), and oomycetes (right). RA
935 is aggregated at the class (bacteria and fungi) and order (oomycetes) levels and significant differences
936 are marked with an asterisk (Wilcoxon rank sum test, FDR < 0.05). **d**, Comparison of taxa RA between
937 soil (dark red) and root (dark green) samples for bacteria (left), fungi (middle), and oomycetes (right).
938 RA measured in soil and root samples across the 17 *A. thaliana* populations were aggregated at the class
939 (bacteria and fungi) and order (oomycetes) levels. Significant differences are marked with an asterisk
940 (Wilcoxon rank sum test, FDR < 0.05).
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984



985
986
987

988 **Supplementary Fig. 3: Geographically widespread taxa at the soil-root interface. a**, Correlation
989 between OTUs prevalence across sites in Soil, rhizosphere (RS), rhizoplane (RP), Root and averaged
990 OTUs relative abundance (RA, log2). Bacteria: upper panels. Fungi: middle panels. Oomycetes: lower
991 panels. Blue: geographically restricted OTUs (site prevalence < 20%). Orange: geographically common
992 OTUs (site prevalence 20-80%). Red: geographically widespread OTUs (site prevalence > 80%). For
993 calculating averaged RA, only samples where the actual OTUs are present were considered. The
994 different shapes highlight OTUs detected one year, or across two or three years. RA and prevalence
995 were averaged across the years where one OTU is present. OTUs with relative abundance < 0.1% were
996 excluded from the datasets. **b**, For each microbial group (bacteria, fungi, and oomycetes), Spearman's
997 rank correlations ($p < 0.01$) were determined between OTUs prevalence in roots of *A. thaliana* and
998 OTUs prevalence in roots of neighboring grasses. The geographically widespread OTUs detected in
999 roots of *A. thaliana* and grasses are indicated with numbers.

1000
1001



Supplementary Fig. 4: Influence of site, host species, and harvesting year on microbial community structure in *A. thaliana* populations. **a**, Principal coordinate analysis (PCoA) based on Bray Curtis distances for soil-, rhizosphere (RS), rhizoplane- (RP), and root-associated microbial communities detected in 17 sites across three successive years in European *A. thaliana* populations. Microbial communities in each compartment are presented for bacteria (upper panel), fungi (middle panel), and oomycetes (lower panel) and color-coded according to the site. **b**, PCoA based on Bray Curtis distances between samples (bacteria n = 881, fungi n = 803, oomycetes n = 875) harvested across 17 sites, four

1012 compartments and three successive years. Microbial communities are presented for the whole dataset
1013 for bacteria (upper panel), fungi (middle panel), and oomycetes (lower panel) and color-coded either
1014 according to the root pooling strategy (root from a single individual compared to pooled roots from four
1015 individuals), the harvesting year (2015, 2016, 2017), or the compartment. Additional soil, RP, RS, and
1016 Root samples (bacteria n = 238, fungi n = 241, oomycetes n = 236) from neighboring grasses were
1017 included in the PCoA plot where samples are color coded according to the host species. OTUs with
1018 relative abundance < 0.1% were excluded from the datasets.

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061


1062

1063

1064

1065

1066



1067
1068
1069

1070 **Supplementary Fig. 5: Association between local factors and microbial abundance profiles.** **a**,
1071 Heatmap showing the distribution of soil properties among each of the 17 sites. Real property values
1072 were normalized (0 = lowest measured value, 1 = highest measured value). **b**, Heatmap showing
1073 significant correlations detected between properties using data from all sites (Spearman's rank
1074 correlation, $p < 0.01$). **c**, Correlation between properties and RA of bacterial taxa (aggregated at class
1075 level) in soil samples (left panel) and root samples (right panel) (Spearman's rank correlation, $p < 0.01$).
1076 **d**, Correlation between properties and RA of fungal taxa (aggregated at class level, if available) in soil

1077 samples (left panel) and root samples (right panel) (Spearman's rank correlation, $p < 0.01$). **e**,
1078 Correlation between properties and RA of oomycetal taxa (aggregated at order level, if available) in soil
1079 samples (left panel) and root samples (right panel) (Spearman's rank correlation $p < 0.01$). Size of circles
1080 in **c**, **d**, and **e** is proportional to measured r values. The respective barplots show the cumulative
1081 correlation score for each variable.

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124


1125

1126


1127

1128

1129



Supplementary Fig. 6: Correlation between bacterial and fungal alpha diversity across compartments. **a**, Spearman's rank correlation between the number of observed fungal OTUs and the number of observed bacterial OTUs in soil, rhizosphere (RS), rhizoplane (RP), and root samples. All soil, RS, RP, and root samples from the 17 sites were taken into account and the datasets were rarefied to 1,000 reads (soil: n = 212, RS: n = 197, RP: n = 104, Root: n = 131). The Spearman's rank correlation coefficient and associated p-values are depicted above each graph. **b**, Spearman's rank correlation between the fungal Shannon index and the bacterial Shannon index in Soil, RS, RP, and root samples. All samples from the 17 sites were taken into account and the datasets were rarefied to 1,000 reads (soil: n = 212, RS: n = 197, RP: n = 104, root: n = 131). The Spearman's rank correlation coefficient and associated p-values are depicted above each graphs.



1164

1165

1166

1167 **Supplementary Fig. 7: Reciprocal transplant between two *A. thaliana* populations in Sweden and**
 1168 **Italy. a, European map showing names and locations of the 17 *A. thaliana* populations. The IT1 and**
 1169 **SW4 sites selected for the reciprocal transplant experiment are highlighted in red and blue, respectively.**
 1170 **b, Schematic overview of the reciprocal transplant experiment. Soils and plant genotypes from IT1 and**
 1171 **SW4 sites were reciprocally transplanted in the two locations (eight different treatment combinations).**
 1172 **The symbols below the schematic view correspond to the symbols used in panels c and d. c, Fitness of**
 1173 **Italian and Swedish genotypes (red and blue color, respectively) when reciprocally planted in Italian and**
 1174 **Swedish soils (circle and triangle symbols, respectively) and grown at Italian and Swedish locations**
 1175 **(filled and open symbols, respectively). Plant survival, fecundity (number of fruits per reproducing**
 1176 **plant), and overall fitness (number of fruits per seedling planted). Means based on block means \pm SE**
 1177 **are given. Note that no Italian plant survived to reproduce at the Swedish site. d, Bray-Curtis distances**
 1178 **constrained by genotype for bacterial, fungal, and oomycetal communities in whole root samples**
 1179 **(cPCoA, see axis 2). Results are shown for Italian and Swedish genotypes (red and blue color,**
 1180 **respectively) planted in Italian and Swedish soils (circle and triangle symbols, respectively) at IT1 site**
 1181 **only since no Italian plant survived at the SW4 site. The percentage of variation explained by the two**
 1182 **genotypes is plotted along the second axis and refers to the fraction of the total variance of the data that**
 1183 **is explained by the constrained factor (i.e: genotype; bacteria $p = 0.001$; fungi $p = 0.026$; oomycetes p**
 1184 **= 0.002**).

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194


1195

1196

1197

1198

1199



1200

1201

1202

1203

Supplementary Fig. 8: OTU distribution pattern across root samples in the transplant experiment.

a, Heatmap depicting the relative abundance (\log_2) of bacterial OTUs in roots of Italian and Swedish genotypes grown in Italian and Swedish soils at IT1 and SW4 locations. OTUs and samples are hierarchical clustered. Enrichment patterns of each OTU was estimated according to the categories described in the lower right side of the figure and highlighted with different colours next to the heatmap. The relative abundance of OTUs falling into one of the six categories is always higher in that category compared to the mean relative abundance measured across all samples. OTUs that are present in all samples (relative abundance $> 0.1\%$) and did not fall in any of the six categories are marked in grey. The heatmap is filtered for OTUs that have at least an average relative abundance of 0.01% across all root samples. Samples have been filtered to contain at least 1,000 reads. Genotype of plants for each sample is indicated below each heatmap. Blue: Swedish genotype. Red: Italian genotype. Note that no Italian plant survived at the Swedish site. **b**, Heatmap depicting the relative abundance (\log_2) of fungal OTUs in roots of Italian and Swedish genotypes grown in Italian and Swedish soils at IT1 and SW4 locations. OTUs and samples are hierarchical clustered. Enrichment patterns of each OTU was estimated according to the categories described in the lower right side of the figure and highlighted with different colours next to the heatmap. The relative abundance of OTUs falling into one of the six categories is always higher in that category compared to the mean relative abundance measured across all samples. OTUs that are present in all samples (relative abundance $> 0.1\%$) and did not fall in any of the six categories are marked in grey. The heatmap is filtered for OTUs that have at least an average relative abundance of 0.01% across all root samples. Samples have been filtered to contain at least 1,000 reads. Genotype of plants for each sample is indicated below each heatmap. Blue: Swedish genotype. Red: Italian genotype. Note that no Italian plant survived at the Swedish site. **c**, Heatmap depicting the relative abundance (\log_2) of Oomycete OTUs in roots of Italian and Swedish genotypes grown in Italian and Swedish soils at IT1 and SW4 locations. OTUs and samples are hierarchical clustered. Enrichment patterns of each OTU was estimated according to the categories described in the lower right side of the figure and highlighted with different colours next to the heatmap. The relative abundance of OTUs falling into one of the six categories is always higher in that category compared to the mean relative abundance measured across all samples. OTUs that are present in all samples (relative abundance $> 0.1\%$) and did not fall in any of the six categories are marked in grey. The heatmap is filtered for OTUs that have at least an average relative abundance of 0.01% across all root samples. Samples have been filtered to contain at least 1,000 reads. Genotype of plants for each sample is indicated below each heatmap. Blue: Swedish genotype. Red: Italian genotype. Note that no Italian plant survived at the Swedish site.

1216 locations. **c**, Heatmap depicting the relative abundance (log2) of oomycetal OTUs in roots of Italian and
1217 Swedish genotypes grown in Italian and Swedish soils at IT1 and SW4 locations **d**, Percentage of OTUs
1218 falling into one of the six categories are presented as pie charts for each main taxonomic classes. The
1219 number of OTUs that belong to each microbial class is given in brackets.

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268 **Supplementary Tables**

1269

1270 **Supplementary table 1: European sites from where *Arabidopsis thaliana* and grasses populations
1271 were harvested.**

1272

1273 **Supplementary table 2: Primers utilized in this study to profile bacterial, fungal and oomycetal
1274 communities in soil and root samples.**

1275

1276 **Supplementary Table 3: Microbial communities' variation explained by several factors across
1277 all compartments.**

1278

1279 **Supplementary Table 4: Description of geographically widespread OTUs detected in *A. thaliana*
1280 Root samples.**

1281

1282 **Supplementary Table 5: Microbial communities' variation explained by several factors and
1283 environmental variables for each individual compartment.**

1284

1285 **Supplementary Table 6: Microbial communities' variation explained by host species at each site
1286 in the root compartment.**

1287

1288 **Supplementary table 7: Microbial communities' variation explained by compartment and by soil,
1289 location, and genotype in a field reciprocal transplant experiment**

1290

1291 **Supplementary Table 8. Survival, number of fruits produced by survivors, and the number of
1292 fruits per seedling planted in the reciprocal transplant experiment conducted at the sites of the
1293 IT1 and SW4 populations.**

1294

1295 **Supplementary Table 9. Analysis of effects of soil (Italian vs. Swedish), and genotype (Italian vs.
1296 Swedish) on total fitness (number of fruits per seedling planted) in a field experiment conducted
1297 at the site of the Italian genotype.**