

ASSOCIATIONS OF ‘RELATIVE CORTICOSTERONE DEFICIENCY’ WITH GENETIC VARIATION IN *CYP17A1* AND METABOLIC SYNDROME FEATURES

Scott D Mackenzie^{1*}, Andrew A Crawford^{1,2,3*}, Daniel Ackermann⁴, Katharina E Schraut¹, Caroline Hayward⁵, Jennifer L Bolton¹, Christopher Saunders¹, Emad Al-Dujaili¹, Bernhard Dick⁴, Geneviève Escher⁴, Bruno Vogt⁴, Menno Pruijm⁶, Belen Ponte⁷, James F Wilson^{5,8}, Mark W J Strachan¹, Jackie F Price⁸, David I W Phillips⁹, Scott M MacKenzie¹⁰, Eleanor Davies¹⁰, Rebecca M Reynolds¹, and Brian R Walker^{1,11}

*These authors contributed equally

(1) BHF Centre for Cardiovascular Science, Queen’s Medical Research Institute, University of Edinburgh, United Kingdom, (2) MRC Integrative Epidemiology Unit at the University of Bristol, Bristol, UK (3) Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK (4) University Clinic for Nephrology, Hypertension and Clinical Pharmacology, Inselspital, Bern University Hospital, University of Bern, Switzerland, (5) MRC Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, United Kingdom, (6) Service of Nephrology, Lausanne University Hospital (CHUV), Switzerland, (7) Service of Nephrology, University Hospitals of Geneva (HUG), Switzerland, (8) Centre for Global Health Research, Usher Institute for Population Health Sciences and Informatics, University of Edinburgh, United Kingdom, (9) MRC Lifecourse Epidemiology Unit, Southampton General Hospital, United Kingdom, (10) Institute of Cardiovascular and Medical Sciences, University of Glasgow, United Kingdom. (11) Institute of Genetic Medicine, Newcastle University, Newcastle upon Tyne, UK

21

22 ABBREVIATED TITLE: Corticosterone, *CYP17A1* and metabolic syndrome

23 KEYWORDS: Glucocorticoids; Metabolic Syndrome; 17 α -Hydroxylase; *CYP17A1*

24 CORRESPONDENCE (including for reprint requests):

25 Professor Brian R Walker

26 Newcastle University

27 Executive Office

28 King's Gate

29 Newcastle upon Tyne

30 NE1 7RU

31 UK

32

33 Tel +44 (0)191 208 7701

34 Email: Brian.Walker@ncl.ac.uk

35

36 FINANCIAL DISCLOSURE:

37 The Wellcome Trust Scottish Translational Medicine and Therapeutics Initiative (STMTI), Medical

38 Research Council, British Heart Foundation, Wellcome Trust and the Swiss National Science

39 Foundation (FN 33CM30-124087, Schweizerischer Nationalfonds (SNF), Wildhainweg 3, Postfach

40 8232, 3001 Bern, Switzerland). The funders had no role in study design, data collection and analysis,

41 decision to publish, or preparation of the manuscript.

42 COMPETING INTERESTS:

43 The authors have declared that no competing interests exist.

ABSTRACT

Context and objective. Common genetic variants in *CYP17A1* associate with higher blood pressure, putatively from impaired 17 α -hydroxylase activity and mineralocorticoid excess. However, the same variants protect against obesity and insulin resistance. We tested whether *CYP17A1* variants that enhance 17 α -hydroxylase activity cause ‘relative corticosterone deficiency’. Since corticosterone is thought to contribute disproportionately to negative feedback in the hypothalamic-pituitary-adrenal axis, we also tested whether lower corticosterone associates with higher cortisol and hence with metabolic syndrome.

Design: Cross-sectional studies within the population-based Orkney Complex Disease Study (ORCADES; n=2018), VIKING Health Study Shetland (VIKING; n=2098), East Hertfordshire study (EHERTS; n=279), Edinburgh Type 2 Diabetes Study (ET2DS; n=903), and the Swiss Kidney Project on Genes in Hypertension (SKIPOGH; n=888).

Outcome measures. Cortisol and corticosterone in morning plasma samples in ORCADES, VIKING and ET2DS, and in EHERTS in plasma following overnight dexamethasone suppression (0.25mg) and 30 mins after ACTH₁₋₂₄ (1 μ g); cortisol and corticosterone metabolites in day and night urine samples in SKIPOGH. Features of the metabolic syndrome including body mass index, systolic blood pressure, lipid profile, fasting glucose, fasting insulin and HOMA-IR.

Results. In ORCADES, ET2DS and SKIPOGH, *CYP17A1* variants were associated with corticosterone:cortisol ratio. In ORCADES, VIKING and ET2DS there were consistent associations of morning plasma cortisol and corticosterone with BMI, blood pressure, lipid profile, fasting glucose and HOMA-IR. In EHERTS, however, after dexamethasone suppression and ACTH₁₋₂₄ stimulation, impaired glucose tolerance and insulin sensitivity were associated with higher cortisol but lower corticosterone. Similarly, in SKIPOGH, low corticosterone:cortisol metabolite ratios were associated with high BMI and dyslipidemia.

Conclusions. ‘Relative corticosterone deficiency’, due to a primary alteration in adrenal steroidogenesis favouring cortisol over corticosterone, may mediate the associations of genetic variation in *CYP17A1* with metabolic syndrome. However, additional determinants of variation in plasma corticosterone are likely to explain its generally positive associations with features of metabolic syndrome.

INTRODUCTION

In genome-wide association studies (GWAS) common variants in the *CYP17A1* locus are consistently associated with hypertension (1-4) and cardiovascular disease (5, 6). *CYP17A1* encodes the steroidogenic enzyme CYP17A1, which is expressed in human but not rat or mouse adrenal cortex and catalyses 17 α -hydroxylase activity, converting precursors for synthesis of 11-deoxycorticosterone and corticosterone into 17-hydroxylated precursors for synthesis of 11-deoxycortisol and cortisol. Rare mutations causing near-absent CYP17A1 activity result in low-renin hypertension (7), attributed to accumulation of the mineralocorticoid 11-deoxycorticosterone. Ratios of steroid metabolites reflecting CYP17A1 activity are highly heritable (8). Associations of more common variation in *CYP17A1* with hypertension might therefore be explained by reduced 17 α -hydroxylase activity.

Intriguingly, *CYP17A1* risk alleles for hypertension are also associated with lower body mass index (BMI) (9-11) and enhanced insulin sensitivity (12). Conversely, alleles which are protective for hypertension, and therefore predicted to increase 17 α -hydroxylase activity, confer increased risk of obesity and insulin resistance. This paradox is unexplained, but may relate to variations in production of the major glucocorticoid, cortisol. Elevated cortisol in Cushing's syndrome causes obesity and insulin resistance, while plasma cortisol is more subtly increased in metabolic syndrome (13). However, if elevated 17 α -hydroxylase activity contributes to obesity and insulin resistance by increasing cortisol production, why would plasma cortisol concentration not be 'corrected' by enhanced negative feedback suppression of ACTH secretion and a compensatory fall in cortisol production? An explanation may be found by considering the role of the second glucocorticoid in humans, corticosterone.

Widely neglected in humans, corticosterone circulates at concentrations of 5-10 % those of cortisol. However, concentrations of corticosterone in human cerebrospinal fluid and brain are relatively high, at ~30% of cortisol (14, 15). This discrepancy has been attributed to differential trans-membrane trafficking of steroids by the ATP-binding cassette (ABC) transporter ABCB1, which is expressed in the human CNS and selectively exports cortisol rather than corticosterone (15). Corticosterone may therefore make a disproportionate contribution to negative feedback control of the hypothalamic-

pituitary-adrenal (HPA) axis within the CNS. Notably, the endogenous negative feedback signal is known to be impaired in people with metabolic syndrome (16).

We hypothesised that individuals with a genetically determined increase in 17 α -hydroxylase activity have relatively low corticosterone which causes impaired negative feedback to the HPA axis, and hence sustains higher plasma cortisol. We predict this pattern of ‘relative corticosterone deficiency’ is associated with obesity and insulin resistance but lower blood pressure. We tested this hypothesis in population-based cohorts, by studying the associations of corticosterone and cortisol in plasma and of their metabolites in urine with: (i) genetic variation in *CYP17A1*; and/or (ii) features of metabolic syndrome.

MATERIALS AND METHODS

Participants

All studies conformed with the Declaration of Helsinki and ethical approval and written informed consent were obtained.

Orkney Complex Disease Study (ORCADES)

The ORCADES study is a genetic epidemiological study based in the Scottish archipelago of Orkney, comprising 2039 subjects aged 18-100 years, with at least two Orcadian grandparents, recruited between 2005 and 2011 (17). Subjects attended a local or mobile venepuncture clinic between 0730h and 1100h (mean 0923 h $\pm$ SD 46 min), after fasting from 2200h the previous night. On another occasion subjects attended for measurement of weight, height and blood pressure. Genotyping was undertaken as described (17).

Viking Health Study – Shetland (VIKING)

The VIKING study is a population cohort study based in an isolated population in the north of Scotland that recruited 2105 volunteers between 2013 and 2015. Subjects were required to have at least two grandparents from the Shetlands in order to participate. Each participant attended a measurement clinic and a venepuncture clinic to give a fasting blood sample. Participants' DNA was genotyped using the Illumina HumanOmniExpressExome8v1-2_A.

Edinburgh Type 2 Diabetes Study (ET2DS)

The Edinburgh Type 2 Diabetes Study (ET2DS) is a prospective cohort study comprising 1066 men and women with type 2 diabetes, living in Lothian, Scotland. Recruitment and study design has been reported (18). Briefly, participants aged 60-75 years with a diagnosis of type 2 diabetes according to WHO criteria (19) were recruited from a clinical database. Following overnight fast, subjects attended a research clinic at 0800-0830h where they underwent venepuncture and physical examination. Genotyping was performed by KBioscience (Herts, UK) using a competitive allele-specific PCR system (KASPar).

East Hertfordshire Study (EHERTS)

All births in EHERTS from 1911 onwards were notified by the attending midwife (20). As described, these records were used to recruit 309 women and 370 men, with no history of diabetes, for a standard 75-g oral glucose tolerance test (OGTT), performed at a clinic at 0800-1020 h following overnight fast from 2100 h (21). In follow up studies (21), 312 of these individuals, aged 67-78 years, ingested 0.25 mg dexamethasone at 2200 h, fasted overnight and attended a local clinic at 0830h next morning. A baseline blood sample was obtained before 1.0µg ACTH₁₋₂₄ (tetracosactrin, Synacthen, Alliance, Chippenham, UK) was administered intravenously, and repeat samples obtained after 30 minutes.

Swiss Kidney Project on Genes in Hypertension (SKIPOGH)

The SKIPOGH study is a family-based cross-sectional study comprising 1093 subjects aged 18-82 years from 2 regions (Bern and Geneva) and 1 city (Lausanne) of Switzerland (8). A random sample of the inhabitants were invited to participate if they were of European ancestry and had at least 1 first degree family member also willing to participate. They attended hospital in the morning after an overnight fast for venepuncture and physical examination. Five consecutive blood pressure (BP) measurements were taken from the arm with the higher BP, and the average of the last four was used. Urine samples were collected separately for day- and night-time periods, as described (8). DNA samples were genotyped on the Illumina MetaboChip array.

Laboratory analyses

In ORCADES , VIKING, EHERTS, and SKIPOGH, the homeostasis model assessment was used to quantify insulin resistance (HOMA-IR) (22). Serum cortisol was measured by radioimmunoassay with Guildhay antisera (23). After exclusion of individuals prescribed glucocorticoid therapy within the previous 3 months (ORCADES n = 21; ETD2S n=163; EHERTS n = 33), there was sufficient sample for analysis of corticosterone in 2018 individuals from ORCADES, 2098 in VIKING, 903 in ETD2S, and 279 in EHERTS. An in-house radioimmunoassay, modified for microtiter plate scintillation proximity assay (SPA), was used to measure plasma corticosterone with assay characteristics as previously reported (24); cross-reactivity for dexamethasone and cortisol was less than 1%. Results were accepted if the coefficient of variation between duplicates was <15%. In SKIPOGH, urinary

steroid metabolites were extracted and analysed by gas chromatography-mass spectrometry (GC-MS) as previously described (8); analyses comprised 888 individuals after exclusion of 205 subjects with missing steroid metabolites. To estimate the apparent CYP17A1 activity, the ratio (tetrahydro-11-dehydrocorticosterone (THA) + tetrahydrocorticosterone (THB) + 5 α -tetrahydrocorticosterone (5 α THB)) / (tetrahydrocortisone (THE) + tetrahydrocortisol (THF) + 5 α -tetrahydrocortisol (5 α THF)) was calculated.

Statistical analysis

Associations with genotype were analysed using Stata v14.1 (Stata Statistical Software: Release 14. College Station, TX: StataCorp LP). Given limited statistical power, we restricted analysis to candidate single nucleotide polymorphisms (SNPs) in *CYP17A1* which have functional effects; these included rs2486758, located in the *CYP17A1* promoter region and associated with ovarian steroidogenesis (25, 26), and the other SNPs in Table 2 which affect CYP17A1 transcription *in vitro* (27). All analyses were adjusted for age and sex. Because clearance of glucocorticoids is increased and plasma cortisol decreased in obesity (28), relevant analyses were adjusted for BMI. Because timing of sampling varied in ORCADES, venesection time (as minutes after the first sample in the cohort) was included as a predictor variable in all models (29). In ORCADES many of the participants are related; analysis of associations of SNPs with plasma glucocorticoid concentrations and metabolic risk factors was therefore undertaken following adjustment after fitting both the first 3 principal components of ancestry and the kinship matrix using a mixed model (mmscore function of GenABEL for the association test in ProbABEL) (30) under an additive model. The kinship matrix used the identity-by-state function of GenABEL (using weight = “freq” option).

Other analyses for ORCADES, VIKING, ET2DS and EHERTS were undertaken using Minitab (version 16; State College PA). Continuous response variables were normalised by log transformation. Student's t-tests were used for group comparisons of corticosterone:cortisol ratios and responses to ACTH. Simple linear regression analysis of corticosterone or cortisol with a given 'response' (or dependent) variable was used to determine unadjusted p values for the regression co-efficient. Analyses were repeated with potentially confounding co-variables in multiple regression analyses, adjusting for age, sex, BMI and time of sampling. Histograms and normal plots of residuals were examined to confirm

the validity of the linear regression model. To maintain this validity, and to minimise confounding due to prescribed medication, those prescribed glucose-lowering medications (see Table 1 for numbers) were excluded from regression analyses involving glucose or insulin in the ORCADES, VIKING and EHERTS cohorts. Because the majority of participants in ET2DS were prescribed these agents, their prescription was encoded as a binary variable and entered into the regression equation. This binary variable approach was also used in all cohorts to adjust for the effect of relevant medications on lipids and blood pressure.

The magnitude of the association of plasma cortisol and corticosterone with response variables is presented using standardised Z scores of log transformed variables in the adjusted regression models. The coefficients from these regression models are interpreted as the SD change in log transformed outcome for a 1SD higher log transformed cortisol or corticosterone. This allows direct comparison of all associations investigated.

In SKIPOGH statistical analyses were conducted using STATA 14.0 (StataCorp, College Station, Texas, USA). Simple mixed linear or multiple regression analyses were used to explore associations of total urinary corticosterone (THA+THB+5 α -THB) or cortisol (THE+THF+5 α -THF) metabolites, while taking familial correlations into account using a random family-effect. In the multiple regression model, analyses were adjusted for age, sex, study center, BMI (unless BMI was the outcome variable) and relevant medication. (THA+THB+5 α THB)/(THE+THF+5 α THF) ratio (i.e. CYP17A1 ratio) was analysed after log transformation in a mixed linear model to investigate associations with variables of interest. Covariates included age, sex, study center, and antihypertensive, lipid- and glucose-lowering treatment, BMI (unless BMI was the outcome variable), estimated GFR, urine flow rate, urinary sodium, potassium and creatinine excretion (24h per kg body weight).

RESULTS

Participant characteristics are summarised in Table 1. Plasma cortisol and corticosterone were positively correlated in ORCADES and ET2DS cohorts (Figure 1A), in VIKING (Coefficient correlation = 0.08, $p < 0.001$) and in both dexamethasone-suppressed and ACTH-stimulated samples in EHERTS (Figure 1B), while urinary cortisol and corticosterone metabolites were positively correlated in SKIPOGH (Figure 1C and 1D). After ACTH stimulation in EHERTS, the increase from baseline was greater for corticosterone than cortisol (6.2 ± 0.3 vs 3.9 ± 0.4 fold, respectively; $p < 0.001$).

Associations of plasma corticosterone and cortisol with *CYP17A1* genotype

In ORCADES, associations of functional SNPs in *CYP17A1* with morning plasma cortisol and corticosterone are shown in Table 2. The minor allele (C; frequency 0.16) of rs2486758 was associated with higher plasma corticosterone and corticosterone:cortisol ratio but not with plasma cortisol. These associations were replicated in ET2DS, in which the minor allele (C; frequency 0.21) of rs2486758 similarly tended to be associated with higher plasma corticosterone ($\beta = 0.074 \pm 0.043$, $p = 0.09$) and corticosterone:cortisol ratio ($\beta = 0.078 \pm 0.040$, $p = 0.05$) but not with cortisol ($\beta = -0.004 \pm 0.017$, $p = 0.83$). Similar associations were observed in SKIPOGH with the minor allele (C; frequency 0.23) of rs2486758 associated with higher overnight urinary corticosterone ($\beta = 11.0 \pm 5.7$, $p = 0.05$) and with day and night urinary corticosterone:cortisol ratio ($\beta = 0.004 \pm 0.002$, $p = 0.03$; $\beta = 0.004 \pm 0.002$, $p = 0.05$), but not with daytime or overnight urinary cortisol ($\beta = -133.5 \pm 96.6$, $p = 0.17$; $\beta = 11.1 \pm 44.6$, $p = 0.80$). There was no evidence of an association between *CYP17A1* variants with morning plasma cortisol or corticosterone in the VIKING cohort.

Associations of morning plasma cortisol and corticosterone with features of metabolic syndrome

In ORCADES (Table 3 and Figure 2A), both higher plasma cortisol and higher plasma corticosterone were associated with lower BMI. After adjustment for BMI, age and sex, both higher plasma cortisol and corticosterone were associated with higher fasting plasma glucose and higher triglycerides, although the magnitude of this association was greater for cortisol than corticosterone. Moreover, higher cortisol, but not corticosterone, was associated with raised systolic blood pressure. Neither cortisol nor corticosterone was significantly associated with total cholesterol or LDL cholesterol. In contrast with

these predominant associations with cortisol, lower corticosterone but not cortisol was associated with lower HDL-cholesterol, while higher corticosterone but not cortisol was associated with higher fasting insulin and HOMA-IR. The corticosterone:cortisol ratio was positively associated with fasting glucose, insulin and HOMA-IR but not with BMI, blood pressure or lipid profile.

In VIKING (Table 3 and Figure 2B), similar associations were observed as in ORCADES. Both cortisol and corticosterone were negatively associated with BMI. Cortisol was more strongly positively associated with systolic blood pressure and triglycerides than corticosterone. The corticosterone:cortisol ratio was negatively associated with BMI and positively associated with fasting glucose.

In ET2DS (Table 3 and Figure 2C), fewer biochemical phenotypes were available, but both higher plasma cortisol and plasma corticosterone were associated with lower BMI and higher fasting plasma glucose, as in ORCADES. However, cortisol and corticosterone were unrelated to systolic blood pressure in ET2DS, and the corticosterone:cortisol ratio was not significantly associated with BMI, blood pressure or glucose.

Associations of ACTH-stimulated cortisol and corticosterone with features of metabolic syndrome

Given the limited associations of the corticosterone:cortisol ratio in morning plasma samples with features of metabolic syndrome, despite its robust association with *CYP17A1* genotype, we sought to test the association of 17 α -hydroxylase activity with features of metabolic syndrome using potentially more sensitive dynamic testing in the EHERTS cohort. See Table 3 and Figures 2D and 2E. After low dose overnight dexamethasone suppression followed by low dose ACTH₁₋₂₄ administration, plasma cortisol and corticosterone levels changed as expected (Table 1). Plasma cortisol was not associated with any features of metabolic syndrome, except that ACTH-stimulated values were positively correlated with plasma glucose 30 min after an oral glucose load. However, plasma corticosterone showed different associations with metabolic syndrome variables. After dexamethasone suppression, lower plasma corticosterone was associated with higher BMI and, independently, with higher fasting insulin and HOMA-IR. After ACTH stimulation, lower plasma corticosterone (and corticosterone:cortisol ratio) was associated with higher plasma glucose before and after an oral glucose

load, and with higher HOMA-IR. There were no significant associations between cortisol, corticosterone and blood pressure in the EHERTS cohort.

Associations of urinary cortisol and corticosterone metabolites with features of metabolic syndrome

In SKIPOGH (Table 4 and Figure 3), after adjustment for confounders (age, sex, study center, relevant medication and BMI), higher cortisol metabolite excretion during daytime and overnight was associated with higher BMI, during the day with higher LDL cholesterol and lower HDL cholesterol, and during the night with higher fasting glucose. Plasma triglycerides were, surprisingly, inversely associated with day and night urinary cortisol metabolite excretion. By contrast, corticosterone metabolite excretion during the day was inversely associated with insulin levels and plasma triglycerides, but during the night was positively associated with fasting glucose and inversely with plasma triglycerides. A lower ratio of corticosterone:cortisol metabolites during both day and night was strongly associated with higher BMI and LDL cholesterol, and during the night was associated with higher systolic BP, higher fasting triglycerides and insulin as well as lower HDL cholesterol levels.

DISCUSSION

This is the largest study to date investigating associations between endogenous glucocorticoids and features of metabolic syndrome. Using 3 cohorts comprising 3200 participants, we confirm previous observations (13, 31-34) that elevated morning plasma cortisol is associated with higher blood pressure and blood glucose in subjects with and without type 2 diabetes. Moreover, we confirm previous observations (35) in by far the largest sample to date that urinary cortisol metabolite excretion is increased in association with obesity and dyslipidemia. With a large number of participants, the chance of statistical errors is much reduced.

This is the first study to investigate whether similar associations exist for corticosterone. We measured plasma corticosterone and analysed variability in *CYP17A1* to determine whether ‘relative corticosterone deficiency’, favouring production of cortisol over corticosterone, underlies associations between variation in *CYP17A1* and features of metabolic syndrome identified in GWAS consortia. The sample size was too small to detect associations at genome-wide significance, but using a candidate gene approach we demonstrated that variation in *CYP17A1* associates with plasma corticosterone:cortisol ratio in a general adult population, and confirmed our findings in two additional cohorts (one in a general adult population and one in a cohort of patients with type 2 diabetes). Moreover, these observations were supported by an association of *CYP17A1* genotype with urinary corticosterone/cortisol metabolite ratios. The association was restricted to just one SNP of 3 that are known to have functional effects on transcription *in vitro* (27), rs2486758; the replication in a second independent cohort, however, makes it highly unlikely to be due to chance. rs2486758 is located in the 5’ regulatory region of *CYP17A1* and in LD with rs2150927 but less so with rs138009835 or the variants previously associated with hypertension by GWAS (27). Paradoxically, the major T allele of rs2486758, which we find associated with ‘relative corticosterone deficiency’, induced lower, rather than higher, *CYP17A1* expression *in vitro* (27) but this may be an artefact of the *in vitro* system or rs2486758 may be linked with other functional variants.

Previous large GWAS consortia have shown opposite directions of association between variants in *CYP17A1* with hypertension versus other features of metabolic syndrome. We therefore anticipated that ‘relative corticosterone excess’ is associated with higher blood pressure while ‘relative corticosterone

deficiency' is associated with cortisol-mediated insulin resistance and obesity. However, despite elevated corticosterone excretion rate being reported previously in hypertension (36), we did not find an association between morning plasma corticosterone and blood pressure. Moreover, our data show that morning plasma cortisol and corticosterone were positively correlated. Further, plasma cortisol and corticosterone showed similar associations with BMI; it is likely that reduced plasma corticosterone in obesity is explained by increased activity of A-ring reductases which also underlie increased clearance of cortisol in obesity (37). When we did find discrepancies in associations with plasma cortisol or corticosterone, it was generally higher, rather than lower, morning plasma corticosterone that was the stronger predictor than morning plasma cortisol, e.g. of fasting plasma glucose, plasma insulin and HOMA-IR. This suggests that *CYP17A1* genotype is not the major determinant of morning plasma corticosterone:cortisol ratio. It is possible that insulin resistance or deficiency causes a shift in steroidogenesis in favour of corticosterone rather than cortisol, since *CYP17A1* expression is up-regulated by insulin (38). Alternatively, the combined elevation of cortisol and corticosterone with insulin resistance could reflect central activation of the HPA axis, consistent with altered responses to habituation and stress (34).

Associations between morning plasma glucocorticoid levels and lipid profile provided limited support for our hypothesis of 'relative corticosterone deficiency'. Thus, elevated triglycerides were consistently associated with plasma cortisol and less strongly with plasma corticosterone, and low HDL-cholesterol was associated only with low corticosterone, although the latter association has been reported previously for cortisol (39). However, effects of glucocorticoids on lipid metabolism are complex (40), and previous studies investigating associations of glucocorticoid excess with dyslipidaemia have been inconsistent (13).

A different pattern emerged when we studied associations of corticosterone and cortisol in EHRTS subjects who had received dexamethasone to suppress any compensatory increase or primary drive to the HPA axis, and a fixed dose of ACTH₁₋₂₄ to stimulate cortisol and corticosterone. The results are striking, with opposite associations of cortisol and corticosterone with metabolic syndrome features, consistent with a primary shift in adrenocortical production in favour of cortisol rather than corticosterone in metabolic syndrome. Since, as a result of the activity of ABCB1 excluding cortisol

from the brain (15), corticosterone is thought to make a disproportionate contribution to negative feedback suppression of the HPA axis in humans, these observations could provide a key insight into the activation of the HPA axis and elevated plasma cortisol in metabolic syndrome. This insight is obscured by measurement of morning plasma corticosterone and cortisol alone, perhaps because of greater variability as a result of unmeasured confounding.

Fasting plasma corticosterone was low in the ET2DS cohort, whereas cortisol was similar to the values in the ORCADES cohort (mean corticosterone:cortisol was 3.0% in ET2DS vs. 7.5% in ORCADES, $p < 0.0001$). This is unlikely to be due to older age of the ET2DS cohort, since corticosterone:cortisol ratio was not associated with age ($r^2 = 0.01$, $p = 0.27$). Again, insulin deficiency might favour cortisol rather than corticosterone secretion, consistent with loss of up-regulation of *CYP17A1* by insulin (38). If confirmed, this mechanism may exacerbate the ‘relative corticosterone deficiency’ apparent with metabolic syndrome in EHERTS, and thereby exacerbate HPA axis activation in type 2 diabetes.

Given the limitations of morning plasma cortisol and corticosterone, we investigated the hypothesis of relative corticosterone deficiency in another cohort, the SKIPOGH study, in which integrated measurements of cortisol and corticosterone metabolite excretion in urine were available in a large number of participants. These results confirmed that the pattern observed for plasma glucocorticoids after ACTH₁₋₂₄ stimulation in EHERTS, with relatively low corticosterone and high cortisol, is similarly associated in urine with glucose and lipid metabolism and, strikingly, with obesity. Surprisingly, ‘relative corticosterone deficiency’ was also associated with higher systolic clinic blood pressure, although previous studies in this cohort have suggested that ambulatory blood pressure is inversely associated with *CYP17A1* activity only in the presence of high sodium intake (8).

An important limitation of these studies is the cross-sectional design which precludes inferences of causation. Moreover, genotype data was not available for EHERTS subjects because samples of DNA from this cohort have been exhausted. Indeed, no sample of sufficient size exists in which ACTH stimulation tests have been conducted to be adequately powered to detect associations of steroidogenic response with *CYP17A1* genotype. Also, effects of intensive lipid-lowering and antihypertensive treatments in the ET2DS may have obscured associations with corticosterone and cortisol. A further limitation is that the dynamic HPA axis tests and urinary steroid analyses were conducted in different

cohorts from the measurement of fasting plasma glucocorticoids. The corollary is, however, that we replicated associations of candidate *CYP17A1* genotype with phenotype in three independent studies and we have been drawn upon comprehensive datasets to adjust for a number of potentially confounding variables and assess the corticosterone:cortisol relationship in urine as well as plasma. We present our findings unadjusted for multiple testing because, as discussed above, there are distinct associations of glucocorticoid levels with individual components of metabolic syndrome such that each analysis tests a distinct hypothesis.

In conclusion, elevated morning plasma corticosterone accompanies insulin resistance and elevated cortisol in metabolic syndrome, consistent with activation of the HPA axis at the time of blood sampling. However, in addition, there are marked discrepancies between associations with ACTH-stimulated cortisol and corticosterone with metabolic syndrome which likely reflect genetically determined differences in adrenal steroidogenesis, particularly by 17 α -hydroxylase, and may also be influenced by dysregulated insulin signaling. When sustained HPA axis activation is assessed by urinary metabolites, the variation in adrenal steroidogenesis predominates in the associations with metabolic syndrome and, in particular, obesity. These findings throw the spotlight on corticosterone and suggest that further dissection of its biology in humans may be fruitful in understanding the basis for altered glucocorticoid signaling in metabolic syndrome.

ACKNOWLEDGEMENTS

We are grateful to: Lynne Ramage and Jill Harrison for technical support; Stela McLachlan for data processing in ET2DS; the study nurses Marie-Odile Levy, Guler Gök-Sogüt, Ulla Schüpbach, Dominique Siminski and Sandrine Estoppey for the data collection in SKIPOGH; and all of the participants and researchers involved. We also acknowledge the investigators in the Swiss Kidney Project on Genes in Hypertension (SKIPOGH) team: (in alphabetical order) Daniel Ackermann¹, Murielle Bochud², Michel Burnier³, Georg Ehret⁴, Idris Guessous⁵, Pierre-Yves Martin⁶, Markus Mohaupt¹, Fred Paccaud², Antoinette Pechère-Bertschi⁷, Belen Ponte⁶, Menno Pruijm³, and Bruno Vogt¹.

¹University Clinic for Nephrology, Hypertension and Clinical Pharmacology, Inselspital, Bern University Hospital, University of Bern, Switzerland

²Institute of Social and Preventive Medicine (IUMSP), Lausanne University Hospital and University of Lausanne, Switzerland

³Department of Nephrology, Lausanne University Hospital, Lausanne, Switzerland.

⁴Cardiology, Department of Specialties of Internal Medicine, Geneva University Hospitals, Geneva, Switzerland.

⁵Unit of Population Epidemiology, Department of Community Medicine and Primary Care and Emergency Medicine, Geneva University Hospitals, Geneva, Switzerland.

⁶Department of Nephrology, Geneva University Hospitals, Geneva, Switzerland.

⁷Unit of Hypertension, Departments of Community Medicine and Primary Care and Emergency Medicine, Geneva University Hospitals, Switzerland.

409

REFERENCES

- 410 1. Ehret GB, Munroe PB, Rice KM. Genetic variants in novel pathways influence blood
411 pressure and cardiovascular disease risk. *Nature*. 2011;478(7367 SRC - BaiduScholar):103-9.
- 412 2. Kato N, Takeuchi F, Tabara Y, Kelly TN, Go MJ, Sim X, et al. Meta-analysis of genome-
413 wide association studies identifies common variants associated with blood pressure variation in east
414 Asians. *Nature genetics*. 2011;43(6):531-8.
- 415 3. Fedorowski A, Franceschini N, Brody J, Liu C, Verwoert GC, Boerwinkle E, et al.
416 Orthostatic hypotension and novel blood pressure-associated gene variants: Genetics of Postural
417 Hemodynamics (GPH) Consortium. *Eur Heart J*. 2012;33(18):2331-41.
- 418 4. Newton-Cheh C, Johnson T, Gateva V, Tobin MD, Bochud M, Coin L, et al. Genome-wide
419 association study identifies eight loci associated with blood pressure. *Nature genetics*.
420 2009;41(6):666-76.
- 421 5. Schunkert H, König IR, Kathiresan S, Reilly MP, Assimes TL, Holm H, et al. Large-scale
422 association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nature genetics*.
423 2011;43(4):333-8.
- 424 6. Dai C-F, Xie X, Yang Y-N, Li X-M, Zheng Y-Y, Fu Z-Y, et al. Relationship between
425 CYP17A1 genetic polymorphism and coronary artery disease in a Chinese Han population.
426 2015;14(1):16.
- 427 7. Yanase T, Simpson ER, Waterman MR. 17 alpha-hydroxylase/17,20-lyase deficiency: from
428 clinical investigation to molecular definition. *Endocr Rev*. 1991;12(1):91-108.
- 429 8. Ackermann D, Pruijm M, Ponte B, Guessous I, Ehret G, Escher G, et al. CYP17A1 Enzyme
430 Activity Is Linked to Ambulatory Blood Pressure in a Family-Based Population Study. *Am J*
431 *Hypertens*. 2016;29(4):484-93.
- 432 9. Yan H, Guo Y, Yang T-L, Zhao L-J, Deng H-WJG, GMR mr. A family-based association
433 study identified CYP17 as a candidate gene for obesity susceptibility in Caucasians. 2012;11(3):1967.
- 434 10. Hotta K, Kitamoto A, Kitamoto T, Mizusawa S, Teranishi H, Matsuo T, et al. Genetic
435 variations in the CYP17A1 and NT5C2 genes are associated with a reduction in visceral and
436 subcutaneous fat areas in Japanese women. *J Hum Genet*. 2012;57(1):46-51.
- 437 11. Wen W, Zheng W, Okada Y, Takeuchi F, Tabara Y, Hwang J-Y, et al. Meta-analysis of
438 genome-wide association studies in East Asian-ancestry populations identifies four new loci for body
439 mass index. *Hum Mol Genet*. 2014;23(20):5492-504.

12. Echiburú B, Pérez-Bravo F, Maliqueo M, Sánchez F, Crisosto N, Sir-Petermann TJM. Polymorphism T→C (– 34 base pairs) of gene CYP17 promoter in women with polycystic ovary syndrome is associated with increased body weight and insulin resistance: a preliminary study. 2008;57(12):1765-71.
13. Walker BR. Glucocorticoids and cardiovascular disease. European journal of endocrinology. 2007;157(5):545-59.
14. Raubenheimer PJ, Young EA, Andrew R, Seckl JR. The role of corticosterone in human hypothalamic-pituitary-adrenal axis feedback. Clin Endocrinol (Oxf). 2006;65(1):22-6.
15. Karssen AM, Meijer OC, van der Sandt IC, Lucassen PJ, de Lange EC, de Boer AG, et al. Multidrug resistance P-glycoprotein hampers the access of cortisol but not of corticosterone to mouse and human brain. Endocrinology. 2001;142(6):2686-94.
16. Mattsson C, Reynolds RM, Simonyte K, Olsson T, Walker BR. Combined receptor antagonist stimulation of the hypothalamic-pituitary-adrenal axis test identifies impaired negative feedback sensitivity to cortisol in obese men. The Journal of clinical endocrinology and metabolism. 2009;94(4):1347-52.
17. McQuillan R, Leutenegger A-L, Abdel-Rahman R, Franklin CS, Pericic M, Barac-Lauc L, et al. Runs of homozygosity in European populations. Am J Hum Genet. 2008;83(3):359-72.
18. Price JF, Reynolds RM, Mitchell RJ, Williamson RM, Fowkes FGR, Deary IJ, et al. The Edinburgh Type 2 Diabetes Study: study protocol. BMC endocrine disorders. 2008;8:18.
19. Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. Diabetic medicine : a journal of the British Diabetic Association. 1998;15(7):539-53.
20. Barker DJ, Winter PD, Osmond C, Margetts B, Simmonds SJ. Weight in infancy and death from ischaemic heart disease. Lancet (London, England). 1989;2(8663):577-80.
21. Reynolds RM, Walker BR, Syddall HE, Andrew R, Wood PJ, Phillips DI. Is there a gender difference in the associations of birthweight and adult hypothalamic–pituitary–adrenal axis activity? European Journal of Endocrinology. 2005;152(2):249-53.
22. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. Diabetologia. 1985;28(7):412-9.

23. Moore A, Aitken R, Burke C, Gaskell S, Groom G, Holder G, et al. Cortisol assays: guidelines for the provision of a clinical biochemistry service. *Ann Clin Biochem.* 1985;22 (Pt 5):435-54.
24. Al-Dujaili EAS, Mullins LJ, Bailey MA, Andrew R, Kenyon CJ. Physiological and pathophysiological applications of sensitive ELISA methods for urinary deoxycorticosterone and corticosterone in rodents. *Steroids.* 2009;74(12):938-44.
25. Flote VG, Furberg A-S, McTiernan A, Frydenberg H, Ursin G, Iversen A, et al. Gene variations in oestrogen pathways, CYP19A1, daily 17 β -estradiol and mammographic density phenotypes in premenopausal women. *Breast cancer research : BCR.* 2014;16(6):499.
26. Iversen A, Thune I, McTiernan A, Makar KW, Wilsgaard T, Ellison PT, et al. Genetic polymorphism CYP17 rs2486758 and metabolic risk factors predict daily salivary 17 β -estradiol concentration in healthy premenopausal Norwegian women. The EBBA-I study. *The Journal of clinical endocrinology and metabolism.* 2012;97(5):E852-7.
27. Diver LA, MacKenzie SM, Fraser R, McManus F, Freel EM, Alvarez-Madrado S, et al. Common Polymorphisms at the CYP17A1 Locus Associate With Steroid Phenotype: Support for Blood Pressure Genome-Wide Association Study Signals at This Locus. *Hypertension.* 2016;67(4):724-32.
28. Lottenberg SA, Giannella-Neto D, Derendorf H, Rocha M, Bosco A, Carvalho SV, et al. Effect of fat distribution on the pharmacokinetics of cortisol in obesity. *Int J Clin Pharmacol Ther.* 1998;36(9):501-5.
29. Reynolds RM, Fischbacher C, Bhopal R, Byrne CD, White M, Unwin N, et al. Differences in cortisol concentrations in South Asian and European men living in the United Kingdom. *Clin Endocrinol (Oxf).* 2006;64(5):530-4.
30. Aulchenko YS, Ripke S, Isaacs A, van Duijn CM. GenABEL: an R library for genome-wide association analysis. *Bioinformatics.* 2007;23(10):1294-6.
31. Phillips D, Barker D, Fall C, Whorwood C, Seckl J, Wood P, et al. Elevated plasma cortisol concentrations: an explanation for the relationship between low birthweight and adult cardiovascular risk factors. *J Clin Endocrinol Metab.* 1998;83(757):60.
32. Reynolds RM, Labad J, Strachan MW, Braun A, Fowkes FGR, Lee AJ, et al. Elevated fasting plasma cortisol is associated with ischemic heart disease and its risk factors in people with type 2 diabetes: the Edinburgh type 2 diabetes study. 2010;95(4):1602-8.
33. Ward AMV, Fall CHD, Stein CE, Kumaran K, Veena SR, Wood PJ, et al. Cortisol and the metabolic syndrome in South Asians. *Clin Endocrinol (Oxf).* 2003;58(4):500-5.

34. Reynolds RM, Walker BR, Syddall HE, Whorwood CB, Wood PJ, Phillips DI. Elevated plasma cortisol in glucose-intolerant men: differences in responses to glucose and habituation to venepuncture. *The Journal of clinical endocrinology and metabolism*. 2001;86(3):1149-53.
35. Reynolds RM, Walker BR, Syddall HE, Andrew R, Wood PJ, Whorwood CB, et al. Altered Control of Cortisol Secretion in Adult Men with Low Birth Weight and Cardiovascular Risk Factors. *The Journal of Clinical Endocrinology & Metabolism*. 2001;86(1):245-50.
36. Soro A, Ingram MC, Tonolo G, Glorioso N, Fraser R. Mildly raised corticosterone excretion rates in patients with essential hypertension. *J Hum Hypertens*. 1995;9(6):391-3.
37. Andrew R, Phillips DI, Walker BR. Obesity and gender influence cortisol secretion and metabolism in man. *The Journal of clinical endocrinology and metabolism*. 1998;83(5):1806-9.
38. Baptiste CG, Battista M-C, Trottier A, Baillargeon J-P. Insulin and hyperandrogenism in women with polycystic ovary syndrome. *The Journal of steroid biochemistry and molecular biology*. 2010;122(1-3):42-52.
39. Varma VK, Rushing JT, Ettinger WH. High density lipoprotein cholesterol is associated with serum cortisol in older people. *J Am Geriatr Soc*. 1995;43(12):1345-9.
40. Macfarlane DP, Forbes S, Walker BR. Glucocorticoids and fatty acid metabolism in humans: fuelling fat redistribution in the metabolic syndrome. *The Journal of endocrinology*. 2008;197(2):189-204.

Figure Legends

Figure 1. Relationships between plasma cortisol and corticosterone

A) Plasma cortisol and corticosterone in the Orkney complex diseases study group (ORCADES, open symbols) and the Edinburgh Type 2 Diabetes Study (ET2DS, closed symbols). Lines indicate simple linear regression with 95% confidence bands. Coefficient of determination (r^2): ORCADES = 0.062, $p < 0.001$; ET2DS = 0.070, $p < 0.001$.

B) Cortisol and corticosterone in the East Hertfordshire Cohort. Sampling was undertaken following overnight dexamethasone (0.25mg) suppression testing (pre synacthen, open symbols); and 30 minutes following intravenous injection of ACTH₁₋₂₄ (synacthen 1µg, closed symbols). Lines indicate simple linear regression with 95% confidence bands. Coefficient of determination (r^2): pre-synacthen = 0.488, $p < 0.001$; post synacthen = 0.168, $p < 0.001$.

C and D) Total urinary cortisol and corticosterone metabolites in the Swiss Kidney Project on Genes in Hypertension. Sampling was undertaken for day (C, open symbols) and night (D, closed symbols) separately. Lines indicate simple linear regression with 95% confidence bands. Coefficient of determination (r^2): day = 0.641, $p < 0.001$; night = 0.69, $p < 0.001$.

Figure 2. Associations of plasma cortisol and corticosterone with features of metabolic syndrome

Data in A-E are for plasma cortisol (solid bars) and corticosterone (striped bars) from A) Orkney complex diseases study (ORCADES), B) Viking Health Study Shetland (VIKING), C) Edinburgh Type 2 Diabetes Study (ET2DS), D) East Hertfordshire study (EHERTS) after overnight dexamethasone (250 µg) suppression, and E) EHERTS 30 mins after ACTH₁₋₂₄ (1µg). The plots show the SD change (and standard error) in log transformed outcome for a 1SD higher log transformed cortisol or corticosterone.

Figure 3. Associations of urine cortisol and corticosterone metabolites with features of metabolic syndrome

Data are from 888 participants in the Swiss Kidney Project on Genes in Hypertension (SKIPOGH). Results are from multiple regression analysis of daytime log-transformed ratio (THA+THB+5αTHB)/(THE+THF+5αTHF) with BMI (left panel) and plasma LDL cholesterol (right panel). The model was adjusted for age, sex, center, lipid- and glucose-lowering and antihypertensive treatment, BMI (for LDL cholesterol only), estimated GFR, urine flow rate, urinary sodium, potassium and creatinine excretion. Results were similar for night-time corticosteroid excretion.

Table 1. Characteristics of participants

	Cohort				
	ORCADES	EHERTS	ET2DS	SKIPOGH	VIKING
Sex (M/F)	794/1224 (39/61%)	189/90 (68/32%)	476/427 (53/47%)	380/508 (43/57%)	838/1260 (40/60%)
Age (years)	53.3 ± 0.3 (range 16-91)	71.3 ± 0.2 (range 67-78)	67.9 ± 0.1 (range 60-68)	47.1 ± 0.6 (range 18-82)	49.8 ± 0.3 (range 18-92)
BMI (kg/m ²)	27.7 ± 0.1	27.1 ± 0.2	31.2 ± 0.2	24.7 ± 0.1	27.4 ± 0.1
Taking anti-hypertensive (n (%))	412 (20%)	105 (38%)	704 (79%)	124 (14%)	344 (16%)
Systolic BP (mmHg)	129.9 ± 0.4	159.3 ± 1.3	133.3 ± 0.6	116.8 ± 0.6	127.0 ± 0.4
Diastolic BP (mmHg)	75.5 ± 0.2	86.9 ± 0.7	69.2 ± 0.3	75.1 ± 0.3	74.7 ± 0.2
Plasma corticosterone at baseline (nM)	50.7 ± 1.2	20.2 ± 1.0*	22.3 ± 0.5	n/a	20.3 ± 1.4
Plasma corticosterone 30 min after ACTH ₁₋₂₄ (nM)	n/a	84.0 ± 2.5	n/a	n/a	n/a
Plasma cortisol at baseline (nM)	724 ± 7	188 ± 6*	732 ± 6	n/a	293 ± 4
Plasma cortisol 30 min after ACTH ₁₋₂₄ (nM)	n/a	460 ± 8	n/a	n/a	n/a
Lipid lowering therapy (n (%))	255 (13%)	71 (25%)	755 (84%)	34 (3.8%)	n/a
Total cholesterol (mM)	5.39 ± 0.03	6.82 ± 0.07	4.28 ± 0.03	5.1 ± 0.03	5.30 ± 0.02
LDL cholesterol (mM)	3.37 ± 0.02	4.89 ± 0.08	n/a	3.11 ± 0.03	3.33 ± 0.02
HDL cholesterol (mM)	1.50 ± 0.01	1.33 ± 0.02	1.28 ± 0.01	1.55 ± 0.01	1.52 ± 0.01
Triglycerides (mM)	1.15 ± 0.02	1.54 ± 0.05	n/a	1.02 ± 0.02	1.00 ± 0.01
Medication for diabetes (n)	47 (2.3%)	10 (3.6%)	734 (81%)	8 (0.8%)	44 (2.1%)
Oral agents			660 (74%)	7 (0.7%)	
Insulin			156 (18%)	1 (0.1%)	
Fasting plasma glucose (mM)	5.33 ± 0.02	5.99 ± 0.08	7.57 ± 0.07	5.14 ± 0.02	4.88 ± 0.02
Plasma glucose 30 min after oral glucose (mM)	n/a	9.38 ± 0.14	n/a	n/a	n/a
Plasma glucose 120 min after oral glucose (mM)	n/a	7.13 ± 0.18	n/a	n/a	n/a
Fasting plasma insulin mU/L	6.44 ± 0.10	7.18 ± 0.35	n/a	6.19 ± 0.21	8.31 ± 0.80
HOMA-IR	1.59 ± 0.34	1.96 ± 0.11	n/a	1.48 ± 0.06	1.86 ± 0.17
Daytime urinary steroids					
Tetrahydrodehydrocorticosterone (µg/day)	n/a	n/a	n/a	67.3 ± 1.3	n/a
Tetrahydrocorticosterone (µg/day)	n/a	n/a	n/a	91.7 ± 1.7	n/a

5 α -Tetrahydrocorticosterone (μ g/day)	n/a	n/a	n/a	188.5 $\pm$ 4.2	n/a
Tetrahydrocortisol (μ g/day)	n/a	n/a	n/a	1080.5 $\pm$ 18.2	n/a
5 α -Tetrahydrocortisol (μ g/day)	n/a	n/a	n/a	779.9 $\pm$ 20.8	n/a
Tetrahydrocortisone (μ g/day)	n/a	n/a	n/a	1861.2 $\pm$ 33.8	n/a
Nighttime urinary steroids					
Tetrahydrodehydrocorticosterone (μ g/night)	n/a	n/a	n/a	27.3 $\pm$ 0.7	n/a
Tetrahydrocorticosterone (μ g/night)	n/a	n/a	n/a	41.8 $\pm$ 1	n/a
5 α -Tetrahydrocorticosterone (μ g/night)	n/a	n/a	n/a	60.7 $\pm$ 1.7	n/a
Tetrahydrocortisol (μ g/night)	n/a	n/a	n/a	362.9 $\pm$ 8.2	n/a
5 α -Tetrahydrocortisol (μ g/night)	n/a	n/a	n/a	247.1 $\pm$ 6.8	n/a
Tetrahydrocortisone (μ g/night)	n/a	n/a	n/a	642.6 $\pm$ 14.2	n/a

Data presented as mean $\pm$ SEM. n/a = not available.

* Dexamethasone suppressed in EHRTS cohort

Table 2. Associations of functionally significant SNPs in *CYP17A1* with morning plasma cortisol and corticosterone

Study	SNP ID	Minor allele frequency	Cortisol		Corticosterone		Corticosterone:cortisol ratio	
			β	P	β	p	β	p
ORCADES	rs2486758	0.16	0.000	0.98	0.074	0.01	0.070	0.01
	rs138009835	0.09	0.003	0.89	0.004	0.91	0.003	0.93
	rs2150927	0.36	-0.008	0.52	0.023	0.29	0.032	0.14
ET2D	rs2486758	0.21	-0.004	0.83	0.074	0.09	0.077	0.05
	rs138009835	0.08	0.006	0.82	0.004	0.95	-0.002	0.97
	rs2150927	0.36	-0.005	0.70	-0.025	0.49	-0.019	0.56
VIKING	rs2486758	0.21	-0.003	0.85	-0.009	0.77	-0.004	0.88
	rs138009835	0.09	0.006	0.78	-0.024	0.60	0.040	0.26
	rs2150927	0.35	0.012	0.35	0.019	0.51	-0.012	0.58
SKIPOGH (day)	rs2486758	0.23	-133.5	0.17	6.86	0.60	0.004	0.03
	rs138009835	0.10	-119.7	0.37	-27.58	0.14	-0.004	0.22
	rs2150927	0.42	20.71	0.80	4.66	0.68	0.000	0.90
SKIPOGH (night)	rs2486758	0.23	11.09	0.80	11.03	0.05	0.004	0.05
	rs138009835	0.10	-29.08	0.64	-9.60	0.24	-0.006	0.06
	rs2150927	0.42	-0.04	0.99	-1.71	0.73	0.001	0.61

β = regression coefficient. Analyses were adjusted for age, sex, BMI, time of sampling, and relatedness.

Table 3. Associations of plasma cortisol and corticosterone with features of the metabolic syndrome

		Cortisol				Corticosterone				Corticosterone:cortisol ratio			
		Unadjusted		Adjusted		Unadjusted		Adjusted		Unadjusted		Adjusted	
		β	P	β	p	β	p	B	p	β	p	β	p
ORCADES (n=2018)	BMI	-0.069	<0.001	-0.064	<0.001	-0.026	<0.001	-0.028	<0.001	-0.006	0.33	-0.009	0.11
	Systolic BP	-0.021	0.01	0.022	0.001	-0.004	0.41	0.005	0.25	0.003	0.48	-0.002	0.57
	Total cholesterol	-0.006	0.61	0.021	0.09	-0.002	0.73	-0.002	0.73	-0.010	0.17	-0.008	0.27
	LDL cholesterol	-0.042	0.02	0.006	0.74	-0.030	0.004	-0.018	0.08	-0.021	0.06	-0.018	0.09
	HDL cholesterol	0.063	<0.001	0.022	0.14	0.026	0.004	0.021	0.02	0.007	0.45	0.014	0.12
	Triglycerides	0.047	0.12	0.176	<0.001	0.016	0.35	0.057	0.001	0.012	0.52	0.010	0.56
	Fasting glucose	-0.007	0.31	0.014	0.02	0.011	0.003	0.014	<0.001	0.015	<0.001	0.011	0.004
	Insulin	-0.066	0.03	0.030	0.27	0.009	0.58	0.042	0.007	0.038	0.03	0.038	0.02
	HOMA-IR	-0.073	0.03	0.045	0.13	0.020	0.27	0.057	0.001	0.053	0.01	0.048	0.01
VIKING (n=2093)	BMI	-0.047	<0.001	-0.035	<0.001	-0.018	<0.001	-0.014	<0.001	-0.012	0.001	-0.010	0.004
	Systolic BP	-0.009	0.11	0.020	<0.001	-0.003	0.30	0.004	0.04	-0.002	0.49	0.001	0.80
	Total cholesterol	-0.019	0.03	-0.001	0.89	-0.006	0.11	-0.002	0.64	-0.005	0.27	-0.003	0.46
	LDL cholesterol	-0.048	<0.001	-0.021	0.09	-0.013	0.02	-0.005	0.35	-0.006	0.32	-0.002	0.69
	HDL cholesterol	0.035	0.002	0.013	0.21	0.014	0.004	0.003	0.41	0.007	0.18	-0.001	0.91
	Triglycerides	0.017	0.48	0.112	<0.001	-0.015	0.13	0.014	0.13	-0.022	0.04	-0.006	0.59
	Fasting glucose	-0.006	0.22	0.012	0.008	0.001	0.72	0.006	0.002	0.002	0.32	0.005	0.02
	Insulin	-0.041	0.10	0.037	0.09	-0.020	0.06	0.008	0.37	-0.017	0.15	0.001	0.96
	HOMA-IR	-0.047	0.08	0.048	0.04	-0.020	0.09	0.014	0.15	-0.015	0.25	0.005	0.62
ET2DS (n=903)	BMI	-0.064	0.003	-0.055	0.01	-0.018	0.05	-0.013	0.12	-0.010	0.30	-0.006	0.50
	Systolic BP	0.015	0.34	0.014	0.37	-0.003	0.66	-0.003	0.66	-0.005	0.40	-0.005	0.42
	Fasting glucose	0.179	<0.001	0.190	<0.001	0.027	0.03	0.030	0.02	0.001	0.95	0.003	0.84
EHERTS dex suppressed (n=279)	BMI	-0.018	0.13	-0.017	0.18	-0.036	0.003	-0.036	0.004	-0.023	0.14	-0.024	0.12
	Systolic BP	0.008	0.49	0.005	0.67	0.008	0.49	0.008	0.49	-0.002	0.89	0.000	1.00
	Fasting glucose	0.018	0.06	0.010	0.28	0.003	0.74	-0.001	0.89	-0.019	0.12	-0.012	0.30
	30 min glucose	0.033	0.05	0.027	0.12	0.015	0.39	0.015	0.38	-0.027	0.23	-0.014	0.51
	120 min glucose	-0.049	0.04	-0.034	0.17	-0.050	0.04	-0.032	0.20	0.016	0.61	0.012	0.69
	Insulin	-0.138	0.01	-0.096	0.06	-0.192	<0.001	-0.131	0.01	-0.046	0.80	-0.040	0.52
	HOMA-IR	-0.119	0.03	-0.085	0.11	-0.189	0.001	-0.133	0.01	-0.105	0.59	-0.053	0.42
EHERTS after ACTH ₁₋₂₄ (n=279)	BMI	0.012	0.71	-0.020	0.61	-0.015	0.19	-0.018	0.12	-0.019	0.11	-0.017	0.15
	Systolic BP	0.014	0.65	0.048	0.19	-0.007	0.58	-0.004	0.77	-0.014	0.32	-0.011	0.41
	Fasting glucose	-0.031	0.19	0.037	0.20	-0.033	0.003	-0.027	0.01	-0.021	0.06	-0.026	0.02
	30 min glucose	0.019	0.66	0.146	0.01	-0.023	0.25	-0.010	0.63	-0.021	0.32	-0.019	0.35
	120 min glucose	0.092	0.15	0.010	0.90	-0.054	0.06	-0.056	0.05	-0.072	0.02	-0.059	0.05
	Insulin	0.136	0.32	0.036	0.82	-0.120	0.04	-0.099	0.07	-0.046	0.02	-0.096	0.09
	HOMA-IR	0.103	0.48	0.072	0.66	-0.153	0.01	-0.126	0.03	-0.485	0.02	-0.123	0.04

β = regression coefficient. P values refer to regression co-efficient, unadjusted or adjusted in multiple regression analyses for co-variables (age, sex, time of sample, relevant medication, and BMI (unless BMI is the outcome variable)). All continuous variables were normalised by log transformation.

Table 4. Associations of day- and night-time excretion of urinary cortisol and corticosterone metabolites with features of the metabolic syndrome

		Cortisol metabolites (THE+THF+5 α THF) day				Corticosterone metabolites (THA+THB+5 α THB) day				Corticosterone:cortisol metabolites (THA+THB+5 α THB)/(THE+THF+5 α THF) day			
		Unadjusted		Adjusted		Unadjusted		Adjusted		Unadjusted		Adjusted	
		β	p	B	p	β	p	B	p	β	p	β	p
SKIPOGH (n=888)	BMI	2.351	<0.001	1.824	<0.001	0.563	0.02	0.143	0.564	-4.618	<0.001	-3.388	<0.001
	Systolic BP	5.996	<0.001	0.385	0.688	0.428	0.65	-0.631	0.449	-13.97	<0.001	-2.524	0.067
	Total cholesterol	-0.045	0.47	0.096	0.173	-0.239	<0.001	-0.032	0.6	-0.542	<0.001	-0.290	0.004
	LDL cholesterol	0.166	0.002	0.174	0.005	-0.052	0.32	0.025	0.64	-0.569	<0.001	-0.295	0.001
	HDL cholesterol	-0.240	<0.001	-0.020	0.026	-0.157	<0.001	0.003	0.908	0.190	<0.001	0.049	0.188
	Triglycerides	0.093	0.006	-0.118	0.002	-0.045	0.17	-0.120	<0.001	-0.368	<0.001	-0.085	0.123
	Fasting glucose	0.244	<0.001	0.055	0.16	0.093	0.01	0.028	0.401	-0.380	<0.001	-0.036	0.523
	Insulin	1.173	0.001	-0.562	0.163	-0.278	0.43	-0.805	0.021	-3.821	<0.001	-1.050	0.071
	HOMA-IR	0.360	0.001	-0.157	0.184	-0.036	0.73	-0.196	0.058	-1.033	<0.001	-0.212	0.217
		(THE+THF+5 α -THF) night				(THA+THB+5 α -THB) night				(THA+THB+5 α THB)/(THE+THF+5 α THF) night			
	BMI	2.094	<0.001	1.274	<0.001	0.676	0.002	-0.025	0.910	-4.252	<0.001	-3.123	<0.001
	Systolic BP	8.146	<0.001	1.444	0.111	2.765	0.002	-0.091	0.903	-13.58	<0.001	-3.152	0.023
	Total cholesterol	0.011	0.85	0.005	0.938	-0.096	0.088	-0.036	0.508	-0.331	0.001	-0.153	0.136
	LDL cholesterol	0.179	0.001	0.090	0.129	0.026	0.596	0.001	0.982	-0.423	<0.001	-0.207	0.023
	HDL cholesterol	-0.195	<0.001	-0.031	0.212	-0.095	<0.001	0.021	0.294	0.281	<0.001	0.137	<0.001
	Triglycerides	0.078	0.018	-0.114	0.002	-0.047	0.121	-0.128	<0.001	-0.420	<0.001	-0.176	0.001
	Fasting glucose	0.299	<0.001	0.093	0.011	0.170	<0.001	0.060	0.047	-0.294	<0.001	0.013	0.824
	Insulin	1.213	0.001	-0.135	0.726	0.008	0.979	-0.499	0.110	-3.850	<0.001	-1.355	0.022
	HOMA-IR	0.394	<0.001	-0.033	0.772	0.060	0.533	-0.110	0.230	-1.041	<0.001	-0.306	0.079

β = regression coefficient. P values refer to regression co-efficient, unadjusted or adjusted in multiple regression analyses for co-variables (age, sex, center, relevant medication, and BMI (unless BMI is the outcome variable)). The ratio (THA+THB+5 α THB)/(THE+THF+5 α THF) was normalised by log transformation before regression analyses.





Figure 1

A


ORCADES

bioRxiv preprint doi: <https://doi.org/10.1101/654269>; this version posted May 29, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.




B

VIKING




C

ET2DS



D

EHERTS (dex-suppressed)



E

EHERTS (post-synacthen)




Figure 2





Figure 3