

1 **Risk practices for bovine tuberculosis transmission to cattle and**
2 **livestock farming communities living at wildlife-livestock-human**
3 **interface in northern KwaZulu Natal, South Africa**

4

5 Petronillah Rudo Sichewo^{1,2*} ^{¶#a}, Catiane Vander Kelen^{3¶}, Séverine Thys^{4¶}, Anita Luise Michel^{1,5¶}

6 1. Department of Veterinary Tropical Diseases, Bovine Tuberculosis and Brucellosis Research
7 Programme, Faculty of Veterinary Sciences, University of Pretoria, South Africa.

8 2. Department of Animal Sciences, Faculty of Natural Resources Management and Agriculture,
9 Midlands State University, Zimbabwe

10 3. Department of Public Health, Neglected Tropical Diseases Unit, Institute of Tropical Medicine,
11 Belgium

12 4. Department of Virology, Parasitology, and Immunology, Faculty of Veterinary Medicine, Ghent
13 University, Belgium;

14 5. Research Associate at the National Zoological Gardens of South Africa, Pretoria, South Africa.

15

16 ^{#a} Current address: Department of Veterinary Tropical Diseases, Level 1, Paraclinical Building,
17 University of Pretoria, Private Bag X04, Onderstepoort 0110, South Africa

18 * Corresponding author

19 E-mail: psichewo@gmail.com

20 ¶ These authors contributed equally to this work.

21 Abstract

22 Bovine tuberculosis (bTB) is a disease of cattle that is transmitted through direct contact
23 with an infected animal or ingestion of contaminated food or water. This study seeks to
24 explore the local knowledge on the disease and establish the risk practices that lead to
25 its transmission to cattle and humans (zoonotic TB) in a traditional livestock farming
26 community with a history of bTB diagnosis in cattle and wildlife. Information was
27 collected using a qualitative approach of Focus Group Discussions (FGDs) targeting
28 household members of livestock farmers that owned either bTB infected or uninfected
29 herds. We conducted fourteen FGDs (150 individuals) across four dip tanks that
30 included the following categories of participants from cattle owning households: heads
31 of households, cattle keepers, dip tank committee members and women. The qualitative
32 data was managed using NVivo Version 12 Pro®software. Social and cultural practices
33 were identified as major risky practices for bTB transmission to people, such as the
34 consumption of undercooked meat, consumption of soured /raw milk and lack of
35 protective measures during slaughtering of cattle. The acceptance of animals into a
36 herd without bTB pre-movement testing following traditional practices (e.g. *lobola*, 'bride
37 price', the temporary introduction of a bull for 'breeding'), the sharing of grazing and
38 watering points amongst the herds and with wildlife were identified as risky practices for
39 bTB transmission to cattle. Overall, knowledge of bTB in cattle and modes of
40 transmission to people and livestock was found to be high. However, the community
41 was still involved in risky practices that expose people and cattle to bovine TB. An inter-
42 disciplinary 'One Health' approach that engages the community is recommended, to
43 provide locally relevant interventions that allows the community to keep their traditional

44 practices and socio-economic systems whilst avoiding disease transmission to cattle
45 and people.

46 **Author summary**

47 Bovine tuberculosis (bTB) is a respiratory disease of cattle that is transmitted to
48 other animals as well as humans (zoonotic TB) through direct contact with infected
49 animals, and consumption of contaminated food (animal products) or water. The study
50 explains the complexities of human-animal relations, reflects on how people understand
51 and conceptualize risk of bovine tuberculosis (bTB) in an endemic area considering the
52 economic value of livestock keeping as well as social and cultural practices of
53 importance to the community. The results of this study identified socio-cultural practices
54 that involved consumption of raw or undercooked animal products and handling of
55 infected animal products during animal slaughter as major risky practices for bTB
56 transmission to people. Introduction of animals into a herd without bTB testing for socio-
57 cultural purposes and sharing of resources amongst the communal herd and with
58 wildlife were identified as risky practices for bTB transmission to cattle. The findings of
59 this study illustrate the need for a One Health strategy that develops appropriate public
60 health policy and related education campaigns for the community as control of zoonotic
61 TB in people depends on the successful control of bovine TB in cattle.

62

63

64

65 Introduction

66 Bovine tuberculosis (bTB) is chiefly a chronic respiratory disease of cattle caused
67 by the bacteria *Mycobacterium bovis* (*M. bovis*) that has multiple incidental hosts
68 including humans, goats, cats, dogs and wild animals [1–3]. In South Africa, the disease
69 is a state controlled disease in cattle due to its negative impact on livestock production,
70 export and local market in animal and animal products, wildlife conservation efforts and
71 could increase human health costs [4,5]. Similar to most African countries where bTB is
72 prevalent, the surveillance and ‘test and slaughter’ programs are not optimally
73 implemented in South Africa due to a lack of resources, despite the potential hazard to
74 human health [6,7]. *M. bovis* infection in humans is referred to as zoonotic tuberculosis
75 and has been classified by the World Health Organisation as a neglected zoonotic
76 disease [8]. Zoonotic TB transmission to humans is predominantly through the
77 consumption of contaminated animal products such as unpasteurised dairy products
78 and less frequently attributed to animal-to-human or human-to-human through direct
79 contact [9].

80 The study area is surrounded by conservation areas where *M. bovis* infection has
81 been established in African buffalo which is wildlife maintenance host [10]. Recent
82 studies have revealed *M. bovis* infection in communal cattle in the same area with 28%
83 of the farmers having at least one test positive animal in their herd [11]. The presence of
84 *M. bovis* in cattle and wildlife increases the risk of zoonotic TB transmission to susceptible
85 human populations living at the wildlife/livestock/human interface [5,12]. A lack of or
86 insufficient implementation of the ‘test and slaughter’ disease control scheme,
87 consumption of uncooked meat products and soured milk, poor understanding of zoonotic

88 TB and poor sanitary conditions are some of the potential risk factors for *M. bovis* infection
89 and disease in humans [13].

90 Due to an increase in land use there is infringement of human activities into
91 conservation areas that results in the sharing of natural resources with wildlife [14].
92 Studies in other countries have suggested that transmission of bTB to cattle from wildlife
93 occurs either through direct contact at shared resources such as watering points or
94 indirectly, when cattle graze on contaminated pastures [15]. Therefore, the wildlife-
95 livestock interface has been defined as a high-risk area for bTB transmission from wildlife
96 to cattle [16]. It is likely that local farming practices will impact on the bTB prevalence
97 therefore it is important to identify local risk factors to *M. bovis* transmission [17].

98 Despite reports of isolation of *M. bovis* from livestock and wildlife, limited
99 information is available in South Africa on the level of bTB knowledge and risk practices
100 of livestock farming communities that influence the transmission of bTB to cattle and
101 humans. The understanding of these practices will provide information for the
102 development of informed grassroots programs that integrate the local and scientific
103 knowledge towards bTB control in animal and zoonotic TB control in human populations
104 living at the wildlife-livestock-human interface. The present study was therefore designed
105 to understand the local knowledge on bTB among livestock farming communities and
106 investigate the risk practices that were associated with bTB infection in cattle and in
107 people co-existing with wildlife.

108

109

110 **Methodology**

111 **Study area and population**

112 A qualitative study based on focus group discussions (FGDs) was carried out in
113 Big Five False Bay Municipality, uMkhanyakude District in the northern part of KwaZulu-
114 Natal, South Africa. This community was part of a One Health investigation into the
115 epidemiology of bTB at the wildlife-livestock-human interface. The Municipality is sparsely
116 populated, and most of the population that occupy the north-eastern part are rural
117 traditional communities with a cattle population that was estimated at 11 000 from a total
118 of 456 owners (W. McCall personal Communication, 28 August 2017). The population is
119 engaged primarily in crop- livestock farming and the main domestic animals are cattle
120 and goats. The Municipality is surrounded by game and nature reserves such as St. Lucia
121 (iSimangaliso), Hluhluwe/Imfolozi, Munyawana and Mkuze, which attract a high number
122 of local and international tourists.

123 The study participants were purposively selected according to the inclusive criteria
124 of being a household member of a farmer owning cattle that were tested for bTB in
125 September 2016/ March 2017 at one of the four dip tanks in the area (Masakeni, Mpempe,
126 Nibela and Nkomo) and owning either a bTB positive or negative herd [11]. A dip tank is
127 a communal cattle handling facility where animals from several villages assemble weekly
128 or once per fortnight for disease inspection and are also dipped in an insecticide plunge
129 tank for external parasites control, primarily ticks.

130 The four-focus group categories that were selected from each dip tank included
131 women that belong to households owning cattle, cattle keepers (male and female), heads
132 of households and dip tank committee members. The group of adult female members of

133 households was selected because in this socio-cultural context women are often solely
134 responsible for the handling of food and food preparation for their families hence
135 determining their consumption behavior. Cattle keepers were included because they are
136 responsible for taking care of the animals (animal husbandry practices). These include
137 young boys that are employees or members of the household (male/female). The heads
138 of households are usually the decision-making group within the household that determine
139 the movement, introduction and selection of animals for slaughter. Finally, the dip tank
140 committee is a group of farmers selected by the community for each dip tank to assist the
141 government animal health technicians in the management of dipping activities and
142 reporting of animal diseases.

143

144 **Data collection**

145 Fourteen focus group discussions (FGDs) were carried out in November and
146 December 2017. In each FGD we examined the knowledge, awareness of bovine and
147 zoonotic tuberculosis and the risk practices for bTB transmission to cattle, humans and
148 wildlife among communal (rural) cattle farmers. A FGD guide was used and pretested in
149 one FGD at Masakeni dip tank that consisted of dip tank committee members and the
150 final topics examined include:

151 i) Local knowledge and awareness of bTB in terms of symptoms, transmission and control
152 measures

153 ii) Food handling, preparation and consumption behaviors, cattle slaughtering procedures

154 iii) Livestock management practices in terms of cattle and wildlife interactions, cattle and
155 cattle interactions and introduction of cattle into a herd.

156 A total of 150 people above 16 years were recruited by the local Animal Health
157 technician into the study based on their various roles in the household and participation
158 in the FGDs depended on their availability and or willingness. The participants for each
159 group belonged to different households within the same dip tank, had similar socio-
160 characteristics and were comfortable to discuss issues among themselves and the
161 facilitator. FGDs were carried out at community halls that were centrally located and
162 where village meetings or activities were commonly conducted.

163 The FGDs consisted of a facilitator, two observers and the preselected group of
164 participants from the target community. The facilitator (TM), the first observer/note taker
165 (PZ), and the principal investigator (PS) who was also a second observer in the discussion
166 were trained in focus group interview principles and techniques by a qualified social
167 science researcher (CV) from the Institute of Tropical Medicine, Belgium. The discussions
168 lasted for about one hour, used the local language of isiZulu and were steered in a flexible
169 and iterative manner. All the FGDs were led by the trained facilitator and the two
170 observers, audio-recorded by digital voice recorder with the permission of the
171 participants, transcribed and translated from local language into English. The facilitator
172 and the observers were fluent in both English and isiZulu.

173

174 **Data management and analysis**

175 All the discussions were transcribed into a Word document and these were cross
176 checked and supervised by the principal investigator. The transcripts were read several
177 times to get an overall understanding and to identify the main and salient themes until no
178 new themes were found. The list of themes was used to code all the transcripts including

179 the hand-written notes from the observers and the process was managed using NVivo
180 Version 12 Pro®software. We examined the relationships between themes and sub-
181 themes, patterns in the views expressed by the various groups or dip tanks in terms of
182 differences or similarities. Each theme was described in detail and exemplary quotes
183 were used to illustrate the meaning of the themes. The six main themes are represented
184 in detail in the results section with quotations from the participants.

185 **Ethical consideration**

186 The University of Pretoria-Faculty of Humanities-Research Ethics Committee
187 approved the study (Reference:16394624/GWO170814HSA). At the commencement of
188 each FGD the study was explained to the participants regarding the research purpose,
189 FGD process, confidentiality and uses of the data. They could ask questions for clarity
190 and thereafter they were given the consent form to read or it was read to them in the local
191 language. A written consent was obtained from the participants to conduct the
192 discussions and oral permission obtained to audio-record all the discussions.

193 **Results**

194 The 14 FGDs involved the following groups from each dip tank; dip tank committee
195 members, head of households, women from cattle owning households and cattle keepers
196 from the 4 dip tanks; Masakeni, Nibela, Mpempe and Nkomo as shown in Table 1.

197

198

199

200 Table 1: Description of Focus Groups according to dip tank, number of participants and
201 gender

Dip tank name	FGDs category	Number of participants	Gender
Masakeni	cattle keepers	11	male
	women	12	female
Nkomo	head of households	12	male
	cattle keepers	11	male & female
	dip tank committee	10	male
	women	12	female
Mpempe	head of households	9	male & female
	cattle keepers	12	male
	dip tank committee	10	male & female
	women	8	female
Nibela	head of household	12	male & female
	cattle keepers	12	male
	dip tank committee	10	male
	women	9	female

202
203 The results are presented according to the themes that were identified from the
204 analysis of the transcripts.

205 **Knowledge on bovine tuberculosis (bTB): symptoms, transmission and control**
206 **measures**

207 Most of the respondents indicated that they knew about bTB and considered it as
208 one of the significant diseases of cattle found in their area. All the groups were able to
209 discuss symptoms of the diseases. Knowledge on the transmission of bTB to cattle or
210 humans was evident in all FGDs except the cattle keepers that were not sure of how this
211 disease was transmitted to humans. Knowledge on bTB was shown by the ability of the
212 participants to mention the correct clinical signs.

213 *“We have seen some of these from some of our animals. The animals cough a lot, I have
214 heard the animals cough especially in the morning, the animal has difficulty in breathing
215 making a loud sound, and it loses weight and has no appetite. That is all I know I do not
216 know about the others”.* (Nibela dip tank, head of household, December 2017)

217 Most of the FGD participants were aware of possible transmission of zoonotic
218 diseases from livestock to people through contact with infected animals or consumption
219 of infected animal products, however they could not give specific examples of zoonotic
220 diseases in livestock or wildlife.

221 *“[...] Let’s say there is a little child in the yard and the child approaches the cow as he or
222 she (the child) is used to the cow and the cow is also used to this child, we might not see
223 it at the time that the cow has a sickness. The child playing next to the cow can inhale
224 whatever the cow coughs”.* (Masakeni dip tank, female member of household, December
225 2017)

226 The discussants from all FGDs groups, except the ones that included the women
227 from cattle owning households, highlighted that transmission from infected cattle or
228 wildlife to uninfected cattle was through direct contact or indirectly by grazing on
229 contaminated pastures.

230 *[...] The cows salivate uncontrollably and more than usual, so an infected cow can leave*
231 *traces of saliva on the grass in which other cows can graze on and then become infected*
232 *as well. TB can be transmitted because an infected cow is always coughing and lacks the*
233 *means to cover itself when it does; the fact that it does not live in isolation means other*
234 *cows and animals can easily be infected". (Mpempe dip tank, dip tank committee*
235 *member, December 2017)*

236 The participants were not conversant with bTB control methods instead mentioned
237 that animal testing was needed since clinical signs could not be used for the diagnosis of
238 the disease.

239 *"The problem with this disease that we are discussing is that it is only visible inside the*
240 *animal and not outside. My biggest fear is that we may end up killing a lot of animals*
241 *trying to figure out what is really bothering them. All I'm trying to say is that the knowledge*
242 *that we have is not enough for us to make sound decisions". (Masakeni dip tank, cattle*
243 *keeper, December 2017)*

244 **Cattle slaughtering and meat inspection**

245 The respondents from all the FGDs revealed that the selection of an animal for
246 slaughter depended on the general health of the animal as observed by the owner,

247 animal's productivity (fertility), age and purpose i.e. for general consumption or for a
248 traditional ceremony.

249 *“1. The reason for the selection of a cow is the size of the family and the number of guests
250 expected. 2. Would be the overall health of the cow [...]3. The cows are slaughtered or
251 selected for slaughtering on a chronological basis where older cows stand a better chance
252 of being selected than younger ones because of expected levels of yield from the young
253 ones”.* (Mpempe dip tank head of household, December 2017)

254 The cattle keepers revealed that they did not wear protective clothing during
255 slaughtering.

256 *“We handle the meat with these bare hands (shows hands), we don't wear gloves for this.
257 This is not a human corpse that has AIDS”.* (Mpempe dip tank cattle keeper, December
258 2017)

259 However, some respondents particularly from the dip tank committee members
260 group had a different opinion after attending past educative meetings on bovine TB.

261 *“Back when we grew up that used to be the case, but now wearing sandals during
262 slaughtering is no different than not wearing gloves. From all the meetings and
263 information sessions we attend, the advice we get is 'wear your protective clothing and
264 gloves if you have' and what is also different is that you have an animal inspected prior
265 to it being slaughtered. In the olden days even, a sickly-looking animal was not spared
266 [...]”.* (Nkomo dip tank committee member, December 2017)

267 The respondents differed in their practices concerning the management of meat
268 with spots. The cattle keepers revealed that in most cases the meat was not inspected
269 and was consumed as quickly as possible once it was slaughtered:

270 *“Ours is different from the meat you get at the butchery where it is checked for*
271 *abnormalities and illnesses. Once we have slaughtered the animal it is only correct for us*
272 *to eat its meat”*. (Nkomo dip tank cattle keeper, December 2017)

273 The cattle keepers also highlighted that the consumption of uninspected meat was
274 a common practice particularly during traditional ceremonies that could not be delayed or
275 postponed.

276 *“The general procedure is that a cow must be checked before slaughtering. This isn’t*
277 *followed because we want to eat meat during a ceremony. The way I see it, when it is*
278 *said that we should check these cows before slaughter; what would happen if we find*
279 *them to be sick and there is a huge possibility that many other are sick because they live*
280 *together. Does this mean I must reschedule a settled date for an ancestral ceremony now*
281 *that it was found that the cows are sick?”*. (Nibela dip tank cattle keepers, December
282 2017)

283 In contrast the FGDs that included cattle owners, dip tank committee members
284 and the female household members indicated that they inspected the meat but that the
285 occurrence of abnormalities was rare. In the case they were coming across unusual spots
286 on organs such as the liver or lungs, they were taking the necessary precautions. These
287 included discarding the affected organ by burying it in the ground, throwing it away or

288 sending the organ to the animal health technician for testing but at the same time they
289 proceeded to eat the rest of the meat that did not show visible spots.

290 *“We have a discussion amongst ourselves on what could possibly be wrong with the*
291 *animal. After this we remove the affected area and continue with the healthy-looking*
292 *parts”.* (Mpempe dip tank head of household, December 2017)

293 **Introduction of cattle into a herd by communal farmers**

294 During the discussions it was clear that the circumstances that led to the
295 introduction of cattle into the herds were common across all the dip tanks. These included
296 the receiving of cattle as a ‘bride price’ as part of the marriage gifts from the groom’s
297 family to the bride’s family, that is locally called “*lobola*”, when performing traditional
298 marriage ceremonies and the commercial exchange of animals.

299 *“We do it for lobola or sometimes we sell amongst each other when we need money”.*
300 (Nkomo dip, head of household, December 2017)

301 Other situations involved individuals that offered help to take care of their neighbor
302 or relative’s cattle or when animals were exchanged to obtain a special bull for ‘breeding’
303 or a specific animal for a traditional ceremony as described below:

304 *“Maybe my neighbor needs a big cow for slaughter and to compensate for this exchange*
305 *I may take two from his herd. You mentioned that there is a drought, another circumstance*
306 *would be to move the cows to where conditions are better whether it is at a friend’s farm,*
307 *neighbor or relative”.* (Masakeni dip tank cattle keeper, December 2017)

308 The animals that were used for compensation for a crime or paid as a fine were
309 not mixed with the herd but were immediately slaughtered.

310 *“No, a cow for compensation does not stay long. It is slaughtered on the very same day*
311 *and it doesn’t enter the yard because it is slaughtered at the gate on arrival. It does not*
312 *stay long because it is an insult”*. (Mpempe dip tank, female member of household,
313 December 2017)

314 **Criterion for accepting animals into the herd and the role of veterinary services**

315 The criteria of acceptance of cattle into a herd was similar as explained in all the
316 FGDs across the dip tanks and depended on the purpose of the introduction of the
317 animals into the herd. For instance, cattle that were brought to the family to pay the bride
318 price were not rejected due to social and cultural reasons.

319 *“Because everyone is happy during this time (Lobola and ultimately marriage), no one*
320 *takes the time to thoroughly inspect the animals with a sober mind because of the*
321 *happiness that the family may be feeling at the time. In a nutshell we do accept such cows*
322 *knowingly or unknowingly”*. (Nibela dip tank cattle owner, December 2017)

323 *“[...] It is tradition, we live according to ancient rules and customs. When your prospective*
324 *son-in-law comes to pay lobola and you refuse his cows you are guilty of a crime in this*
325 *whole exchange. He (son-in-law) will never again be expected to pay Lobola because*
326 *you refused to accept his cows initially. Customary law also agrees if he takes your*
327 *daughter and goes on to live with her for free. He has the grounds and can state this at*
328 *the tribal council that you refused to accept his cows when he was then willing. That is*
329 *the primary reason why we accept cows. You are in part afraid of giving your son-in-law*
330 *powers to walk away freely with your daughter while also demeaning your daughter’s*
331 *character”*. (Mpempe dip tank cattle owner, December 2017)

332 The same principle of accepting the cattle without concern for their health status
333 was also applied in circumstances whereby the transfer of animal was meant to help a
334 neighbor or relative with animal care.

335 *“Another circumstance would be the exchange of cattle between owners when one has*
336 *better grazing land and water conditions than the other. And during such an exchange it*
337 *might be that my herd is infected and moving into uninfected herd and vice versa”.* (Nibela
338 dip tank women, December 2017)

339 In response to the commercial exchange of animals all the groups concurred that
340 in this scenario it was within their right to accept or reject an animal after inspecting its
341 health status and if possible, involve their local state veterinary officers to grant the
342 permission for movement of animals into the area.

343 *“What is better is the buying of cattle because I cannot buy something that does not satisfy*
344 *me. But a cow that is to be given to me by someone else or one that needs help because*
345 *of the drought, I would not turn back such person and his cow”.* (Masakeni dip tank head
346 of household, November 2017)

347 **Cattle-to-cattle and cattle-to-wildlife interactions**

348 Regardless of the group category, participants acknowledged that cattle-to-cattle
349 contact from different herds was common in the area when their animals gather at the dip
350 tank, communal watering points and nearby pastures.

351 *“Yes, our cattle interact with other cattle from herds different from ours. The interaction*
352 *happens especially at grazing points and at the dip tank”.* (Nibela dip tank, head of
353 household, December 2017)

354 Contact with wildlife was also noted when farmers were granted permission by the
355 game park authorities to move their cattle into the game park for water and grazing. This
356 information was provided by the respondents from FGDs that were mainly involved in the
357 herding of animals i. e the cattle keepers, cattle owners and dip tank committee members.
358 This situation was more generally the case of farmers from Nibela dip tank in our study.

359 *“There is a nearby wetland and a game park called iSimangaliso. Because of the*
360 *difficulties posed by the drought, we have arranged with the park’s management to allow*
361 *us to enter and let our cows graze and drink water within the park. And you would find*
362 *our cows interacting with wildlife, especially buffalos, zebras and wildebeest drinking and*
363 *grazing together”*. (Nibela dip tank committee members, December 2017)

364 Cattle to wildlife contact was also occurring through the broken-down fences
365 surrounding some game parks in the area, allowing free movement of animals into and
366 out of the game park.

367 *“The fence around the reserve is not properly erected to keep livestock out of the reserve*
368 *and wild animals in. So, cattle love walking with zebras and with buffalos as well. This is*
369 *even though buffalos are dangerous to the cow because they will kill it. Nevertheless,*
370 *they do interact a lot”*. (Nkomo dip cattle keeper, December 2017)

371 However, one of the dip tanks (Masakeni) indicated that the fence that surrounded
372 the game park in their area was not damaged and movement of animals into the game
373 park was impossible.

374 “In this area there is no wildlife-livestock interactions. No, they do not because wild
375 animals from the game parks or lodges are well fenced in. It can be the cows and goats
376 that interact with each other”. (Masakeni dip tank cattle keepers, December 2017)

377 **Food preparation and consumption practices of communal farmers**

378 Reference to the preparation and consumption of animal products for food,
379 medicine and traditional practices was discussed with various groups.

380

381 Milk and meat were clearly stated as the main products that are obtained from the cattle.
382 “It has to be meat, milk and amasi (soured milk)”. (Mpempe dip tank women, December
383 2017)

384 Other animal products mentioned included cattle hide, horns, bile (for seasoning
385 of meat and traditional rituals), gall bladder (for seasoning of meat and cultural wrist
386 bands), urine (as medicine for a bad cough), cow dung (to neutralize poison) and blood
387 which was mostly used following cultural practices.

388 “Traditional healers and their trainees drink it (blood). Others drink the bile from the cow”.
389 (Nkomo dip tank women, December 2017)

390 The meat was prepared in various ways as illustrated by the responses given and
391 whereby the meat was either boiled or grilled (“braaing”).

392 “My child, the meat is cut into different pieces, the portions that need grilling are cut out
393 and portions that require to be boiled are also cut out. We leave it to boil in the pot till it is
394 soft”. (Masakeni dip tank women, December 2017)

395 The men, that mostly take part in the slaughtering of the cow, chose the meat
396 portions for their *braaiing* or special stew and this determined the success of a traditional
397 ceremony. Afterwards the rest of the meat is cooked by women for the whole family.

398 “*Methods of preparation vary with the kind of meat that is to be consumed. The norm is*
399 *that there is meat called ‘stolen’ meat. What I mean by this is that very small portions*
400 *from the entire cow’s carcass are skillfully cut off to make a stew of all the portions cut off*
401 *and the other half is flame-grilled (braai)*”. (Mpempe dip tank committee, December 2017)

402 When the participants were asked about their view on the consumption of
403 undercooked meat the majority responded that it was the best and most preferred
404 preparation, especially the liver, as expressed by a female participant:

405 “*Braai meat shouldn’t be overcooked my child, cooked so much that it becomes dry. The*
406 *same applies to the liver. It must not dry up; it needs to have that ounce of blood on it*
407 *because when it is well-done it becomes very hard*”. (Nkomo dip tank women, December
408 2017)

409 It was also clear that raw liver was consumed because they believed that it
410 supplied nutrients.

411 “*Like those that eat a cow’s liver raw. Yes, raw. It is believed that there are more nutrients*
412 *in it at this state than when it is cooked*”. (Mpempe dip tank committee member, December
413 2017)

414 Participants revealed that consumption of raw meat from the other organs of the
415 cow was also done despite them knowing that this practice placed them at a risk for
416 zoonotic diseases.

417 *“This would mean that the meat should not be eaten. The chances of us adhering to this*
418 *condition are very minimal, because when we slaughter there is meat in which we*
419 *consume sometimes raw or half-cooked. There are other parts in the cow’s intestine that*
420 *are eaten raw”.* (Nibela dip tank, cattle keepers, December 2017)

421 A minority of the respondents disclosed that they were no longer eating
422 undercooked meat although it was a difficult decision for them.

423 *“For health reasons, I no longer eat braai meat that is half cooked which oozes blood. It*
424 *is difficult I will not lie, but I love my liver and as you would know it is not consumed well-*
425 *done sometimes, it is eaten raw as this is believed that the liver in this state is very*
426 *nutritious. Another meat consumed raw is the intestines, but a lot has changed since the*
427 *discovery of diseases [...]”.* (Nibela dip tank cattle owner, December 2017)

428 The discussants also acknowledged that in some traditional practices certain
429 organs were meant for a specific group of people such as the chest for women, cow heels
430 (trotters) and head of the cow for both young and elderly men and the liver which was
431 given to the mother of a pregnant girl during marriage ceremonies or the elderly women
432 in the family.

433 Milk is the other main product from the cattle that is consumed as either raw milk,
434 boiled milk or as sour milk (“*aMasi*”).

435 *“We boil the milk but not always. When you are milking the cow you sometimes take*
436 *straight shots into your mouth from the cow’s udder and it is nice and warm”.* (Nkomo dip
437 tank cattle keeper, December 2017)

438 The preparation of “*aMasi*” was common to all groups and following traditional
439 methods.

440 “*We first milk the cow and pour the milk into a traditional (calabash) or into 2-liter bottle*
441 *of coke and it sits in a dry warm place maybe for three days and after that the “aMasi”*
442 *should be ready*”. (Nkomo dip tank women, December 2017)

443 Milk was also used for traditional rituals for example during the cleansing ceremony
444 of a widow, whereby the women used it to bath, and during funerals for the washing of
445 hands.

446 “*Other families use it at funerals at the gate for the washing of hands when coming back*
447 *from the gravesite. It is mixed with water*”. (Masakeni dip tank women, December 2017)

448 In some cases when a person had ingested poison or was constipated, milk was
449 also used as a neutralizing substance or laxative respectively.

450 “*When you have ingested something poisonous or poison itself you mix the milk with*
451 *some dung and then drink*”. (Masakeni dip tank, cattle keepers, December 2017)

452 **Discussion**

453 A qualitative research approach was used to investigate the risk factors for bovine
454 TB transmission to cattle to and humans using FGDs in the livestock-keeping community
455 living at the wildlife-livestock-human interface. The awareness and knowledge of bovine
456 TB in cattle and humans was also assessed using appropriate themes during the
457 discussions. The purpose of the study was to document the community’s perception of
458 bTB within their social and cultural context for the development of suitable interventions
459 targeted towards the control of zoonotic TB. This is in line with the initiative by

460 WHO/OIE/FAO/IUATLD (Road Map for Zoonotic Tuberculosis) to identify people at risk
461 of zoonotic TB especially in sub-Saharan Africa in support of the WHO's goal towards
462 eradication of TB by 2050 [18].

463 Despite the participants being aware of zoonotic diseases they could not state
464 specific examples of zoonotic diseases in livestock and wildlife. Contrastingly, a study in
465 the Gauteng province of South Africa documented historical knowledge of zoonotic
466 diseases amongst small-scale farmers and these recognized brucellosis as a zoonotic
467 disease [19]. In other studies in sub-Saharan region cattle owners stated d rabies,
468 anthrax, tuberculosis and brucellosis as major zoonotic diseases [20–22]. This lack of
469 knowledge could be linked to the limited awareness campaigns on zoonoses, the
470 absence of local information on zoonoses, inadequate communication between
471 veterinary and human health professionals as described by Cripps [23]. Documentation
472 from elsewhere in Africa indicates that the awareness, knowledge, attitude and perception
473 of zoonoses determines the increase or decrease in zoonotic risk in livestock keeping
474 communities and the general public [22,24].

475 Generally, the results showed a high awareness of bTB in cattle, the symptoms in
476 cattle and mode of bTB transmission to cattle as well as to humans, although this was
477 coupled with poor preventive practices. This is in contrast with other cattle keeping
478 communities where the knowledge of bTB was generally found to be low as revealed by
479 studies in Zambia, Tanzania and Ethiopia [25–27]. The high awareness in the community
480 could possibly be attributed to the bTB activities associated with the research program on
481 bTB conducted by our team which included the successful bTB information day and bTB
482 testing of cattle at the dip tanks. Prior education on zoonotic diseases has been

483 associated with a display of good knowledge of the disease by the cattle farming
484 community as demonstrated in a study in Uganda [28].

485 Through the FGDs it was noted that most people are at least involved in one
486 practice placing them at risk of bTB or other zoonotic diseases such as brucellosis. The
487 risky practices that are characteristic of pastoral communities particularly in sub-Saharan
488 Africa included consumption of raw (soured milk) or undercooked meat, poor handling
489 practices and absence of trained veterinary personnel during slaughtering of animals [29–
490 31]. Most of the participants indicated that they consumed boiled milk, but this practice
491 might have been overstated due to awareness by the participants of the accepted
492 practice. Research has shown that people are inclined to display paradoxical behavior
493 where one maybe aware but not apply precautionary measures [32]. Inadequate
494 precautionary measures during production, processing, handling of animal products and
495 slaughtering of cattle exposes individuals to bTB through direct contact with infected
496 carcasses [8,20,33]. The poor practices might also be linked to the socio-economic status
497 of the farmers whereby soured milk is readily available and cheap. Indeed, in contexts of
498 fragile livelihoods, disease risk might not be people's sole concern, or they might not have
499 the resources to take up protection measures such as wearing protective clothing during
500 slaughter [34]. The groups that were dominated by males, i.e. cattle owners, cattle
501 keepers and dip tank committee members were at a greater risk of *M. bovis* infection than
502 their female counterparts; as these were involved in unprotected practices during the
503 slaughter of animals and consumption of undercooked meat during “*braaing*”.

504 Consistent with other studies livestock keeping activities that were associated with
505 bTB transmission to cattle involved uncontrolled movement of animals and introduction

506 of animals into a herd without bTB pre-testing for social benefit, cultural or commercial
507 purpose [7,15,35–37]. The free movement of cattle in communal farming within the village
508 parameters is a common practice and movement to areas adjacent or into game reserves
509 in search of pasture was attributed to the frequent drought conditions in the area.
510 Livestock management practices that result in contact of uninfected herds with infected
511 herds as well as livestock-wildlife contact have been observed in bTB endemic settings
512 (South Africa and Zambia) and in the European context where occasional outbreaks have
513 been reported [38–40].

514 The findings from this study were similar across all the dip tanks with minor
515 differences concerning livestock-wildlife contact. Contact at the wildlife-livestock interface
516 has been suggested to be caused by porous boundaries that promote sharing of
517 resources that might result in the exchange of diseases between livestock and wildlife
518 [41]. There were no extensive differences in animal husbandry, cultural practices, food
519 consumption and handling behavior amongst the four different groups of participants
520 since these groups belong to the same geographical location and tribe. However, there
521 was a need to include all areas (dip tanks) with *M. bovis* infected herds for an in-depth
522 analysis of risky practices.

523 Most of the opinions expressed by the participants in the four categories were
524 similar with knowledge gaps or different views being expressed by the cattle keepers.
525 Amongst the differences identified were the poor knowledge on bTB transmission to
526 people, consumption of uninspected meat and the habit of drinking raw milk during
527 milking, implying that the cattle keepers are more at risk of *M. bovis* infection. Several
528 investigations have reported the potential risk of bTB transmission through drinking of raw

529 milk and contact with infected cattle [42–44]. Poor practices regarding TB have been
530 reported amongst high risk groups such as the cattle keepers and *M. bovis* transmission
531 to people was confirmed in a study of livestock workers in Nigeria [29,45].

532 The study highlighted the influence of belief, habits and socio-cultural aspects on
533 food processing (e.g. fermented milk), food consumption (undercooked or raw meat) and
534 introduction of animals into a herd (lobola ‘bride price’). The consumption of raw or
535 undercooked organs such as lymph nodes as influenced by socio-cultural practices has
536 been previously reported in another study in South Africa [19]. Therefore, cultural issues
537 related to drinking of raw blood, milk and meat (organs such as the liver and intestines);
538 selection of animals for slaughter during traditional ceremonies and acceptance of
539 animals into a herd without bTB pre-testing might be difficult to alter. Instead farmers
540 should be informed about potential consequences of certain practices in the spread of
541 bTB in cattle and people. Cultural practices are recognized as impediments in *M. bovis*
542 control strategies in developing countries, consequently 10-15% of human TB is
543 potentially caused by this pathogen [46]. Thus, policy makers need to be familiar with the
544 cultural practices associated with the cattle owners’ actions that influence bTB control to
545 be able to design effective preventive solutions.

546

547 Most of the data on bTB risk practices in previous studies was obtained using
548 quantitative methods only or mixed with qualitative methods, whilst in this study a
549 qualitative method of FGDs was applied. FGDs provide a platform for participants to
550 easily discuss their beliefs within the group atmosphere than individual interviews [47].
551 Using the different categories according to the people’s role in the community/households

552 allowed the participants to be comfortable and freely express their opinions. The
553 advantage of using FGDs is that the researcher can reach many people on one goal,
554 explore people's knowledge, experiences and collect a large amount of data within a short
555 period of time [48]. The limitation of this method is that the researcher has no control over
556 the information that is generated during the discussion and it's not the best way to obtain
557 responses on sensitive issues [49]. More participant observations would have
558 triangulated some data regarding practices (e.g. milk consumption), hence avoiding bias
559 raised during the discussions.

560 Using disease risk as a departure point in One Health studies of zoonoses is very
561 pertinent as reducing risk has a real public health, economic and social rationales.
562 However, it is very important to avoid that people's cultural logics or social practices are
563 cast in negative terms, as ignorance or superstition, or behaviors that exacerbate risk and
564 require changing [34]. Furthermore, how risk is perceived by livestock owners or by
565 animal and human health workers are not necessarily the same [50]. The endeavor of
566 addressing risky practices is complex and cannot involves a simple linear process of
567 social engineering because knowledge alone does not drive behavior change. Derived
568 from the social and psychological sciences, Kelly and Barker (2016) proposed to start
569 with the behavior, identify who is behaving and where, and working backwards using
570 regressive inference (understand the preceding conditions of the specifics) [51]. It is a
571 much more profitable avenue for developing interventions instead of predictive single
572 causal models such as the model of Theory of Planned Behavior which is elaborated on
573 rational assessment alone based on economic utility theory [51,52].

574

575 **Conclusion**

576 Our qualitative study allowed us to inform about a nuanced understanding of how
577 people experience and indeed conceptualize risk within specific socio-cultural practices
578 and a wider web of structural factors such as the intersection of poverty and gendered
579 divisions of labor affecting risk of bTB in an endemic area at the wildlife-livestock-human
580 interface in northern Kwa-Zulu Natal, South Africa. These findings are keys for the
581 elaboration of appropriate public health policy and related education campaigns. The
582 strengths of using a qualitative approach is its attention to complexity, questioning the
583 familiar, helping with language and translation, reconfiguring boundaries (e.g. human-
584 animal relations) to create novel frameworks and being reflective [53]. The participants
585 displayed good knowledge of bTB in cattle and its transmission to humans and cattle.
586 However, the perceived risks to humans and cattle were not translated into protective
587 practices as these were influenced by socio-cultural aspects and economic value of
588 livestock keeping in the community. Finally, concerted effort is required from all
589 stakeholders using the One Health strategy for the implementation of an all-
590 encompassing disease control program as control of zoonotic TB transmission to humans
591 is ultimately linked to the control of bTB in cattle.

592

593 **Recommendations**

594 A community-based animal health delivery system that provide basic instructions
595 on practices that reduce risk of disease transmission to cattle and people through
596 increased awareness campaigns is suggested. We suggest community involvement in

597 the planning of appropriate educational programs that consider human-animal relations,
598 social, cultural and economic reality of the community. These programs should involve
599 the community leaders (tribal leaders) that are respected by the community, health
600 caregivers and local veterinary personnel. In addition, targeted educational programs
601 would benefit risk groups, for example cattle keepers to improve their perception towards
602 zoonotic TB since the group is actively involved in livestock keeping activities.

603 **Acknowledgements**

604 The authors are indebted to the team of investigators involved in the study Tebogo
605 Mogoru (TM) who was the facilitator and Philani Zulu (PZ) first observer during the FGDs.
606 We acknowledge the veterinary staff at Big 5 False Bay Municipality offices especially Mr.
607 Simon Mkwamubi (Animal Health Technician) for their contribution to the success of the
608 study. We also appreciate the local tribal authorities for granting us the permission and
609 the farmers who agreed to participate in the study.

610

611 **References**

- 612 1. Michel AL, Müller B, van Helden PD. *Mycobacterium bovis* at the animal-human
613 interface: A problem, or not? *Veterinary Microbiology*. 2010;140(3–4):371–81.
- 614 2. Humblet MH, Boschioli ML, Saegerman C. Review article Classification of
615 worldwide bovine tuberculosis risk factors in cattle : a stratified approach. *Vet Res*.
616 2009;40(50):1–24.
- 617 3. Smith NH, Gordon S V, de la Rua-Domenech R, Clifton-Hadley RS, Glyn HR.

618 Bottlenecks and broomsticks: the molecular evolution of *Mycobacterium bovis*.
619 Nature Publishing Group. 2006;4:670–81.

620 4. Department of Agriculture, Forestry and Fisheries R of SA (DAFF). Bovine
621 Tuberculosis Scheme Manual 6. 2013.

622 5. Caron A, , de Garine-Wichatitsky M and RF. Bovine tuberculosis: a double-edged
623 issue at the human/livestock/wildlife interface in Africa. Empres -animal health.
624 2014;44(2):10–3.

625 6. Cosivi O, Grange JM, Daborn CJ, Raviglione MC, Fujikura T, Cousins D, et al.
626 Zoonotic tuberculosis due to *Mycobacterium bovis* in developing countries.
627 Emerging Infectious Diseases. 1998;4(1):59–70.

628 7. Munyeme M, Muma JB, Skjerve E. Risk factors associated with bovine
629 tuberculosis in traditional cattle of the livestock / wildlife interface areas in the
630 Kafue basin of Zambia. Preventive Veterinary Medicine. 2008;85(2008):317–28.

631 8. WHO. The Control of Neglected Zoonotic Diseases A route to poverty alleviation.
632 World Health. 2006.

633 9. Bobadilla-Del Valle M, Torres-González P, Cervera-Hernández ME, Martínez-
634 Gamboa A, Crabtree-Ramirez B, Chávez-Mazari B, et al. Trends of
635 *Mycobacterium bovis* Isolation and First-Line Anti-tuberculosis Drug Susceptibility
636 Profile: A Fifteen-Year Laboratory-Based Surveillance. PLOS Neglected Tropical
637 Diseases. 2015;9(9).

638 10. Miller M. Tuberculosis in South African wildlife: Why is it important? 2015.

639 11. Sichewo P, Etter E, Michel A. Prevalence of bovine tuberculosis in cattle at the
640 wildlife/livestock/human interface in Northern Kwazulu-Natal province, South
641 Africa. In: Proceedings of the 9th Veterinary, Paraveterinary & Southern African
642 Society for Veterinary Epidemiology and Preventive Medicine (SASVEPM)
643 Congress 24-27 July 2017 Boksburg, South Africa. 2017.

644 12. Shitaye JE, Tsegaye W, Pavlik I. Bovine tuberculosis infection in animal and
645 human populations in Ethiopia: a review. Veterinární Medicína. 2007;52(8):317–
646 32.

647 13. Mfinanga S, Morkve O, Kazwala R, Cleaveland S, Sharp M, Kunda J NR.
648 Mycobacterial Adenitis : Role of *Mycobacterium bovis*, Non-Tuberculous
649 Mycobacteria, HIV Infection, and risk factors in Arusha,Tanzania.
650 2004;81(4):171–8.

651 14. Katale BZ, Mbugi E V, Kendal S, Fyumagwa RD, Kibiki GS, Godfrey-Faussett P,
652 et al. Bovine tuberculosis at the human-livestock-wildlife interface: is it a public
653 health problem in Tanzania? A review. The Onderstepoort journal of veterinary
654 research. 2012;79(2):463.

655 15. Dejene SW, Heitkönig IMA, Prins HHT, Lemma FA, Mekonnen DA, Alemu ZE, et
656 al. Risk factors for bovine tuberculosis (bTB) in cattle in Ethiopia. PLoS ONE.
657 2016;11(7):1–16.

658 16. Munyeme M, Muma JB, Samui KL. Prevalence of bovine tuberculosis and animal
659 level risk factors for indigenous cattle under different grazing strategies in the
660 livestock / wildlife interface areas of Zambia. 2009;345–52.

661 17. Phillips CJ., Foster CR., Morris P., Teverson R. The transmission of
662 Mycobacterium bovis infection to cattle. Research in Veterinary Science.
663 2003;74(1):1–15.

664 18. World Health Organization (WHO) F and AO of the UN (FAO), World Organisation
665 for Animal Health (OIE) IUAT and LD, (IUATLD). Road map for zoonotic
666 tuberculosis. 48th Union World Conference on Lung Health; Guadalajara: Mexico;
667 2017. 2017.

668 19. Declercq G. Descriptive human health risk assessment of informal slaughter by
669 small-scale farmers of Gauteng , focussing on Brucella abortus. 2018.

670 20. Swai E, Schoonman L, Daborn.C. Knowledge and attitude towards zoonoses
671 among animal health workers and livestock keepers in Arusha and Tanga,
672 Tanzania. Tanzanian Journal of Health Research. 2010;12(4):8.

673 21. Gadaga BM, Etter EMC, Mukamuri B, Makwangudze KJ, Pfukenyi DM, Matope G.
674 Living at the edge of an interface area in Zimbabwe : cattle owners , commodity
675 chain and health workers ' awareness , perceptions and practices on zoonoses.
676 BMC Public Health. 2016;16(84):1–10.

677 22. Shirima GM, Fitzpatrick J, Kunda J, Mfinanga G, Kazwala R, Kambarage D, et al.
678 SHORT COMMUNICATIONS The role of livestock keeping in human brucellosis
679 trends in livestock keeping communities in Tanzania. Tanzanian journal health
680 research. 2010;12(3):1–6.

681 23. Cripps PJ. Veterinary education, zoonoses and public health: a personal
682 perspective. Acta, Tropica. 2000;76(1):77–80.

683 24. John K, Kazwala R, Mfinanga GS. Knowledge of causes , clinical features and
684 diagnosis of common zoonoses among medical practitioners in Tanzania. BMC
685 Infectious Diseases. 2008;8(162):1–8.

686 25. Grange JM. *Mycobacterium bovis* infection in human beings. 2001;81:71–7.

687 26. Munyeme M, Muma JB, Munang HM, Kankya C, Skjerve E. Cattle owners ' awareness of bovine tuberculosis in high and low prevalence settings of the
688 689 wildlife-livestock interface areas in Zambia. BMC veterinary research.
690 2010;6(21):1–9.

691 27. Getahun Asebe. EG. Community knowledge , attitudes and practices on bovine
692 tuberculosis and associated risk factors in Gambella Regional State Lare Woreda
693 , South West Ethiopia. Veterinaria. 2018;67(1):27–25.

694 28. Kazoora HB, Majalija S, Kiwanuka N, Kaneene JB. Knowledge, Attitudes and
695 Practices Regarding Risk to Human Infection due to *Mycobacterium bovis* among
696 Cattle Farming Communities in Western Uganda. Zoonoses and Public Health.
697 2016;63(8):616–23.

698 29. Mfinanga SG, Mørkve O, Kazwala RR, Cleaveland S, Sharp JM, Shirima G, et al.
699 The role of livestock keeping in tuberculosis trends in Arusha , Tanzania.
700 International Journal of Tuberculosis and Lung Disease. 2003;7(7):695–704.

701 30. Ayele WY, Neill SD, Zinsstag J, Weiss MG, Pavlik I. Bovine tuberculosis: an old
702 disease but a new threat to Africa. The international journal of tuberculosis and
703 lung disease : the official journal of the International Union against Tuberculosis
704 and Lung Disease. 2004;8(8):924–37.

705 31. Hambolu D, Freeman J, Taddese HB. Predictors of Bovine TB Risk Behaviour
706 amongst Meat Handlers in Nigeria: A Cross-Sectional Study Guided by the Health
707 Belief Model. *PLoS ONE*. 2013;8(2).

708 32. Wachinger G, Renn O, Begg C, Kuhlicke C. The Risk Perception Paradox —
709 Implications for Governance and Communication of Natural Hazards. *Risk
710 Analysis*. 2013;33(6):1049–65.

711 33. Chowdhury TA, Shanzana P, Zahan FN. Knowledge , Awareness and Risks of
712 Zoonotic Diseases Among the Smallholder Livestock Farmers in Suburban.
713 *Advances in Biology and Earth Sciences*. 2018;3(1):69–84.

714 34. Macgregor H, Waldman L, Macgregor H. Views from many worlds : unsettling
715 categories in interdisciplinary research on endemic zoonotic diseases.
716 *Philosophical Transactions Royal Society B*. 2017;372(20160170).

717 35. Tschopp R, Schelling E, Hattendorf J, Aseffa A, Zinsstag J. Risk factors of bovine
718 tuberculosis in cattle in rural livestock production systems of Ethiopia. *Preventive
719 Veterinary Medicine*. 2009;89(2009):205–11.

720 36. Kazwala RR, Kambarage DM, Daborn CJ, Nyange J, Jiwa SFH, Sharp JM. Risk
721 Factors Associated with the Occurrence of Bovine Tuberculosis in Cattle in the
722 Southern Highlands of Tanzania.

723 37. Oloya J, Muma JB, Opuda-asibo J, Djønne B. Risk factors for herd-level bovine-
724 tuberculosis seropositivity in transhumant cattle in Uganda. *Preventive Veterinary
725 Medicine*. 2007;80(2007):318–29.

726 38. Munyeme M, Muma JB, Siamudaala VM, Skjerve E, Munang'andu HM, Tryland
727 M. Tuberculosis in Kafue lechwe antelopes (*Kobus leche Kafuensis*) of the Kafue
728 Basin in Zambia. *Preventive Veterinary Medicine*. 2010;95(3–4):305–8.

729 39. Hlokwe TM, van Helden P, Michel AL. Evidence of increasing intra and inter-
730 species transmission of *Mycobacterium bovis* in South Africa: Are we losing the
731 battle? *Preventive Veterinary Medicine*. 2014;115(1–2):10–7.

732 40. Kaneene JB, Colleen ;, Bruning-Fann S, Granger LM, Miller R, Porter-Spalding
733 BA. Environmental and farm management factors associated with tuberculosis on
734 cattle farms in northeastern Michigan. Vol. 221, *JAVMA*. 2002.

735 41. de Garine -Wichatitsky M, Caron A, Kock R, Tschopp R, Munyeme M, Hofmeyer
736 M, et al. A review of bovine tuberculosis at the wildlife–livestock–human interface
737 in sub-Saharan Africa. *Epidemiology and Infection*. 2013 Jul 15;141(07):1342–56.

738 42. Cadmus SIB, Yakubu MK, Soolingen D Van. *Mycobacterium bovis* , but also *M .*
739 *africanum* present in raw milk of pastoral cattle in north-central Nigeria. *Tropical*
740 *Animal Health Production*. 2010;42(2010):1047–8.

741 43. Ereqat S, Nasereddin A, Levine H, Azmi K, Al-Jawabreh A, Greenblatt CL, et al.
742 First-Time Detection of *Mycobacterium bovis* in Livestock Tissues and Milk in the
743 West Bank, Palestinian Territories. Phillips RO, editor. *PLoS Neglected Tropical*
744 *Diseases*. 2013 Sep 12;7(9):e2417.

745 44. Nuru A, Mamo G, Zewude A, Mulat Y, Yitayew G, Admasu A, et al. Preliminary
746 investigation of the transmission of tuberculosis between farmers and their cattle
747 in smallholder farms in northwestern Ethiopia: A cross-sectional study. *BMC*

748 Research Notes. 2017;10(1):1–7.

749 45. Adesokan HK, Jenkins AO, Soolingen D Van, Cadmus SIB. *Mycobacterium bovis*
750 infection in livestock workers in Ibadan , Nigeria : evidence of occupational
751 exposure. *INT J TUBERC LUNG DIS.* 2012;16(February):1388–92.

752 46. de la Rua-Domenech R. Human *Mycobacterium bovis* infection in the United
753 Kingdom: Incidence, risks, control measures and review of the zoonotic aspects
754 of bovine tuberculosis. *Tuberculosis.* 2006 Mar 1;86(2):77–109.

755 47. M.E.Edginton, C.S. Sekatane SJG. Patients' beliefs: Do they affect tuberculosis
756 control? A study in a rural district of South Africa. *International Journal of*
757 *Tuberculosis and Lung Disease.* 2002;6(12):1075–82.

758 48. Masadeh MA. Focus Group : Reviews and Practices. *International Journal of*
759 *Applied Science and Technology.* 2012;2(10):63–8.

760 49. Mfinanga SG, Mørkve O, Kazwala RR, Cleaveland S, Sharp JM, Shirima G, et al.
761 Tribal differences in perception of tuberculosis : a possible role in tuberculosis
762 control in Arusha , Tanzania. *International Journal of Tuberculosis and Lung*
763 *Disease.* 2003;7(10):933–41.

764 50. Ainslie A. Farming Cattle , Cultivating Relationships : Cattle Ownership and
765 Cultural Politics in Peddie District , Eastern Cape Farming Cattle , CULTivating
766 Relationships : Cattle Ownership and Cultural Politics in Peddie District , Eastern
767 Cape Introduction : Situ. *Social Dynamics.* 2008;31(1):129–56.

768 51. Kelly MP, Barker M. Europe PMC Funders Group Why is changing health-related

769 behaviour so difficult ? Public Health. 2017;July(136):109–16.

770 52. Ajzen I. Attitudes, Personality and Behaviour. Chicago: 1988. 1988.

771 53. Porter JDH. Epidemiological reflections of the contribution of anthropology to
772 public health policy and practice. Journal of Biosocial Science. 2006;38:133–44.

773

774