

A polygenic and phenotypic risk prediction for Polycystic Ovary Syndrome evaluated by Phenome-wide association studies

Short title: PRS & PheWAS of PCOS in 124,852 adults from electronic health records

Yoonjung Yoonie Joo¹, Ky'era Actkins², Jennifer A. Pacheco³, Anna O. Basile⁴, Robert Carroll⁵, David R. Crosslin⁶, Felix Day⁷, Joshua C. Denny⁵, Digna R. Velez Edwards^{5,8}, Hakon Hakonarson^{9,10}, John B. Harley^{11,12}, Scott J Hebring¹³, Kevin Ho¹⁴, Gail P. Jarvik¹⁵, Michelle Jones¹⁶, Tugce Karder¹⁷, Frank D. Mentch⁹, Cindy Meun¹⁸, Bahram Namjou¹¹, Sarah Pendergrass¹⁴, Marylyn D. Ritchie¹⁹, Ian B. Stanaway⁶, Margrit Urbanek¹, Theresa L. Walunas²⁰, Maureen Smith³, Rex L. Chisholm³, International PCOS Consortium, Abel N. Kho²⁰, Lea Davis⁵, M. Geoffrey Hayes^{1,3,21}

1. Division of Endocrinology, Metabolism, and Molecular Medicine, Department of Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL, 60611, USA
2. Department of Microbiology, Immunology, and Physiology, Meharry Medical College, Nashville, TN, 37203, USA
3. Center for Genetic Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL, 60611, USA
4. Department of Biomedical Informatics, Columbia University New York, NY, 10032, USA
5. Departments of Biomedical Informatics and Medicine, Vanderbilt University Medical Center, Nashville, TN, 37203, USA
6. Department of Biomedical Informatics and Medical Education, University of Washington School of Medicine, Seattle, WA, 98195, USA
7. MRC Epidemiology Unit, Cambridge Biomedical Campus, University of Cambridge School of Clinical Medicine, Cambridge, CB2 0QQ, United Kingdom
8. Division of Quantitative Sciences, Department of Obstetrics and Gynecology, Vanderbilt University Medical Center, Nashville, TN, 37203, USA
9. Center for Applied Genomics, Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA

10. Department of Pediatrics, The Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, 19104, USA
11. Center for Autoimmune Genomics and Etiology (CAGE), Cincinnati Children's Hospital Medical Center, Cincinnati, OH 45229, USA
12. Department of Pediatrics, University of Cincinnati College of Medicine; US Department of Veterans Affairs, Cincinnati, OH 45229, USA
13. Center for Precision Medicine Research, Marshfield Clinic Research Institute, Marshfield, WI, 54449, USA
14. Biomedical and Translational Informatics, Geisinger, Danville, PA, 17822, USA
15. Division of Medical Genetics, Department of Medicine (Medical Genetics) and Genome Sciences, University of Washington Medical School, Seattle, WA, 98195, USA
16. Center for Bioinformatics & Functional Genomics, Department of Biomedical Sciences, Cedars-Sinai Medical Center, Los Angeles, CA, 90048, USA
17. The Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, OX3 7BN, United Kingdom
18. Department of Obstetrics and Gynecology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands
19. Department of Genetics, University of Pennsylvania, Philadelphia, PA, 19104, USA
20. Division of General Internal Medicine and Geriatrics, Department of Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL, 60611, USA
21. Department of Anthropology, Northwestern University, Evanston, IL 60208, USA

Keywords

Phenome-wide association study, Genomic prediction, Polygenic risk score, Polycystic Ovary Syndrome

Abstract (~300 words)

Purpose

As many as 75% of patients with Polycystic ovary syndrome (PCOS) are estimated to be unidentified in clinical practice. Utilizing polygenic risk prediction, we aim to identify the phenome-wide comorbidity patterns characteristic of PCOS to improve accurate diagnosis and preventive treatment.

Methods and Findings

Leveraging the electronic health records (EHRs) of 124,852 individuals, we developed a PCOS risk prediction algorithm by combining polygenic risk scores (PRS) with PCOS component phenotypes into a polygenic and phenotypic risk score (PPRS). We evaluated its predictive capability across different ancestries and perform a PRS-based phenome-wide association study (PheWAS) to assess the phenomic expression of the heightened risk of PCOS. The integrated polygenic prediction improved the average performance (pseudo- R^2) for PCOS detection by 0.228 (61.5-fold), 0.224 (58.8-fold), 0.211 (57.0-fold) over the null model across European, African, and multi-ancestry participants respectively. The subsequent PRS-powered PheWAS identified a high level of shared biology between PCOS and a range of metabolic and endocrine outcomes, especially with obesity and diabetes: ‘morbid obesity’, ‘type 2 diabetes’, ‘hypercholesterolemia’, ‘disorders of lipid metabolism’, ‘hypertension’ and ‘sleep apnea’ reaching phenome-wide significance.

Conclusions

Our study has expanded the methodological utility of PRS in patient stratification and risk prediction, especially in a multifactorial condition like PCOS, across different

genetic origins. By utilizing the individual genome-phenome data available from the EHR, our approach also demonstrates that polygenic prediction by PRS can provide valuable opportunities to discover the pleiotropic phenomic network associated with PCOS pathogenesis.

Introduction

Polycystic ovary syndrome (PCOS) is the most common reproductive metabolic disorders, affecting 5-15% of reproductive age women worldwide [1]. The estimated cost of diagnosing and treating American women with PCOS is \$5.46 billion annually as of 2017 [2, 3]. In addition to being a major cause of female infertility, the disease is a well-known risk factor for endocrine complications, such as type 2 diabetes, impaired glucose tolerance, and metabolic syndrome before age 40 [4]. Monozygotic twin studies of PCOS have suggested that PCOS is highly heritable ($h^2 = \sim 70\%$) [5] and the genetic architecture is polygenic with complex genetic inheritance pattern [6, 7]. Despite its clinical importance and high heritability, the underlying genetic etiology of PCOS remains incompletely understood. The phenotypic manifestations of PCOS are heterogeneous and exhibit considerable variation across race and ethnicity, further complicating the clinical diagnosis. Currently, it is estimated that up to 75% of women with PCOS remain undiagnosed in part due to varying diagnostic criteria from the National Institutes of Health (NIH), Rotterdam, or Androgen Excess Society, [8-12] which use different combinations of hyperandrogenism, ovulatory dysfunction, and/or polycystic ovarian morphology. Despite shared genetic risk across the criteria [13], the disagreement regarding PCOS phenotypic criteria presents a significant challenge for both clinical practice and research [14, 15]. The commonalities and differences between the phenotypic characteristics of PCOS may be better understood with an integrative observation of phenome-wide pleiotropies and co-morbidities.

Polygenic risk scores (PRS) built from well-powered genome-wide association studies (GWAS) have demonstrated operationalizing potential as biological risk predictors for patient stratification and risk prediction [16-19]. PRS represents the cumulative effect of common genetic variation summed per individual into a single risk score, providing an intuitive way to translate GWAS findings into clinically relevant information such as a patient's risk of disease [20, 21]. From a precision medicine perspective, PRS hold significant promise especially for a multifactorial condition with complicated clinical manifestations, such as PCOS. However, several practical challenges remain in the equitable translation of PRS into clinical practice [22, 23]. For instance, most GWAS have been performed in samples of primarily European ancestry, resulting in PRS statistics that systematically perform worse in populations of different ancestry, including African ancestry populations. This underperformance is due to a combination of population-specific genetic effects that are undetected in a Euro-centric GWAS, and differences in the patterns of linkage disequilibrium (LD) between populations of differing biogeographic ancestry [24-27]. Thus, the evaluation of PRS from existing GWAS in both European and non-European ancestry samples is a critical step in setting priorities for equitable precision medicine initiatives.

The widespread deployment of Electronic Health Records (EHRs) and the availability of these multi-dimensional records enables evaluation of PRS in a research context that mimics a clinical hospital setting. Using these data, the predictive capability of PRS can be assessed regarding many possible diagnoses that can accumulate during an individual's lifespan (i.e., the phenome). The eMERGE (electronic MEDical Records and GEnomics) Network is a nationwide consortium of multiple medical

institutions that link DNA biobanks to EHRs [28], which is an important resource for determining the clinical utility of genomic findings, and enabling exploration of the range of phenotypes associated with genetic variation [29, 30].

The aim of this study is to systematically examine the utility of PRS derived from a GWAS meta-analysis by the International PCOS Consortium [13] for risk prediction across multiple ancestries and to further characterize the other EHR phenotypes that are clinically associated with PCOS genetic risk in both women and men. We first developed the integrative polygenic and phenotypic risk score (PPRS) for PCOS by combining the patient DNA genotype information and PCOS phenotypic elements from the EHR. Then we tested the predictive utility of the algorithm within European ancestry (EA) samples and further evaluated its performance in African ancestry (AA) and combined multi-ancestry (MA) participants which included EA, AA, and other ancestries. In addition, we performed a Phenome-Wide Association Study (PheWAS) of the PPRS for PCOS to identify the range of phenotypic indicators associated with PCOS and evaluated the predictive characteristics of PPRS to identify underlying PCOS pathophysiological pathways.

Materials and Methods

PCOS Polygenic Risk Score (PRS) Development

We obtained the full summary statistics of the largest meta-GWAS of PCOS through the International PCOS consortium and developed a PRS for PCOS [13].

(Supplementary table 1) The GWAS was conducted in 5,209 cases and 32,055 controls of EA women who were diagnosed according to either NIH or Rotterdam criteria. All variant positions were converted to GrCh37 and we excluded any entries with missing ORs or risk allele frequency (RAF) information. The RAF of each variant was calculated using PLINK [31], and we excluded the entries which RAF deviates more than 15% than our eMERGE data in order to ensure additional quality control (QC). PRSice software [32] was used to filter any correlated SNVs in pairwise Linkage Disequilibrium (LD) ($r^2 > 0.2$) and constructed PRS for PCOS by summing the best-guess imputed genotype data of PCOS risk variants in each individual weighted by the reported effect sizes. We used eight different subsets of PCOS susceptibility SNVs to build the model based on p-value cutoff and compared for their predictive accuracy in the following validation step: 5×10^{-8} , 5×10^{-7} , 5×10^{-6} , 5×10^{-5} , 5×10^{-4} , 5×10^{-3} , 5×10^{-2} , and 1 (All SNVs).

PRS/PPRS Evaluation & PheWAS Discovery Cohort

Our cohort included genotypes and clinical diagnosis records of 99,185 individuals collected from 12 EHR-linked biobanks nationwide through the eMERGE consortium [29]. After identity-by-descent (IBD) analysis, we removed 8,019 related individuals that

were not in canonical IBD position or genetically identical individuals near the origins ($Z_0 > 0.83$ and $Z_1 < 0.1$). The cohort was composed of multiple self-reported and 3rd party observed ancestries and we defined them into three main genetic ancestral groups using the intersection of self-reported ancestries and principal component analysis (PCA) based k-mean clusters: European (71.7%), African (15.0%), and Asian (1.0%). We excluded any self-reported or genetically Hispanic participants for ancestry-stratified analysis for better homogeneity. Throughout this study, the first four principal components (PCs) were used to correct population structure, explaining over 17% of the variances among different genetic origins.

The phenome data of the participants were collected from the EHR including diagnostic records and basic demographic information. The data collection was performed under local institutional review board approval with informed consent from the patients. Diagnostic information was structured in the format of the International Classification of Diseases, Clinical Modification (ICD-CM) codes, in both 9th and 10th edition, and aggregated into a higher level of 1,711 phecodes for a standardized categorical analysis of diseases (Phecode map version 1.2) [33, 34]. We excluded 23 individuals under the age of 14, the clinically plausible age for PCOS diagnosis, which is defined as two years after the first menstruation. A demographic information of the 91,144 participants after filtering criteria is presented in **Table 1**.

Genotype data and Quality Control

The participants provided their saliva samples for genotyping, which were genotyped on 78 genotype Illumina or Affymetrix array batches from 12 medical sites.

Table 1: Demographic and clinical characteristics of discovery cohorts (eMERGE) and replication cohort (BioVU).

Site*	N Subjects	Sex (Female)	Ancestry (EA)	Ancestry (AA)	Age Average	Age SD	BMI Average	BMI SD	PCOS cases	Hirsutism cases	Irregular Mense cases	Female Infertility cases
BSCH**	862	362 (42.2%)	623	74	N/A	N/A	N/A	N/A	2	5	20	0
CCHMC**	5385	2320 (43.2%)	4058	523	8.9	6.7	20.9	6.2	11	24	54	2
CHOP**	9528	4376 (46.0%)	4898	4105	9.8	5.3	21.1	6.2	47	39	205	2
Columbia	2029	989 (48.8%)	519	143	56.1	19.8	27.0	5.4	3	4	15	1
Geisinger	2785	1320 (47.5%)	2439	8	62.8	15.7	32.6	8.1	77	48	158	8
Harvard	23922	13135 (55.0%)	20727	1343	55.3	16.5	28.3	5.8	417	322	2284	217
KPW/UW	3225	1829 (56.7%)	2891	109	76.1	8.9	26.4	4.8	2	25	10	18
Mayo Clinic	9307	4672 (50.2%)	6680	17	61.7	15.4	29.3	5.8	48	85	217	17
Marshfield	3725	2255 (60.9%)	3696	2	69.3	11.0	29.6	6.0	6	84	476	43
Mt. Sinai	5765	3362 (58.8%)	702	3643	59.6	10.0	30.6	7.4	51	45	200	15
Northwestern	4719	3913 (82.9%)	2250	301	53.7	14.8	28.7	7.2	65	83	280	51
Vanderbilt***	19892	10810 (54.4%)	15902	3371	56.6	17.1	29.4	7.1	220	144	1017	48
All (Discovery Cohort)	91144	49343	65385	13639	.	.	.	.	949	908	4936	422

	N Subjects	Sex (Female)	Ancestry (EA)	Ancestry (AA)	Age Average	Age SD	BMI Average	BMI SD	PCOS cases	Hirsutism cases	Irregular Mense cases	Female Infertility cases
VUMC Replication Sample (BioVU)	33708	18096 (54%)	33708	N/A	55.7	20	28.2	6.8	284	225	4330	48

* BSCH = Boston Children's Hospital, CCHMC=Cincinnati Children's Hospital Medical Center, CHOP= Children's Hospital of Philadelphia, KPW/UW = Group Health Cooperative/University of Washington

** Children's hospital with low average age

*** No Sample Overlap with replication cohort (BioVU)

We used the Michigan Imputation Server(MIS) [35] with the minimac3 missing genotype variant imputation algorithm to impute missing genotypes in our sample based on the Haplotype Reference Consortium (HRC1.1) which includes ~65,000 individuals of diverse ancestry [36]. The imputation resulted in a genome-wide set of ~40 million SNVs. We filtered the poorly imputed genetic variants with the r-squared imputation quality threshold (mean variant r-square) less than 0.3, minor allele frequency (MAF) less than 0.05 and genotype call rate lower than 95%, which resulted in 5,760,270 autosomal polymorphic variants for subsequent analysis. The detailed data collection and QC report for the eMERGE network is reported elsewhere [29].

Validation of Polygenic Risk Score

A. Predictive ability of each prediction model with different PRS

We performed logistic regression analysis to demonstrate the prediction ability of PRS for PCOS diagnosis in the female population of three different genetic racial cohorts: European (n=33,869), African (n=8,198), and the entire admixed cohort (n=49,365). Each cohort was randomly divided into 75% training and 25% testing set to separately calculate the regression statistics and out-of-sample prediction error. Using generalized linear model, the residuals of PRS after covariate adjustments (first four PCs, sites) were obtained and scaled to build the logistic regression model in the training set. Regression coefficients and p-value of PRS variable, and pseudo-R² of the eight different PRS models were measured.

We applied the regression model built out of the training set to measure out-of-sample performance in the testing dataset. We predicted the individuals as ‘PCOS cases’ if their fitted scores are higher than the average fitted score and calculated the accuracy by comparing with their actual diagnosis records of PCOS. The overall accuracy, sensitivity, specificity of each model were measured and structured through confusion matrix. The area under the receiving operating characteristic (ROC) curve, AUC, was also measured for classifier performance of each model.

B. Stratification ability of each prediction model with different PRS

To evaluate the phenotypic stratification ability of PRS, we divided the cohort into ten quantiles based on PRS of each individual and compared the average phenotypic values (e.g. proportion of PCOS diagnosed patients, body mass index (BMI), PRS) among the groups. The proportion of PCOS patients in each quantile, average PRS values, and average BMI measures of each individual were analyzed. We also performed independent t-test to assess if the average PRS score differences between PCOS cases and controls were statistically significant.

C. Performance improvement by the PRS variable

To estimate the performance of the PRS variable, we built a null regression model without the PRS variable for PCOS prediction (PRS model vs. Null model). The incremental pseudo- R^2 by McFadden’s [37] were calculated between the PRS models and the null logistic regression only with first 4 PCs and site variables. The analysis of variance (ANOVA) was performed to examine how significant PRS variable impacts the PCOS diagnosis prediction model compared to the null model.

156

157 PRS model:

$$158 \quad \text{logit}(\text{PCOS diagnosis} = 1) = \beta_0 * \text{PRS} + \beta_1 * \text{Site} + \beta_2 * 4\text{PCs} + \beta_3$$

159 Null model:

$$160 \quad \text{logit}(\text{PCOS diagnosis} = 1) = \beta_0 * \text{Site} + \beta_1 * 4\text{PCs} + \beta_2$$

161

162 ***Development of prediction algorithms with PRS and PCOS component***

163 ***phenotypes (PPRS)***

164 We built an integrative polygenic and phenotypic risk score (PPRS) with PRS and
 165 PCOS component phenotypes in the EHR to maximize the utility of PRS for risk
 166 prediction. Additional dichotomous phenotypic variables to each individual from their
 167 EHR diagnosis records: hirsutism (ICD9 code 704.1, ICD10 code L68.0), irregular
 168 menstruation (ICD9 code 626.4, ICD10 code N92.6), and female infertility (ICD9 code
 169 627, ICD10 code N97.0) were selected, all of which are well-established clinical
 170 components of PCOS. A total 908 individuals with hirsutism, 4,936 individuals with
 171 irregular menstruation, and 422 individuals with female infertility ICD diagnosis codes
 172 were identified in the eMERGE consortium database.

173 Firstly, the logistic regression adjusted for first four PCs and sites were examined
 174 for their effect coefficients and variable p-values. Psuedo-R² of each model was
 175 calculated for measuring the improvement over the normal PRS model. ANOVA
 176 between the integrative model and normal PRS model were examined.

PPRS model:

$$\text{logit}(\text{PCOS diagnosis} = 1) = \beta_0 * \text{PRS} + \beta_1 * \text{Site} + \beta_2 * 4\text{PCs} + \beta_3 * \text{Hirsutism} \\ + \beta_4 * \text{Irregular menstruation} + \beta_5 * \text{Female infertility} + \beta_6$$

PPRS null model:

$$\text{logit}(\text{PCOS diagnosis} = 1) = \beta_0 * \text{Site} + \beta_1 * 4\text{PCs} + \beta_2 * \text{Hirsutism} + \\ \beta_3 * \text{Irregular menstruation} + \beta_4 * \text{Female infertility} + \beta_5$$

Phenome-wide analysis

To investigate the potential pleiotropy of PCOS, PCOS components, and other diseases in the EMR phenome, we selected the best performing PRS model that presented a validated predictive accuracy and stratification ability across ancestries based on the examination results above. PheWAS was performed on the mapped 1,711 representative EHR phenotypes with a minimum of 30 case patients from the discovery cohort of 91,144 participants after QC criteria. Case group for a given phecode is defined by the presence of at least one assignment of the corresponding ICD codes from EHR as defined in the phecode map v1.2. Controls for each phecode are defined by the absence of the same ICD codes that defined cases and the absence of clinically related phenotypes. Based on the assumption that a participant with higher PCOS-PRS conveys greater genetic risk, our main sex-stratified PheWAS interrogated the comorbid networks of high-risk predictive phenotypes for PCOS (**PheWAS-1**). 49,343 female participants and 41,669 male participants were used for the analysis. Logistic

regression was used adjusting for genotyping site and the first four PCs of ancestry to correct for population stratification in the MA cohort [$\text{logit}(\text{Clinical Phenotype} = 1 \mid \text{PRS}, \text{Site}, 4\text{PCs}) = \beta_0 + \beta_1 \cdot \text{PRS} + \beta_2 \cdot \text{Site} + \beta_3 \cdot 4\text{PCs}$].

In this study, phenome-wide significance refers to either (1) the Bonferroni corrected threshold of $p\text{-value} = 2.9 \times 10^{-5}$ adjusting for multiple testing, which is determined by using the $p\text{-value}$ of 0.05 divided by the 1,711 phenotypes interrogated, or (2) the false discovery rate (FDR) significance of 0.05, which is a popular alternative threshold to the stringent Bonferroni correction in reporting PheWAS. Manhattan PheWAS plots of $-\log_{10}(p\text{-value})$ were generated for visual inspection of significant clinical traits. All the analyses were performed in the R statistical software environment (ver 2.1.2).

Sensitivity Analysis

We performed several comparative PheWAS in an effort to interrogate different phenome-wide aspects of the PRS in clinical phenome.

Firstly, to distinguish secondary or symptomatic phenotypes derived from the PCOS-diagnosed patients, we removed the clinical diagnosis records of the 949 individuals with PCOS (phecode 256.4, ICD9 256.4 and ICD10 E28.2) and performed the same PheWAS analysis. **(PheWAS-2)**. Additionally, to gauge the contrasting performance of polygenic prediction over a single-variant approach, we performed traditional PheWAS of each genome-wide significant susceptibility loci ($p\text{-value} < 5 \times 10^{-8}$) for PCOS (RAF > 0.05). This analysis aims to compare the clinical phenotypes associated with the cumulative effects of multiple genetic variants on PRS versus a

single genetic signal generated by an individual PCOS susceptibility locus. Among genome-wide significant loci ($p\text{-value} < 5 \times 10^{-8}$) for PCOS, (**Supplementary Table 1**) we filtered the entries with $MAF > 0.05$ and genotype call rate > 0.90 in our discovery cohort and $MAF > 0.01$ in summary statistics. 85 SNVs were selected and used for the subsequent PheWAS analysis (**PheWAS-3**).

PRS PheWAS Replication

To confirm the predictive performance of our PRS algorithm and its effect on clinical phenome, replication analyses were performed at Vanderbilt University Medical Center on an independent genotyped sample of 33,708 European descent individuals (BioVU). The participants were genotyped on the Illumina MEGAEX platform (~2 million markers) and we applied filters for individual call rates $< 98\%$, batch effects ($p\text{-value} < 5 \times 10^{-5}$), heterozygosity ($|F_{het}| > 0.2$), and sample relatedness ($\pi_{hat} > 0.2$). After imputation with 1000G reference panel, we excluded any genetic variants with missingness > 0.02 , certainty < 0.9 , or imputation info score < 0.95 . The genetic ancestry of the samples were restricted to only EA, based on comparison to 1000G European population and a K-means clustering definition. The final samples included 33,708 individuals of European descent genotyped on 5,550,390 SNVs. Using the same PRS generation methodology in discovery samples, we took the identical phenome-wide approach to identify the associated phenotypic networks with PRS among the replication samples. Logistic regression was used adjusting for first four ancestry PCs.

Results

Polygenic risk scores for PCOS are normally distributed in European and multi-ancestry participants

A total of 5,760,270 autosomal single nucleotide variants (SNVs) were considered for the PCOS-PRS construction, which displays the genetic architecture of effect size (beta) by risk allele frequency (RAF) presented in **Figure 1**. There was a significant negative correlation between RAF and effect size, which is generally anticipated in common quantitative traits and supports the use of methodology of PRS to explore the extreme of the common polygenic liability spectrum. According to the central limit theorem, PRS in a large population will show normality when the genetic architecture of the target trait is polygenic, i.e. produced by the addition of many genetic variants of small effect [38, 39].

PRS were calculated at eight different p-value cutoffs from the PCOS GWAS summary statistics (5×10^{-8} , 5×10^{-7} , 5×10^{-6} , 5×10^{-5} , 5×10^{-4} , 5×10^{-3} , 5×10^{-2} , 1) for all the discovery eMERGE participants (n=91,144). Each set of scores were adjusted for participant site and first four PCs. All the polygenic scores were evaluated for their predictive performance in the female populations of EA (n=33,869), AA (n=8,198) and MA cohorts (n=49,365). The covariate-adjusted PCOS-PRS generally presented a normal distribution across each ancestry cohort (**Supplementary Figure 1**). PRS models with trimodal or skewed distributions (PRS p-value cutoff: 5×10^{-7} , 5×10^{-6} , 5×10^{-5}), which may be a function of poor representation of risk variants across populations, were not considered for the subsequent phenome-wide analysis.




Fig 1. Effect distribution of PCOS susceptibility variants in samples from the International PCOS consortium by risk allele frequency. (a) The 120,340 PCOS autosomal SNVs with p-value < 0.05, and **(b)** the 139 PCOS genome-wide significant SNVs (p-value < 5×10⁻⁸). The dark green line and grey band around it are the linear regression fit and its 95% confidence interval, respectively, between risk allele frequency and effect size (beta).

265 **Validation of PCOS PPRS in European ancestry participants**

A. Predictive ability of each prediction model with different PRS

In the PRS prediction models using the training set of the female EA cohort (n=33,869 with 632 PCOS cases), all the coefficient p-values of the PRS variables are statistically significant except for two PRS models of SNVs with p-value $< 5 \times 10^{-7}$ and p-value $< 5 \times 10^{-6}$ that do not show PRS normality (**Supplementary Figure 1**). The average odds ratios (OR) of the significant PRS variable across EA was 1.13 (average SE=0.046) and the average pseudo- R^2 value was 0.044, which indicates 4.4% of the phenotypic variances in the training sample could be explained by PRS (**Table 2**).

The regression models built in the training set were then used to predict PCOS case status in the testing dataset. A model including PRS yielded average prediction accuracy of 0.55, sensitivity of 0.55, specificity of 0.76 with an average area under the receiving operating characteristic curve (AUC) of 0.72 in the EA participants (**Table 3**).

B. Stratification ability of each prediction model with different PRS

The percentage of PCOS-diagnosed patients increases in higher PRS quantiles, and the individuals in the highest PRS group tend to have higher average BMI. In the genome-wide PRS calculation with SNVs with p-value ≤ 1 , the average BMI of the individuals in highest PRS quantile is 1.1 kg/m² higher than the individuals in the lowest PRS group (Cohen's d=0.16, t-test p-value= 1.06×10^{-9}) (**Figure 2 and Table 4**). The finding confirms the positive correlation between elevated generic risk for PCOS, actual PCOS diagnosis, and higher risk for increased BMI.

Table 2: Regression results of the PRS and PPRS models in PCOS prediction.

	PRS*				PPRS*			
PRS/PPRS p-value Cutoff	OR	Std. Error	P	R ²	OR	Std. Error	P	R ²
<u>European Ancestry</u>								
5E-08	1.14	0.047	4.76E-03	0.045	1.14	0.052	1.40E-02	0.232
5E-07	1.04	0.042	3.78E-01	0.043	1.04	0.046	3.89E-01	0.230
5E-06	1.08	0.039	6.41E-02	0.044	1.08	0.044	7.26E-02	0.231
5E-05	1.10	0.041	2.13E-02	0.044	1.10	0.046	3.59E-02	0.231
5E-04	1.13	0.045	6.12E-03	0.044	1.11	0.049	2.85E-02	0.231
5E-03	1.11	0.047	2.70E-02	0.044	1.08	0.051	1.45E-01	0.231
5E-02	1.16	0.048	2.11E-03	0.045	1.12	0.052	3.21E-02	0.231
1	1.15	0.049	3.13E-03	0.045	1.11	0.052	4.04E-02	0.231
<u>Multiancestry</u>								
5E-08	1.16	0.038	1.15E-04	0.040	1.15	0.042	1.19E-03	0.228
5E-07	1.08	0.037	4.28E-02	0.038	1.09	0.038	2.99E-02	0.227
5E-06	1.09	0.037	1.60E-02	0.038	1.10	0.038	1.19E-02	0.227
5E-05	1.12	0.037	2.35E-03	0.039	1.12	0.038	3.67E-03	0.228
5E-04	1.12	0.038	1.88E-03	0.039	1.11	0.039	8.59E-03	0.228
5E-03	1.16	0.039	1.25E-04	0.040	1.13	0.041	2.54E-03	0.228
5E-02	1.20	0.039	5.03E-06	0.041	1.16	0.042	3.81E-04	0.228
1	1.22	0.040	5.33E-07	0.041	1.19	0.043	5.91E-05	0.229
<u>African Ancestry</u>								
5E-08	1.14	0.090	1.42E-01	0.031	1.15	0.099	1.62E-01	0.211
5E-07	1.24	0.086	1.22E-02	0.034	1.30	0.093	4.63E-03	0.215
5E-06	1.25	0.086	9.80E-03	0.034	1.30	0.092	3.95E-03	0.216
5E-05	1.23	0.086	1.71E-02	0.034	1.27	0.093	1.08E-02	0.214
5E-04	1.19	0.088	4.38E-02	0.032	1.17	0.094	9.82E-02	0.211
5E-03	1.18	0.090	6.74E-02	0.032	1.18	0.098	9.32E-02	0.211
5E-02	1.25	0.091	1.23E-02	0.034	1.17	0.097	1.07E-01	0.211
1	1.30	0.091	3.33E-03	0.036	1.26	0.097	1.56E-02	0.214

Average of the significant models (regression coefficient p-value < 0.05)

PRS	Average OR	Average R ²	Null R ² **	Incremental R ^{2***} over PRS null model
EA	1.13	0.044	0.004	0.041
MA	1.14	0.039	0.004	0.036
AA	1.25	0.034	0.004	0.030

PPRS	Average OR	Average R^2	PPRS Null R^2 **	Incremental R^{2***} over null model	Incremental R^{2***} over PPRS null model
EA	1.12	0.231	0.193	0.228 (61.5-fold)	0.038 (19.6%)
MA	1.13	0.228	0.201	0.224 (58.8-fold)	0.027 (13.2%)
AA	1.28	0.215	0.193	0.211 (57.0-fold)	0.021 (11.0%)

OR = odds ratio; SE = standard error; R^2 = psuedo- R^2

* PRS: PRS + basic covariates [Model(1) = PCOS ~ PRS + PC1-4 + Site]

* PPRS: PRS + PCOS component phenotypes + basic covariates [PPRS = PCOS ~ PRS + PC1-4 + Site + Hirsutism + Female Infertility + Irregular Menses]

** Null model: basic covariates only [Null Model = PCOS ~ PC1-4 + Site]

** PPRS Null model: PCOS component phenotypes + basic covariates [PPRS Null Model = PCOS ~ PC1-4 + Site + Hirsutism + Female Infertility + Irregular Menses]

*** Improvement rate: (Incremental change in pseudo- R^2 between the model with PRS/PPRS and the null model without PRS/PPRS) / (Pseudo- R^2 of the null model without PRS/PPRS)

Table 3: Average performance of PRS prediction algorithms in the female cohorts of European (n=33,869), Multiancestry (n= 49,365) and African (n=8,198) participants.

Summary - Average

PRS*	Accuracy	Sensitivity	Specificity	Balanced Accuracy***	AUC****
European (n=33,869)	0.551	0.547	0.755	0.651	0.715
Multiancestry (n= 49,365)	0.533	0.529	0.736	0.632	0.693
African (n=8,198)	0.496	0.494	0.590	0.542	0.543
PPRS**	Accuracy	Sensitivity	Specificity	Balanced Accuracy***	AUC****
European (n=33,869)	0.873	0.876	0.717	0.797	0.870
Multiancestry (n= 49,365)	0.881	0.886	0.640	0.763	0.823
African (n=8,198)	0.864	0.872	0.522	0.697	0.706

* PRS: PRS + basic covariates [PRS Model = PCOS ~ PRS + PC14 + Site]

** PPRS: PRS + PCOS component phenotypes + basic covariates [PPRS Model = PCOS ~ PRS + PC1-4 + Site + Hirsutism + Female Infertility + Irregular Menses]

*** Balanced Accuracy = (Sensitivity + Specificity)/2

**** AUC = Area Under the Curve




Fig 2. Stratification performance by quantile of PRS models, including PCs 1-4 and site as covariates, in (a) EA, (b) MA, and (c) AA populations. Group 1 includes those with the lowest PRS, and group 10 includes those with the highest. Bar colors indicate the average BMI in the quantile (darker indicates higher BMI), while the proportion of PCOS-diagnosed patients in each group is indicated at the top of each bar.

Table 4: Quantile analysis of PRS in the female European cohort (n=33,869) (PRS SNVs' p-value $<5 \times 10^{-8}$ and p-value ≤ 1 only).

	GROUP*	PCOS cases	PCOS prop**	Average BMI (kg/m ²)	Average PRS
PRS P < 5×10⁻⁸	1	45	1.3%	27.9	-1.750
	2	57	1.7%	27.9	-0.950
	3	54	1.6%	27.6	-0.813
	4	60	1.8%	28.1	-0.239
	5	61	1.8%	28.2	-0.068
	6	62	1.8%	28.0	0.014
	7	75	2.2%	27.6	0.248
	8	65	1.9%	28.1	0.810
	9	82	2.4%	27.9	0.952
	10	71	2.1%	27.8	1.790
...					
PRS P ≤ 1	1	50	1.5%	27.3	-1.790
	2	49	1.5%	27.5	-1.020
	3	61	1.8%	27.8	-0.654
	4	48	1.4%	27.9	-0.369
	5	58	1.7%	28.0	-0.113
	6	66	2.0%	27.8	0.132
	7	68	2.0%	28.0	0.386
	8	65	1.9%	28.1	0.672
	9	85	2.5%	28.4	1.040
	10	82	2.4%	28.4	1.720

PRS is adjusted with covariates and scaled for standardization.

* Higher group number indicates higher PRS

** Proportion of PCOS case patients in the quantile

The subsequent t-test reveals that PRS of case patients are significantly higher than the controls in all the nominally significant PRS models with regression p-value < 0.05, implying that higher genetic risk scores indicate higher occurrence of PCOS diagnosis (p-value= 2.15×10^{-4} , 7.75×10^{-4} , 2.43×10^{-4} , 2.51×10^{-5} , 3.12×10^{-5} in PRS model SNVs' p-value < 5×10^{-8} , 5×10^{-4} , 5×10^{-3} , 5×10^{-2} , 1 respectively) (**Supplementary Table 2**).

C. Performance improvement by the PRS variable

All the PRS models containing PCOS-PRS provided an improved fit over the null model by increasing the estimated explained sum of squares (pseudo- R^2) by McFadden's [37]. The average increase of pseudo- R^2 by the PRS variable in EA samples is 0.040, which is a 10-fold improvement ($=0.040/0.004$) over the null model. The ANOVA p-values of differentiating the PRS models from the null model are all under 1×10^{-31} , which validate the statistical significance of the performance improvement over the null model (**Table 2 and Supplementary Table 3**).

Evaluation of PRS in multi-ancestry and African ancestry participants

A. Predictive ability of each prediction model with different PRS

In the training set of the MA cohort (n=49,365 with 949 PCOS cases), the coefficient p-values of all PRS variables remain significant with positive beta coefficients (**Table 2; model1**). The average OR of PRS is 1.14 (average SE=0.038) and the average pseudo- R^2 value is 0.039, indicating that 3.9% of the phenotypic variance in the MA cohort could be explained by the PRS model. In the training set of AA participants (n=8,198 with 172 PCOS cases), the coefficient p-values of PRS variables remain

overall significant except for two PRS models of SNVs with p-value $< 5 \times 10^{-8}$ and p-value $< 5 \times 10^{-3}$ which may be due to the smaller sample size. Even though the regression p-values of the PRS variable do not show uniform performance in AA as compared to EA, the nominally significant PRS models generate a higher effect size in the AA samples compared to the other ancestry groups. The average OR of PRS models in the AA is 1.25 (SE=0.089), higher than both the EA (OR=1.13) and MA (OR=1.14). This is possibly due to the low RAF of PCOS risk variants in AA compared to EA (**Supplementary Table 1**).

For the testing dataset, PRS prediction displays an average 0.533 of accuracy, 0.529 of sensitivity, 0.736 of specificity with an average AUC of 0.693 in the multi-ancestry cohort. The out-of-sample performance in AA yielded an average AUC of 0.543 and showed an overall lower average accuracy (0.496), sensitivity (0.494) and specificity (0.590) compared to other ancestry groups (**Table 3**).

B. Stratification ability of each prediction model with different PRS

In the MA cohort, the proportion of PCOS patients increases from 1.5% in the lowest quantile to 2.6% in the highest quantile in the PRS calculation of SNVs with p-value ≤ 1 . The average BMI of the participants in the highest PRS quantile is 1.2 kg/m² higher (Cohen's d=0.17, t-test p-value= 1.62×10^{-13}) than the participants in the lowest PRS group (**Supplementary Table 4, Figure 2(b)**).

In the AA cohort, the number of PCOS patients does not always increase with higher PRS quantile, but the observation of an excess of PCOS patients in the highest PRS quantile is generally consistent across the models (**Figure 2c**). In the full-inclusive PRS model (SNVs with p-value ≤ 1), the rate of PCOS patients increases from 1.7% in

the lowest quantile to 3.1% in the highest PRS quantile (**Supplementary table 4**). The observed increase of the rate of PCOS patients is most pronounced in the PRS model with genome-wide significant variants (SNVs with $p\text{-value} < 5 \times 10^{-8}$), as the PCOS case rate doubles from 1.7% in the lowest quantile to 3.5% in the highest PRS quantile. We did not identify any notable trends in BMI in AA participants, which is depicted by the quantile color changes in **Figure 2(c)**.

An independent t-test confirms the significant differences of average PRS between PCOS cases and controls in MA across the models. The PRS difference between PCOS MA cases and controls is 0.165 after scaling with a full-inclusive PRS model, SNVs with $p\text{-value} \leq 1$ (Cohen's $d=0.201$, t-test $p\text{-value}=2.62 \times 10^{-6}$). In AA, only the full-inclusive PRS model shows statistically significant difference between PCOS cases and controls with a PRS difference of 0.175 (Cohen's $d=0.191$, t-test $p\text{-value}=2.90 \times 10^{-2}$) (**Supplementary Table 2**).

C. Performance improvement by the PRS variable

In the joint ancestry participants, all the prediction models containing the PRS variable provide a better fit over the null model by increasing the average pseudo- R^2 to 0.039, which is an 8.75-fold increase ($=0.035/0.004$) in explanatory power (**Table 2**). The subsequent ANOVA analysis confirms the statistical significance of the improved fits over the null model with all $p\text{-values} < 1 \times 10^{-46}$ (**Supplementary table 3**).

In the AA samples, the statistically significant PRS models show the average pseudo- R^2 of 0.034, which has the poorest fit among the ancestries. The models show an average pseudo- R^2 improvement of 7.5-fold increase ($=0.030/0.004$) from the null model without PRS (**Table 2**). Even with the lowest average incremental pseudo- R^2

(0.030) among the ancestries, the significant difference between the PRS models and the null model in Africans are confirmed with all ANOVA p-values $< 5 \times 10^{-3}$ (**Supplementary table 3**).

Development of PPRS prediction algorithms with PRS and PCOS component phenotypes

The addition of PCOS component EHR phenotypes to polygenic risk prediction significantly improved the predictive accuracy (**Table 2; model2 and Figure 3**). The average pseudo- R^2 of the PPRS is 0.231 in EA, 0.228 in MA, and 0.215 in AA samples, which indicates an average 14.7% improvement in pseudo- R^2 (19.6% in EA, 13.2% in AA, 11.0% in MA) over the PPRS null model by the inclusion of PCOS component phenotypes. Compared to the basic null model, the PPRS prediction boosts the average predictive performance (pseudo- R^2) by approximately 60 times (61.5-fold in EA, 58.8-fold in AA, 57.0-fold in MA) by the combinational use of PCOS component EHR phenotypes and PRS. Of note, the PRS variable's p-values in every PPRS model remain consistently valid in the MA samples (p-values $< 5 \times 10^{-3}$), whereas it was not always significant in AA or even EA samples. The ORs of the PRS and PPRS remain similar across the ancestries (**Figure 4**).

The subsequent ANOVA tested that all the pairs between PPRS and PPRS null models were statistically distinct across the cohorts and every PPRS model show the improved fit over the PPRS null model (**Supplementary Table 3**). The average ORs of irregular menstruation (ICD9 code 626.4, ICD10 code N92.6), female infertility (ICD9 code 627, ICD10 code N97.0) and hirsutism (ICD9 code 704.1, ICD10 code L68.0) for






Figure 3. Comparison of odds ratios (ORs) for the PRS and PPRS in (a) EA, (b) MA, and (c) AA cohorts, at different PRS/PPRS inclusion thresholds by GWAS p-value. The top row shows OR distributions for the PRS model, which adjusted for basic covariates [PRS Model = PCOS ~ PRS + PC1-4 + Site]. The bottom row shows OR distributions for the PPRS model which adjusted for the same basic covariates as well as PCOS EHR component phenotypes [PPRS Model = PCOS ~ PRS + PC1-4 + Site + Hirsutism + Female Infertility + Irregular Menses].


(a) European ancestry (EA)




(b) Multiancestry (MA)




(c) African ancestry (AA)



(d) European ancestry (EA)



(e) Multiancestry (MA)



(f) African ancestry (AA)




Figure 4. Comparison of Receiving Operating Curves (ROC) of the PPRS and PRS prediction models for PCOS diagnosis. The models with the genome-wide significant SNVs ($p\text{-value} < 5 \times 10^{-8}$) were evaluated in females of (a) EA, (b) MA, and (c) AA cohorts, along with the full-inclusive prediction models ($p\text{-value} < 1$) in females of (d) EA, (e) MA, and (f) AA cohorts. The areas under the curve (AUC) are provided in Table 2 and Supplementary Table 2. PRS model adjusted for basic covariates [PRS Model = PCOS ~ PRS + PC1-4 + Site], and PPRS model adjusted for the same basic covariates as well as PCOS EHR component phenotypes [PPRS Model = PCOS ~ PRS + PC1-4 + Site + Hirsutism + Female Infertility + Irregular Menses]. Null models only included the basic covariates without the PRS variable.

PCOS prediction were, as expected, strong across the cohorts: 5.49, 10.9, and 17.1, respectively (**Supplementary Table 5**).

Clinical phenome analysis


A. Associated phenotypes with PRS (*PheWAS-1*)

The general scheme of our PheWAS analyses are depicted in **Figure 5a**. Based on the model examination described above, the genome-wide PRS that includes all SNVs with $p\text{-value} \leq 1$ was selected as the best performing PRS model and used for phenome-wide analysis. The phenomes of 49,343 female participants and 41,669 male participants were analyzed separately to test for association with high genetic risk for PCOS.


In the female PheWAS with PRS, 75 EHR phenotypes were identified with phenome wide significance (**Figure 5b, Supplementary Table 6a**). ‘Morbid obesity’ (phecode 278.11) and obesity-related endocrine phenotypes, including ‘overweight, obesity, and other hyperalimentation’ (phecode 278), ‘type 2 diabetes’ (phecode 250.2), ‘essential hypertension’ (phecode 401.1) ‘hypercholesterolemia’ (phecode 272.11), ‘hypertension’ (phecode 401), ‘disorders of lipid metabolism’ (phecode 272) are the top-ranked. The phenome-wide significant association of ‘polycystic ovaries’ (phecode 256.4) and PCOS-PRS are observed with one of the largest effect sizes ($OR=1.015$) among the result.

As a complex endocrine disorder, the PCOS pathophysiology seems to be tightly linked to the expression of endocrine or circulatory system manifestation. Among the 75


(a)



(b)



(c)






Fig 5. PheWAS scheme and results using PRS. (a) PheWAS scheme and sample sizes; (b) PheWAS Manhattan plot of PRS (SNVs with $p\text{-value} \leq 1$); (c) PheWAS Manhattan plot of PRS (SNVs with $p\text{-value} < 5E-08$); (d) pie chart summarizing PheWAS groups. In Manhattan plots (b) and (c), the x-axis represents the EHR phenotype categorical group and the y-axis represents the negative $\log_{10}$ of the PheWAS $p\text{-value}$. Red lines indicate the cutoff for phenome-wide significance. For readability, only the most significant associations are annotated. Full lists of phenome-wide significant results are provided in Supplementary Tables 5 and 6, respectively. The pie chart in (d) shows EHR categories for the 72 phenome-wide significant phenotypes identified through PheWAS of the genome-wide PRS (SNVs with $p\text{-value} \leq 1$).

phenome-wide significant traits with PRS, the phenotypes of circulatory system (26.0%) and endocrine/metabolic system (21.0%) appeared the most frequently (**Figure 5d**), while the four highest associated phenotypes are all endocrine/metabolic features.

Among the remainder of the phenome-wide significant phenotypes, associations of musculoskeletal phenotypes like ‘osteoarthritis’ (phecode 740 and 740.9) or ‘calcaneal spur; Exostosis NOS’ (phecode 726.4) possibly imply the hormonal changes on the skeletal system impacted by PCOS epidemiology. Multiple symptomatic genitourinary phenotypes of PCOS were also identified: ‘abnormal mammogram’ (phecode 611.1) or ‘other signs and symptoms in breast’ (phecode 613.7). An obesity-related pulmonary disorder of ‘sleep apnea’ (phecode 327.3) is also observed (classified as neurological phenotype in phecode map) with ‘obstructive sleep apnea’ (phecode 327.32). We could not identify any psychological or depression related phenotype that is known to have genetic correlation with the hormonal changes of PCOS.

The overall low range of OR (1.004~1.010) of the PheWAS results should be noted, which is assumedly due to the aggregated effects of the low impact SNVs for PCOS, especially in the full-inclusive PRS with the entire GWAS SNVs. The ORs from the generic PheWAS of individual PCOS SNVs are observed to be higher before merging them into the cumulative PRS, which is described later (**Supplementary Table 7**).

In the replication analysis on an independent cohort of 18,096 EA females (BioVU), 16 out of 75 phenome-wide signals from the discovery analysis are replicated including ‘PCOS’ (p-value= 1.93×10^{-2} , phecode 256.4) with the positive OR of 1.174 (**Table 5a**). Half of the replicated phenotypes (8 out of 16) belong to the endocrine/metabolic category. In particular, the following obesity-related endocrine phenotypes exhibit strong

Table 5: (a) 16 significant phenotypes of PRS (SNVs' p-value ≤ 1) female-stratified PheWAS that were phenome-wide significant in the discovery cohort (n=49,343) and successfully replicated in the independent VU cohort (n=18,096). (b) 3 phenome-wide significant results of PCOS-PRS (SNPs with $P < 1$) male-stratified PheWAS from the discovery cohort (n=41,669) and replication cohort (n=15,612)

(a)			Discovery analysis					Replication analysis				
phecode	description	group	OR	SE	p	n_total	n_cases	OR	SE	p	n_total	n_cases
278.11	Morbid obesity	endocrine/metabolic	1.010	0.001	9.74E-18	37108	6790	1.116	0.029	1.64E-04	15329	1762
278.1	Obesity	endocrine/metabolic	1.008	0.001	4.14E-17	44267	13949	1.087	0.022	1.29E-04	17051	3484
278	Overweight, obesity and other hyperalimentation	endocrine/metabolic	1.007	0.001	2.20E-16	47803	17485	1.077	0.020	1.44E-04	18096	4529
250.2	Type 2 diabetes	endocrine/metabolic	1.007	0.001	8.18E-13	42874	10800	1.081	0.022	3.70E-04	16562	3660
327.3	Sleep apnea	neurological	1.008	0.001	4.71E-12	40673	6503	1.096	0.028	1.33E-03	15602	1847
250	Diabetes mellitus	endocrine/metabolic	1.007	0.001	5.39E-12	43325	11251	1.079	0.021	3.56E-04	16763	3861
571	Chronic liver disease and cirrhosis	digestive	1.008	0.001	4.17E-09	40531	4582	1.093	0.032	4.64E-03	15369	1463
539	Bariatric surgery	digestive	1.012	0.002	7.59E-09	47803	2034	1.202	0.055	8.00E-04	18096	439
327.32	Obstructive sleep apnea	neurological	1.007	0.001	1.16E-08	39291	5121	1.098	0.032	3.98E-03	15138	1383
571.5	Other chronic nonalcoholic liver disease	digestive	1.008	0.001	2.13E-08	40251	4302	1.112	0.033	1.38E-03	15219	1313
743.9	Osteopenia or other disorder of bone and cartilage	musculoskeletal	1.005	0.001	1.71E-07	43335	11354	0.956	0.022	4.45E-02	16019	3263
256.4	Polycystic ovaries	endocrine/metabolic	1.015	0.003	3.16E-07	40696	942	1.174	0.069	1.93E-02	15637	281
743	Osteoporosis, osteopenia and pathological fracture	musculoskeletal	1.004	0.001	6.38E-07	47803	15822	0.957	0.019	1.84E-02	18096	5340
250.3	Insulin pump user	endocrine/metabolic	1.008	0.002	2.25E-06	35057	2983	1.136	0.036	3.42E-04	14065	1163
250.23	Type 2 diabetes with ophthalmic manifestations	endocrine/metabolic	1.010	0.002	9.20E-06	33663	1589	1.221	0.062	1.20E-03	13272	370
627	Menopausal and postmenopausal disorders	genitourinary	1.004	0.001	1.40E-05	40468	14392	0.947	0.020	7.64E-03	16061	4301

(b)			<u>Discovery analysis</u>					<u>Replication analysis</u>				
phecode	description	group	OR	SE	p	n_total	n_cases	OR	SE	p	n_total	n_cases
278.11	Morbid obesity	endocrine/metabolic	1.009	0.002	5.93E-08	32456	3489	1.049	0.036	1.78E-01	13465	1082
250.2	Type 2 diabetes	endocrine/metabolic	1.005	0.001	1.41E-05	36835	10984	1.031	0.021	1.49E-01	14000	4198
250	Diabetes mellitus	endocrine/metabolic	1.005	0.001	2.47E-05	37199	11348	1.029	0.021	1.70E-01	14180	4378

Phenome-wide significant threshold: p-value < 2.9E-5

evidence of replication after multiple testing correction ($p\text{-value} < 6.7 \times 10^{-5}$, 0.05/75): ‘morbid obesity’ (phecode 278.11), ‘obesity’ (phecode 278.1), ‘overweight, obesity and other hyperalimentation’ (phecode 278). The well-known comorbidity between ‘type 2 diabetes’ (phecode 250.2) and PCOS is also identified along with other diabetic syndromes like ‘diabetes mellitus’ (phecode 250). Other notable replicated phenotypes included multiple neurological and digestive manifestations, which have well-known association to obesity, such as ‘chronic liver disease and cirrhosis’ (phecode 571), ‘bariatric surgery’ (phecode 539) and ‘other chronic nonalcoholic liver disease’ (phecode 571.5). An obesity-related pulmonary disorder of ‘sleep apnea’ (phecode 327.3) is also observed (classified as neurological phenotype in phecode map) with ‘obstructive sleep apnea’ (phecode 327.32).

In male-specific PheWAS with PRS (SNVs with $p\text{-value} \leq 1$) model, three metabolic phenotypes reached phenome-wide significance in the discovery analysis: ‘morbid obesity’ (phecode 278.11), ‘type 2 diabetes’ (phecode 250.2), ‘diabetes mellitus’ (phecode 250) which are known risk factors and/or co-morbidities for PCOS (**Figure 5b, Table 5b, Supplementary Table 6b**). However, none of the associations were replicated in the replication analysis on 15,611 independent males. It is possible that the replication sample remained underpowered and larger sample sizes will be needed to distinguish these results from a true null result.

B. Sensitivity analysis – Case-excluded analysis (PheWAS-2)

After removing 949 PCOS patients in PheWAS investigation, we still identified 68 PRS-phenotype associations that reached phenome-wide significance (**Supplementary**

table 8), which is not very different from PheWAS-1. The result might be due to the challenge of current diagnosis practices in identifying PCOS cases, which implies the control groups are not completely excluding PCOS patients and possibly include some mixed signals from the unidentified PCOS cases. Alternatively, it is possible that genetic risk for PCOS remains a robust risk factor for these phenotypes even in the absence of clinical manifestations of PCOS.

The representative signals of diabetes/obesity-related endocrine traits that are identified in PheWAS-1 remained significant: ‘morbid obesity’ (phecode 278.11), ‘type 2 diabetes’ (phecode 250.2), ‘obesity’ (phecode 278.1), ‘overweight, obesity and other hyperalimentation’ (phecode 278), ‘diabetes mellitus’ (phecode 250), ‘hypercholesterolemia’ (phecode 272.11), ‘disorders of lipid metabolism’ (phecode 272) and ‘hyperlipidemia’ (phecode 272.1) etc.

Four phenotypes no longer remained phenome-wide significant in PheWAS-2 compared to PheWAS-1, including ‘menopausal and postmenopausal disorders’ (phecode 627), ‘iron deficiency anemias, unspecified or not due to blood loss’ (phecode 280.1), ‘sleep disorders’ (phecode 327) and ‘Insomnia’ (phecode 327.4). A new metabolic phenotype of ‘disorders of fluid, electrolyte, and acid-base balance’ (phecode 276) was phenome-wide significance in PheWAS-2 compared to PheWAS-1, but the association did not remain significant in replication analysis. The phenome-wide significant phenotype with the largest effect size in PheWAS-2 is ‘localized adiposity’ (OR=1.014, phecode 278.3), same as for PheWAS-1. It should be of note that the range of OR is low in PRS-PheWAS due to the cumulative effect sum of all PCOS susceptibility loci including low-effect variants.

C. Sensitivity analysis – Associations with individual PCOS susceptibility loci **(PheWAS-3)**

In the individual PheWAS of 85 PCOS genome-wide significant variants, even though no association survives phenome-wide significance, likely due to the multiple testing burden, 11 PCOS variants show notable association to ‘polycystic ovaries’ across the ancestry groups (Most significant variant hg19 chr11:30226528, OR=1.36, phecode 256.4), ranked as the second most significant phenotype (**Supplementary table 7**). Out of top 100 associations in PheWAS-3, the largest number of associations were related to circulatory system for ‘thrombotic microangiopathy’ (31.0%). Endocrine/metabolic related phenotypes were the second most frequent category (21.0%) composed of either ‘PCOS’ or ‘ovarian dysfunction’, and 12% of the top associations were digestive traits, largely devoted to diverticular diseases. We did not identify any associations related to obesity or diabetes, which were the most significant phenotypic features found in PheWAS-1 and PheWAS-2.

Discussion

A key question in precision medicine is how to identify patients at high risk for a given disease for the goal of targeting preventive care. In this study, we examined the ability of PRS to predict PCOS clinical diagnosis and mine comorbid EHR phenotypes with the ultimate goal of improving diagnostic accuracy for PCOS. We show that a PRS

for PCOS can be used (a) to identify patients at elevated risk of PCOS and (b) to determine the comorbid or pleiotropic phenome-wide expression associated with PCOS in a clinical setting.

The primary accomplishment of this study is a systematic enhancement of the polygenic risk prediction by integration of additional disease component phenotypes in the EHR into a PPRS. The onset of hirsutism, menstrual dysfunction, or female infertility are representative symptoms of PCOS and essential in determining clinical hyperandrogenism [10, 40, 41]. They are not required for a diagnosis of PCOS per se, but are useful in suggesting PCOS in a clinical context. The PPRS significantly improves the average explanatory power (pseudo- R^2) of PCOS prediction by 0.221 (59.1-fold increase) compared to the null model without PRS or component phenotypes, and by 0.037 (14.7% increase) over the null model with the component phenotypes alone (**Table 2 and Figure 4**). In contrast to the previous studies that attempted to identify PCOS diagnosis with risk score calculation [13, 42], our algorithm did not limit risk predictor in a single-dimension, using both phenotype and genotype markers with polygenic inheritance, and extensively demonstrated the predictive performance of PPRS with several machine-learning techniques. The findings shown here strengthen the potential clinical utility of PPRS as a disease predictor, particularly when combined with component symptom information available within the EHR.

To date, research has consistently shown that the PRS built from EA GWAS data does not perform as robustly across non-EA samples. In this study, we assessed the performance of a Eurocentrically built PCOS-PRS on the samples of EA, AA, and the joint MA cohorts. Undeniably, validation statistics varied by ancestry group and the

PCOS diagnosis prediction in AA cohort shows the poorest performance. However, it is of note that more than half of the tested models in AA still show statistical significance in terms of regression p-value, and those models display a reliable efficiency for PCOS detection in effect size and AUC (**Table 3**). Interestingly, the ORs for PRS differ across the ancestry cohorts, and somewhat higher in some prediction models in AA (average OR of model1=1.25, model2=1.28) and MA samples (average OR of model1=1.14, model2=1.13) than EA samples (average OR of model1=1.13, model2=1.12). The overall ORs of the PRS variable are fairly stable throughout all polygenic prediction models (OR 1.12~1.28). The observed significance of the PRS variable in the MA cohort, more stable than in the EA or AA participants alone, is likely due to the increased statistical power with larger sample size that counters the sample heterogeneity introduced. In addition, we found that the accumulation of genetic variants did not always increase the predictive capability of PRS in terms of pseudo- R^2 and OR (**Figure 3, Table 2**). This might be due to the different RAF of PCOS risk variants by different PRS p-value cutoffs, and the varying LD structure of the ancestry groups. Previous research has confirmed that the LD pattern varies between EA and Chinese women at the PCOS susceptibility loci encoding LH/choriogonadotropin receptor (*LHCGR*) and FSH receptor (*FSHR*) genes, but the reproducible signals of the loci are consistently associated to PCOS regardless of ancestry[43, 44]. Our sensitivity analysis (PheWAS-3) also suggests the varying phenotypic effect of PCOS loci in different ancestries, but confirms the strong association with PCOS nonetheless. These findings demonstrate the primary role of PCOS-PRS in cumulatively explaining substantial

variation of disease susceptibility across ancestries even with differing LD structures, and extend the general utility of PPRS in disease prediction.

Furthermore, our PRS-based phenome-wide analysis revealed several clinical associations that are tightly linked with obesity, confirming the shared metabolic pathways between PCOS and obesity in a phenomic aspect. As obesity is a common finding which can be found in 50-65% of PCOS patients[10], and previous Mendelian randomization study revealed the causal relationship of BMI on PCOS etiology[45], many of our findings could be interpreted as phenotypic evidence of co-morbid obesity. 'Morbid obesity' (phecode 278.11), 'hypercholesterolemia' (phecode 272.11), 'disorders of lipid metabolism' (phecode 272), 'hyperlipidemia' (phecode 272.1), 'hypertension' (phecode 401) or 'abnormal glucose' (phecode 250.4) are easily understandable with the context of heightened metabolic risks for obesity. 'Sleep apnea' (phecode 327.3) and 'chronic liver disease and cirrhosis' (phecode 571), 'GERD' (phecode 530.11), 'diseases of esophagus' (phecode 530 and 530.1) are either neurological, pulmonary or digestive assorted symptoms that are commonly found in the patients with obesity.

It is also noteworthy that there were 75 significant associations identified in women while in men, there were only three significantly associated diagnosis (morbid obesity, type 2 diabetes, diabetes mellitus) despite a similar sample size for males and females in the analysis. It is possible that the clinical consequences of high androgens in males are less likely to cause symptoms for which medical treatment is sought, or that these genetic variants only elevate androgen levels in a female 'environment' but not a male one. The three identified phenotypes in males additionally suggest that if an individual

harbors high genetic risk for PCOS, the metabolic manifestations are similar regardless of sex.

Consistent with previous studies [13, 45], we identified phenotypic evidence of positive BMI association with genetic risk of PCOS. In the stratification analysis of PRS, our observation of the increased BMI in individuals with high risk of PCOS are evident in both EA and MA cohorts (**Figure 2**). The comorbid phenotypes could be driven by pleiotropy in which PCOS-associated genes also increase BMI, or could be due to under diagnosis of PCOS itself, in which case the association with obesity phenotypes may be a result of comorbidity with undiagnosed PCOS.

Several limitations to this study need to be acknowledged. First, the sample size of AA participants was relatively small which increases the likelihood of both false negative and false positive findings. Further investigation is needed to fully understand the overlap in PCOS genetic factors across multi-ancestry participants and the methodological application of Eurocentric PCOS-PRS to other genetic ancestries considering LD structure. Secondly, the phenotypic components we used for polygenic prediction are currently limited to only three representative phenotypes: hirsutism, irregular menstruation, and female infertility. Fueled by our PheWAS finding, the work could be extended by incorporating the additional phenotypes that might increase the likelihood of an eventual diagnosis. Also, the phecode of PCOS used for PheWAS was converted from ICD-9-CM 256.4 and ICD-10-CM E28.2, which was used as a proxy for capturing PCOS in the EMR. This phecode may not perfectly capture PCOS as they may or may not capture hyperandrogenemia. The selection bias in our discovery cohort should be acknowledged as well. Two of our participating sites (Geisinger and

Marshfield) mainly recruited their patients for the study of obesity and type 2 diabetes, which resulted in a higher proportion of obese patients into their biobank and therefore may inflate the prevalence of PCOS in these subgroups. Lastly, due to the low diagnosis rate of PCOS patients in current EHR system, it is possible that unidentified PCOS cases could reduce power in each analysis.

Our approach has provided a novel methodological opportunity to stratify patients' genetic risk and to discover the phenomic network associated with PCOS pathogenesis. Integrative analysis of the PRS-PheWAS enables the systematic interrogation of PCOS comorbidity patterns across the phenome, which cannot be readily identified by a single-variant approach. The identified phenomic networks could be used at the stage of first screening, prior to the testing of hormones or imaging of ovaries, or to help the patient and physician decide whether more extensive testing would be useful for PCOS diagnosis. From a precision medicine perspective, such an approach may provide a greater understanding of a patient's clinical presentation and suspected diagnosis based on phenotypic or genetic variations.

Reference

1. Davies N. PCOS: Polycystic Ovarian Syndrome. *Diabetes Self Manag.* 2016;33(1):44-7.
2. Azziz R, Marin C, Hoq L, Badamgarav E, Song P. Health care-related economic burden of the polycystic ovary syndrome during the reproductive life span. *J Clin Endocrinol Metab.* 2005;90(8):4650-8.
3. Society TE. *Endocrine Facts and Figures: Reproduction and Development.* 2017.
4. Yawn V. Polycystic ovarian syndrome. *Adv NPs PAs.* 2012;3(12):11-4; quiz 5.
5. Vink JM, Sadrzadeh S, Lambalk CB, Boomsma DI. Heritability of polycystic ovary syndrome in a Dutch twin-family study. *J Clin Endocrinol Metab.* 2006;91(6):2100-4.
6. Jahanfar S, Eden JA, Nguyen T, Wang XL, Wilcken DE. A twin study of polycystic ovary syndrome and lipids. *Gynecol Endocrinol.* 1997;11(2):111-7.
7. Jahanfar S, Eden JA, Warren P, Seppala M, Nguyen TV. A twin study of polycystic ovary syndrome. *Fertil Steril.* 1995;63(3):478-86.
8. Broekmans FJ, Knauff EA, Valkenburg O, Laven JS, Eijkemans MJ, Fauser BC. PCOS according to the Rotterdam consensus criteria: Change in prevalence among WHO-II anovulation and association with metabolic factors. *BJOG.* 2006;113(10):1210-7.
9. Li L, Baek KH. Molecular genetics of polycystic ovary syndrome: an update. *Curr Mol Med.* 2015;15(4):331-42.
10. Futterweit W. Polycystic ovary syndrome: clinical perspectives and management. *Obstet Gynecol Surv.* 1999;54(6):403-13.
11. Wolf WM, Wattick RA, Kinkade ON, Olfert MD. Geographical Prevalence of Polycystic Ovary Syndrome as Determined by Region and Race/Ethnicity. *Int J Env Res Pub He.* 2018;15(11).
12. Carmina E. Diagnosis of polycystic ovary syndrome: from NIH criteria to ESHRE-ASRM guidelines. *Minerva Ginecol.* 2004;56(1):1-6.
13. Day F, Karaderi T, Jones MR, Meun C, He C, Drong A, et al. Large-scale genome-wide meta-analysis of polycystic ovary syndrome suggests shared genetic architecture for different diagnosis criteria. *PLoS Genet.* 2018;14(12):e1007813.
14. Dewailly D. Diagnostic criteria for PCOS: Is there a need for a rethink? *Best Pract Res Clin Obstet Gynaecol.* 2016;37:5-11.
15. Agapova SE, Cameo T, Sopher AB, Oberfield SE. Diagnosis and challenges of polycystic ovary syndrome in adolescence. *Semin Reprod Med.* 2014;32(3):194-201.
16. Fritsche LG, Gruber SB, Wu Z, Schmidt EM, Zawistowski M, Moser SE, et al. Association of Polygenic Risk Scores for Multiple Cancers in a Phenome-wide Study: Results from The Michigan Genomics Initiative. *Am J Hum Genet.* 2018;102(6):1048-61.
17. International Schizophrenia C, Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, et al. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature.* 2009;460(7256):748-52.

18. Khera AV, Chaffin M, Aragam KG, Haas ME, Roselli C, Choi SH, et al. Genome-wide polygenic scores for common diseases identify individuals with risk equivalent to monogenic mutations. *Nat Genet.* 2018;50(9):1219-24.
19. Zheutlin AB, Dennis J, Restrepo N, Straub P, Ruderfer D, Castro VM, et al. Penetrance and pleiotropy of polygenic risk scores for schizophrenia in 90,000 patients across three healthcare systems. *bioRxiv.* 2018:421164.
20. Carroll RJ, Bastarache L, Denny JC. R PheWAS: data analysis and plotting tools for phenome-wide association studies in the R environment. *Bioinformatics.* 2014;30(16):2375-6.
21. Denny JC, Ritchie MD, Basford MA, Pulley JM, Bastarache L, Brown-Gentry K, et al. PheWAS: demonstrating the feasibility of a phenome-wide scan to discover gene-disease associations. *Bioinformatics.* 2010;26(9):1205-10.
22. Martin AR, Kanai M, Kamatani Y, Okada Y, Neale BM, Daly MJ. Hidden 'risk' in polygenic scores: clinical use today could exacerbate health disparities. *bioRxiv.* 2019:441261.
23. Rosenberg NA, Edge MD, Pritchard JK, Feldman MW. Interpreting polygenic scores, polygenic adaptation, and human phenotypic differences. *Evolution, Medicine, and Public Health.* 2018:eoy036-eoy.
24. Duncan L, Shen H, Gelaye B, Ressler K, Feldman M, Peterson R, et al. Analysis of Polygenic Score Usage and Performance in Diverse Human Populations. *bioRxiv.* 2018:398396.
25. Martin AR, Kanai M, Kamatani Y, Okada Y, Neale BM, Daly MJ. Hidden 'risk' in polygenic scores: clinical use today could exacerbate health disparities. *bioRxiv.* 2018:441261.
26. Martin AR, Gignoux CR, Walters RK, Wojcik GL, Neale BM, Gravel S, et al. Human Demographic History Impacts Genetic Risk Prediction across Diverse Populations. *Am J Hum Genet.* 2017;100(4):635-49.
27. Curtis D. Polygenic risk score for schizophrenia is more strongly associated with ancestry than with schizophrenia. *bioRxiv.* 2018:287136.
28. McCarty CA, Chisholm RL, Chute CG, Kullo IJ, Jarvik GP, Larson EB, et al. The eMERGE Network: a consortium of biorepositories linked to electronic medical records data for conducting genomic studies. *BMC Med Genomics.* 2011;4:13.
29. Stanaway IB, Hall TO, Rosenthal EA, Palmer M, Naranbhai V, Knevel R, et al. The eMERGE genotype set of 83,717 subjects imputed to ~40 million variants genome wide and association with the herpes zoster medical record phenotype. *Genet Epidemiol.* 2019;43(1):63-81.
30. Kho AN, Pacheco JA, Peissig PL, Rasmussen L, Newton KM, Weston N, et al. Electronic medical records for genetic research: results of the eMERGE consortium. *Sci Transl Med.* 2011;3(79):79re1.
31. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet.* 2007;81(3):559-75.
32. Euesden J, Lewis CM, O'Reilly PF. PRSice: Polygenic Risk Score software. *Bioinformatics.* 2015;31(9):1466-8.

33. Denny JC, Bastarache L, Ritchie MD, Carroll RJ, Zink R, Mosley JD, et al. Systematic comparison of phenome-wide association study of electronic medical record data and genome-wide association study data. *Nat Biotechnol.* 2013;31(12):1102-10.
34. Wu P, Gifford A, Meng X, Li X, Campbell H, Varley T, et al. Developing and Evaluating Mappings of ICD-10 and ICD-10-CM codes to Phecodes. *bioRxiv.* 2018:462077.
35. Das S, Forer L, Schonherr S, Sidore C, Locke AE, Kwong A, et al. Next-generation genotype imputation service and methods. *Nat Genet.* 2016;48(10):1284-7.
36. McCarthy S, Das S, Kretzschmar W, Delaneau O, Wood AR, Teumer A, et al. A reference panel of 64,976 haplotypes for genotype imputation. *Nat Genet.* 2016;48(10):1279-83.
37. McFadden D. Conditional logit analysis of qualitative choice behavior. *Frontiers in Econometrics.* 1973:105-42.
38. Plomin R, Haworth CM, Davis OS. Common disorders are quantitative traits. *Nat Rev Genet.* 2009;10(12):872-8.
39. Krapohl E, Euesden J, Zabaneh D, Pingault JB, Rimfeld K, von Stumm S, et al. Phenome-wide analysis of genome-wide polygenic scores. *Mol Psychiatry.* 2016;21(9):1188-93.
40. Goodman NF, Cobin RH, Futterweit W, Glueck JS, Legro RS, Carmina E, et al. American Association of Clinical Endocrinologists, American College of Endocrinology, and Androgen Excess and Pcos Society Disease State Clinical Review: Guide to the Best Practices in the Evaluation and Treatment of Polycystic Ovary Syndrome--Part 1. *Endocr Pract.* 2015;21(11):1291-300.
41. Rosenfield RL, Lucky AW. Acne, hirsutism, and alopecia in adolescent girls. Clinical expressions of androgen excess. *Endocrinol Metab Clin North Am.* 1993;22(3):507-32.
42. Deshmukh H, Papageorgiou M, Kilpatrick ES, Atkin SL, Sathyapalan T. Development of a novel risk prediction and risk stratification score for polycystic ovary syndrome. *Clin Endocrinol (Oxf).* 2019;90(1):162-9.
43. Mutharasan P, Galdones E, Penalver Bernabe B, Garcia OA, Jafari N, Shea LD, et al. Evidence for chromosome 2p16.3 polycystic ovary syndrome susceptibility locus in affected women of European ancestry. *J Clin Endocrinol Metab.* 2013;98(1):E185-90.
44. Chen ZJ, Zhao H, He L, Shi Y, Qin Y, Shi Y, et al. Genome-wide association study identifies susceptibility loci for polycystic ovary syndrome on chromosome 2p16.3, 2p21 and 9q33.3. *Nat Genet.* 2011;43(1):55-9.
45. Day FR, Hinds DA, Tung JY, Stolk L, Styrkarsdottir U, Saxena R, et al. Causal mechanisms and balancing selection inferred from genetic associations with polycystic ovary syndrome. *Nat Commun.* 2015;6:8464.

Acknowledgements

The phase III of the eMERGE Network was initiated and funded by the NHGRI through the following grants: U01HG008657 (Kaiser Permanente Washington/University of Washington School of Medicine); U01HG008685 (Brigham and Women's Hospital); U01HG008672 (Vanderbilt University Medical Center); U01HG008666 (Cincinnati Children's Hospital Medical Center); U01HG006379 (Mayo Clinic); U01HG008679 (Geisinger Clinic); U01HG008680 (Columbia University Health Sciences); U01HG008684 (Children's Hospital of Philadelphia); U01HG008673 (Northwestern University); U01HG008701 (Vanderbilt University Medical Center serving as the Coordinating Center); U01HG008676 (Partners Healthcare/Broad Institute); and U01HG008664 (Baylor College of Medicine).

The International PCOS Consortium members

Felix Day, Tugce Karaderi, Michelle R. Jones, Cindy Meun, Chunyan He, Alex Drong, Peter Kraft, Nan Lin, Hongyan Huang, Linda Broer, Reedik Magi, Richa Saxena, Triin Laisk-Podar, Margrit Urbanek, M. Geoffrey Hayes, Gudmar Thorleifsson, Juan Fernandez-Tajes, Anubha Mahajan, Benjamin H. Mullin, Bronwyn G.A. Stuckey, Timothy D. Spector, Scott G. Wilson, Mark O. Goodarzi, Lea Davis, Barbara Obermeyer-Pietsch, André G. Uitterlinden, Verner Anttila, Benjamin M Neale, Marjo-Riitta Jarvelin, Bart Fauser, Irina Kowalska, Jenny A. Visser, Marianne Anderson, Ken Ong, Elisabet Stener-Victorin, David Ehrmann, Richard S. Legro, Andres Salumets, Mark I. McCarthy, Laure Morin-Papunen, Unnur Thorsteinsdottir, Kari Stefansson,

Unnur Styrkarsdottir, John Perry, Andrea Dunaif, Joop Laven, Steve Franks, Cecilia M. Lindgren, Corrine K. Welt

Authors' contributions

YYJ, LD and MGH designed the study; IBS and DRC imputed and quality controlled the genotype array data missing variants with input from GPJ; JAP, AOB, RC, DRC, JCD, DRVE, HH, JBH, SJH, KH, GPJ, FDM, SP, MDR, IBS contributed to eMERGE genotype and phenotype data generation; LD, MGH, FD, MJ, TK, CM generated PCOS GWAS data through the International PCOS consortium; YYJ performed statistical analysis in discovery cohort and validated the algorithms; KA performed statistical analysis in replication cohort; YYJ, ANK, LD and MGH interpreted the results; YYJ, KA, JAP, ANK, LD, MGH drafted the manuscript; YYJ designed the figures and created the tables; All authors critically reviewed the manuscript for important intellectual content; DRC, GPJ, MS and RLC obtained the funding.

Competing interest statement

The authors report no competing interests.