

A machine learning and network framework to discover new indications for small molecules

Coryandar Gilvary^{*1,2,3,4}, Jamal Elkhader^{*1,2,3,4}, Neel Madhukar⁵, Claire Henchcliffe⁶, Marcus D. Goncalves^{3,7}, Olivier Elemento^{1,2,3,4,5,8}

¹ HRH Prince Alwaleed Bin Talal Bin Abdulaziz Alsaud Institute for Computational Biomedicine, Dept. of Physiology and Biophysics, Weill Cornell Medicine, New York, NY 10065, USA;

² Caryl and Israel Englander Institute for Precision Medicine, Weill Cornell Medicine, New York, NY 10065, USA;

³ Sandra and Edward Meyer Cancer Center, Weill Cornell Medicine, New York, NY 10065, USA;

⁴ Tri-Institutional Training Program in Computational Biology and Medicine, New York, NY 10065, USA;

⁵ OneThree Biotech, New York, NY 10021, USA

⁶ Department of Neurology, Weill Cornell Medicine, New York, NY 10065

⁷ Division of Endocrinology, Department of Medicine, Weill Cornell Medicine, New York, NY 10065

⁸ WorldQuant Initiative for Quantitative Prediction, Weill Cornell Medicine, New York, NY 10065, USA;

*These authors contributed equally

Correspondence: ole2001@med.cornell.edu (Olivier Elemento)

Abstract

Drug repurposing, identifying novel indications for drugs, bypasses common drug development pitfalls to ultimately deliver therapies to patients faster. However, most repurposing discoveries have been led by anecdotal observations (e.g. Viagra) or experimental-based repurposing screens, which are costly, time-consuming, and imprecise. Recently, more systematic computational approaches have been proposed, however these rely on utilizing the information from the diseases a drug is already approved to treat. This inherently limits the algorithms, making them unusable for investigational molecules. Here, we present a computational approach to drug repurposing, CATNIP, that requires only biological and chemical information of a molecule. CATNIP is trained with 2,576 diverse small molecules and uses 16 different drug similarity features, such as structural, target, or pathway based similarity. This model obtains significant predictive power (AUC = 0.841). Using our model, we created a repurposing network to identify broad scale repurposing opportunities between drug types. By exploiting this network, we identified literature-supported repurposing candidates, such as the use of systemic hormonal preparations for the treatment of respiratory illnesses. Furthermore, we demonstrated that we can use our approach to identify novel uses for defined drug classes. We found that adrenergic uptake inhibitors, specifically amitriptyline and trimipramine, could be potential therapies for Parkinson's disease. Additionally, using CATNIP, we predicted the kinase inhibitor, vandetanib, as a possible treatment for Type 2 Diabetes. Overall, this systematic approach to drug repurposing lays the groundwork to streamline future drug development efforts.

Introduction

With over \$800 million spent bringing a single drug to market over the course of 15 years, drug development has remained a costly and time-consuming affair¹. In response, there has been an increase in interest in drug repurposing, the identification of novel indications for known, safe drugs. Successes in this area have been seen in the past, most notably in sildenafil (e.g. Viagra), which was originally intended to treat hypertension and angina pectoris but was later repurposed to treat erectile dysfunction. Other examples of compounds repurposed for new therapeutic applications include minoxidil² and raloxifene³, which are now used to treat androgenic alopecia and osteoporosis, respectively. However, most of these repurposing opportunities were discovered through inefficient approaches including anecdotal observations or hypothesis-driven investigations, and a more efficient approach could lead to many more repurposing opportunities.

Computational approaches for repurposing drugs are appealing in that they can be systematically and quickly applied to many drugs at a low cost compared to their experimental counterparts. One computational approach that has proven to be invaluable in other areas of the drug development pipeline is machine learning. Machine learning is the use of computational algorithms to learn from available data to make novel predictions and gain new insight. Using this technique, one can create unbiased algorithms to match seemingly disparate drugs by comparing their common features⁴, such as clinical indication, toxicity profile⁵ or therapeutic target^{6,7}. Previously, our lab used a ‘similarity’ approach, leveraging the principle that similar drugs tend to have similar characteristics, to predict a drug’s target by investigating the known targets of other drugs that were predicted to be “similar” to the investigated drug based on shared features⁶. We found that DRD2, a dopamine receptor, was the predicted target for the compound ONC201. After identifying and experimentally validating this target, clinical trials were shifted to focus on gliomas, which are now successfully completing phase two trials at the time of this publication⁸. The approach of

leveraging drug similarity could immensely aid drug repurposing efforts with the appropriate data.

Others have successfully used this ‘similarity’ approach to repurpose drugs and demonstrated high predictive power when tested against FDA approved drug-diseases⁹. However, these methods have primarily linked drugs together using a disease-centric approach instead of using features related to the drug itself (i.e. drug-centric). These repurposing opportunities are identified by predicting diseases similar to the diseases a drug is already known to treat. Disease similarities can be based on semantic, pathophysiological, or clinical similarities related to the drug’s clinical indication. For example, PREDICT, a repurposing method developed by Gottlieb et al.¹⁰, exploits the semantic similarity of disease terms as a form of disease-disease similarity. Such approaches, while reliable, limit the scope of the repositioning effort in several ways. First, the vast majority of small molecules never reach clinical approval and would be overlooked in this type of analysis. Second, the use of a disease-centric approach biases repurposing predictions toward exclusively similar clinical diseases (i.e.: cancer drugs to other cancer types)¹¹. We postulated that using solely drug information, such as chemical and biological features, would be a more effective and broader approach to drug repurposing.

Here, we propose a novel approach to drug repurposing, which operates by a platform we call, **Creating A Translational Network for Indication Prediction** (CATNIP). CATNIP is a machine-learning algorithm that learns to predict whether two molecules share an indication based solely on the drug’s chemical and biological features, using 2,576 unique drugs. The systematic application of CATNIP to molecule pairs creates a network with ~4.6 million nodes that can then be used to identify potential drug repurposing opportunities. Because CATNIP uses chemical structure and targets as key features, it can effectively bridge between different therapeutic indications. In this report, we have identified various candidate drug classes that are predicted to have therapeutic activity

outside of their intended indication in diseases such as Parkinson's disease and Type 2 Diabetes.

Results

Variance in drug indication nomenclature can be standardized

We collected a wide variety of drugs (N=3,066, including both approved and investigational molecules) with a diverse set of indications to ensure that our drug network covered a large portion of the known chemical space. A subset of these drugs (2,576 FDA approved drugs and 2,492 indications taken from DrugBank¹²) were used as a gold-standard of drug-indication associations in the training set for the model. Disease names are often not standardized, which can lead to many diverse names for the same disease. This problem leads to many drug pairs appearing to not have shared indications, when they are associated with two different names for the same disease. To address inconsistencies in nomenclature for drug indications, such as “prostate carcinoma” and “carcinoma of the prostate”, the MetaMap tool¹³ was applied (**Methods**). Using MetaMap, we clustered the 2,492 DrugBank indications into 1,042 standardized indications. A multitude of indication types were included in this standardization including, but not limited to, oncological, mental health, and neurological diseases (**Figure S1A**). Our rigorous standardization of drug indications ensured an accurate training set, allowing for the discovery and modeling of drug-indication relationships.

Drug pairs sharing indications have other similar characteristics

We hypothesized that pairs of drugs that shared at least one indication would have other similar drug characteristics (**Table S1**). To test this hypothesis, we integrated the similarity of two drugs across chemical and biological drug properties, and created a computational model to predict if two drugs will share an indication (**Figure 1**). All 16 of the drug similarity features (**Table S1**) collected could significantly distinguish between drug pairs known to share an indication

and those not known to share an indication (**Figure S2-5**). For example, we found that drug pairs with a shared clinical indication, according to their listed DrugBank indications, tended to have significant overlap in targets (D-statistic = 0.168, p-value < 0.001, **Figure S2A**). The feature which best discriminated between drug pairs that shared a clinical indication versus drug pairs that do not was the similarity between the KEGG pathways that each drug's targets are involved in (D-statistic = 0.241, p < 0.001, **Figure S4C**). Pathway similarity was calculated as the Jaccard Index between the KEGG pathways that contain each drug's gene targets (**Methods**). The difference in effect size between the target similarity and the pathway similarity (D-statistic= 0.168 vs 0.241, respectively) indicates that the drugs do not necessarily have to target the same exact genes, but rather the same biological pathway, in order to share a clinical indication. Additionally, we found that drug pairs that share an indication had a more similar chemical structure than drug pairs that did not share an indication (D-statistic = 0.105, p-value < 0.001, **Figure S5A**). Overall, these features seem to indicate sufficient power in differentiating drugs that share and do not share indications, which we hypothesized can then be leveraged to create a predictive model.

Drug pairs that share indications can be predicted by model

Using these diverse drug properties as features we trained a Gradient Boosting model to predict if two drugs share a clinical indication. A Gradient Boosting model showed superior results when compared with other algorithms (**Methods, Table S2**). The model output is a drug similarity score (hereby referred to as a "CATNIP score"), which allows us to classify drug pairs that share clinical indications. We performed a 5-fold cross-validation analysis and achieved significant predictive performance with an area-under-the-receiver-operator curve (AUC) of 0.841 (**Figure 2A**). We confirmed the statistical significance of our model with a precision-recall curve (PRC) because of the class imbalance in our dataset between drug pairs that share indications against those that do not (23,840 Shared, 1,299,623 Not Shared). When compared to random predictions,

our model showed significant improvement (0.189 vs 0.0184 area-under PRC, **Figure S6**).

It has been shown before that structurally similar drugs have a high probability of treating the same indication¹⁵. However, there are many examples of drug pairs that defy this rule. For example, tamoxifen¹⁶ and anastrozole¹⁷ are structurally dissimilar compounds (Dice similarity = 0.372) that treat the same indication (Metathesaurus term: Cancer, Breast). To ensure that our model could accurately classify drug pairs that share an indication but are not structurally similar, we recalculated all performance metrics to control for high and low structural similarity. High performance was retained under both of these conditions (high structural similarity AUC = 0.885, low structural similarity AUC = 0.828 AUC, **Figure 2A**). These performance metrics confirm that our model is robust enough to predict if a drug pair will share an indication with or without structural similarity.

Network clusters identify drugs with similar clinical characteristics

We constructed a repurposing network by calculating a CATNIP score for all possible drug pairs found within DrugBank, and assigning the drugs as nodes and the CATNIP score as the edge weight. We pruned the network using a cut-off value of 7.4 for the CATNIP scores (**Figure 2B**), which included 792 different drug pairs. This cut-off is equivalent to a predicted probability of >99% to share an indication and allowed for a balance between confidence within our predictions and drug diversity and availability.

We hypothesized that drugs sharing at least one indication would cluster together in our network. To confirm this theory, we classified each drug per its 1st order Anatomical Therapeutic Chemical (ATC) classification. This identification is a method of distinguishing the clinical use of a drug that is widely used in European and North American chemoinformatics databases¹⁸. Using ATC, we observed clearly defined clusters within the repurposing network (**Figure 2B**). Many

clusters featured multiple ATC classifications, suggesting potential repurposing opportunities. For example, one cluster included the thiazolidinediones, rosiglitazone and pioglitazone (ATC classification: ‘Alimentary Tract and Metabolism’) and the fibrates, fenofibrate and bezafibrate (ATC classification: ‘Cardiovascular system’). These two clustered ATC classifications were connected by a high (7.42) CATNIP score between bezafibrate and pioglitazone, an antidiabetic drug; a relationship driven by the shared targeting of PPAR α and PPAR γ resulting in the improvement of lipid and glucose metabolism. Bezafibrate has shown efficacy in the treatment of Type 2 Diabetes in numerous retrospective and pre-clinical studies, including Phase 2 trials¹⁹⁻²¹, however is still not an approved antidiabetic. The identification of bezafibrate as a potential diabetes treatment is a key example of how CATNIP can be used to identify repurposing opportunities.

We reasoned that the connections between ATC classifications across all the drug clusters could provide additional aid for drug repurposing purposes. Using the pruned network (CATNIP Score > 7.4), we collected all the scores between drugs of differing ATC classifications. From this collection, we were able to determine the median score associated between each pair of ATC classifications. The ATC classifications with the highest median CATNIP scores had literature support for numerous repurposing efforts between them (**Table 1**). For example, drugs with the ATC classifications of “Respiratory System” and “Systemic Hormonal Preparations, excluding sex hormones and insulins” were strongly connected to each other (7.97 median CATNIP score). This connection was driven by highly scored pairs of drugs including rimexolone to mometasone (8.31 CATNIP score) and prednisone to triamcinolone (8.13 CATNIP score). These connections are supported by the fact that hormonal agents like glucocorticoids and beta adrenergic agonists have been used for decades to relax the airway musculature in patients with reactive airways disease and chronic obstructive pulmonary disease²². Interestingly, our analysis identified glucagon, a peptide hormone that increases blood glucose levels, as a candidate

for “Respiratory System” repurposing and this use already has clinical support^{23,24}. Additionally, drugs classified as “Respiratory System” and “Dermatological” were also observed to be highly associated because of interactions such as the one between ciclesonide and hydrocortisone (8.43 CATNIP score). Ciclesonide and hydrocortisone do in fact share a clinical indication, “Asthma Bronchial”, giving added confidence to our findings. These types of network observations are important in laying the groundwork for suggesting novel clinical repurposing strategies for FDA-approved drugs.

CATNIP identifies novel disease areas for drug classes

We investigated the ability to leverage CATNIP scores to identify repurposing opportunities by evaluating specific drug classes. Drug classes are predefined in DrugBank. In order to identify actionable repurposing possibilities, we narrowed this list down to 50 classes containing inhibitors, antagonists, or agonists of specific gene or protein families. We focused our attention on specific disease areas that are attractive for drug repurposing opportunities, due to a lack of current treatments or high rates of acquired resistance. The specific disease areas were: “mental disorders”, “neurological diseases”, “diabetes”, and “cancer” (cancer was further divided into specific cancer types due to the large variance in disease pathology between types, **Methods**). We hypothesized that CATNIP scores could be used to identify specific drug classes that would be efficacious for a new disease area. For each drug class and disease area, we found the statistical difference in the CATNIP score distribution between two sets of drug pairs. The first set included pairs that had one drug within the drug class and the other drug approved for the disease in question, while the other set included drug pairs that had one drug within the drug class and the other drug not approved for the disease in question (**Methods**). We compared the effect size, estimated by the Wilcoxon location shift, for all drug class-disease pairs that had a significant difference in distribution compared to drug class-non-disease pairs (FDR < 0.1, **Figure 3A-B, Figure S7-8**). By using CATNIP scores, we found that many well-known drug class-diseases associations could be recovered. For example,

“muscarinic antagonists” were highly ranked for “neurological diseases” and many such agents are FDA-approved for this indication²⁵. In addition, we found that “kinase inhibitors” were closely associated with the treatment of cancer and “dopamine antagonists” for the treatment of “mental disorders”^{26, 27} (Wilcoxon Location Shift = 0.711-0.945 for “kinase inhibitors” and select cancer types, Location Shift = 0.882 for “dopamine antagonists” and “mental disorders”, p-value < 0.001, **Figure S9**). In fact, almost all drug class-disease associations contained at least one FDA-approved drug for the respective disease, giving us added confidence in our model. Of note, each drug was allowed to be categorized into numerous drug classes, leading to unexpected, yet easily explained, results; for example, “dopamine antagonists” appearing as a top drug class for “neurological diseases”. This is due to risperidone, a drug traditionally used for schizophrenia and mood disorders, also having a secondary indication of Alzheimer’s type severe dementia.

Next, we further interrogated the drug classes associated with “neurological diseases” and “diabetes”, specifically. CATNIP scores were able to correctly identify almost all drug classes known to treat these diseases (**Figure 3A-B**). To identify possible repurposing candidates, we focused our attention on drug classes shown to have a large positive effect size with this CATNIP analysis but are not currently approved for treatment. For “neurological diseases”, the use of adrenergic uptake inhibitors, traditionally used as antidepressants, was the top repurposing candidate (**Figure 3A**). For “diabetes” alpha 1 antagonists and kinase inhibitors were identified as possible novel treatments for diabetes (**Figure 3B**). We believe further investigation into these drug classes and diseases could lead to successful clinical applications.

CATNIP interpretability reveals reasoning for repurposing candidates

From our list of repurposing candidates, we chose two novel drug class-disease associations to further investigate.

Adrenergic uptake inhibitors applied to Parkinson's disease

First, we evaluated the relationship between “neurological diseases” and “adrenergic uptake inhibitors”. We focused on the drug pairs with the highest CATNIP scores, i.e. those predicted with the highest confidence to share at least one indication (**Figure 3C**). Of all the adrenergic uptake inhibitors, we found that amitriptyline and trimipramine, two anti-depressants, had the highest CATNIP scores with the “neurological diseases” drugs. The drugs that shared the strongest connections with amitriptyline and trimipramine were drugs approved for Parkinson's disease (PD). Specifically, metixene, atropine, pergolide and benztropine were associated with amitriptyline, according to CATNIP, and trimipramine was associated to benztropine and rotigotine. Trimipramine was also strongly connected with orphenadrine, which is sometimes used off label in PD, but will not be included in the following analyses.

Using the CATNIP model, we evaluated which features contributed towards the prediction of amitriptyline and trimipramine to share an indication with PD drugs. We found that target, gene ontology, and pathway similarity all strongly contributed to the predictions for both amitriptyline and trimipramine (**Figure 3D**, **Figure S10**). Since target similarity and distance between targets (in a protein-protein interaction network) were among the top contributing features, we investigated which gene targets were shared amongst these drug pairs. We found that amitriptyline targets three specific gene classes that are also targeted by at least one of the PD drugs: muscarinic acetylcholine receptors, G-coupled protein receptors (GPCRs), and alpha adrenergic receptor. Trimipramine also targets muscarinic acetylcholine receptors, alpha-adrenergic receptors, and dopamine transporters, which is similar to benztropine, a PD drug. All these receptors have well-defined relationships with PD and other neurological diseases^{25, 28, 29}, which adds support for repurposing amitriptyline and/or trimipramine.

Amitriptyline may be an ideal candidate for use in PD patients. We evaluated the shared molecular function gene ontology terms shared between amitriptyline and all four PD drugs. GPCR activity was once again identified (**Supplementary Data**). We then interrogated the biological pathways these drug targets are involved in and found many broad GPCR pathways overlapping between amitriptyline and the PD drugs (**Figure S11**) including the Reactome pathway “GASTRIN_CREB_SIGNALLING PATHWAY VIA PKC AND MAPK”. Several recent studies support the link between gastrin-releasing peptide signaling to brain function³⁰. Through CATNIP, we have identified “adrenergic uptake inhibitors” like amitriptyline and trimipramine as a possible treatment for PD.

Kinase inhibitors applied to Diabetes

Our CATNIP analysis identified an opportunity to repurpose “kinase inhibitors” for the treatment of diabetes (**Figure 3B**). Of the drug pairs evaluated in this context, the link between vandetanib, a thyroid cancer drug, and gliclazide, a Type 2 diabetes drug (CATNIP Score = 6.39, **Figure 3E**) was the strongest. This association was driven by target similarity and similarity between KEGG pathways of the drug targets (**Figure 3F**). Vandetanib and gliclazide have an overlapping target, VEGFA. Several KEGG pathways are shared between vandetanib and gliclazide including the “Cytokine cytokine receptor interaction” pathway (**Supplementary Data**). This pathway contains VEGFA, the shared target, and the epidermal growth factor receptor (EGFR), another one of vandetanib’s targets. The similarity between these two drug’s targets and pathway effects leads us to believe there is strong potential for vandetanib to be repurposed.

Discussion

Although considerable improvements have been made in drug repurposing efforts over the past decade, the use of previous disease associations will eventually curtail these improvements due to the imposed restriction of previous knowledge. Our new approach, CATNIP, could provide a highly effective aid to

drug repurposing endeavors. Here, we accurately predicted drugs that shared an indication, while keeping high levels of both sensitivity and specificity. Leveraging our prediction metric enabled us to generate a network for repurposing, identifying, and repurposing predictions based on system-wide drug scopes.

The CATNIP method allows for broad-scale drug repurposing opportunities to be readily identified. By identifying and interpreting our drug similarity features, we can investigate the possible mechanisms behind these repurposing candidates. The benefit of using drug similarity features is two-fold. First, these features are readily available for both approved and investigational drugs, which have been underserved by previous repurposing methods. Second, the interpretability of the features allows us to identify possible mechanisms of action when we back engineer what contributed to high CATNIP scores.

We found strong support for repurposing amitriptyline and trimipramine, both of which are in clinical use as anti-depressants, for PD. These drugs have many functions in addition to being adrenergic uptake inhibitors, such as serotonin blockers, anticholinergics, and the mechanisms overlapping with current PD drugs described above. Movement Disorders Society guidelines found insufficient evidence to support the use of amitriptyline for depression in PD³¹ and a published Practice Parameter found only level C evidence for its use³². However, amitriptyline has been commonly used for not only depression but other off-label indications in neurological disorders, including pain³³. While clinical trials have been conducted for the effect of amitriptyline on depression in PD patients³⁴, currently there are no trials evaluating amitriptyline or trimipramine as a treatment for other symptoms and signs of PD. There have, however, been preclinical studies evaluating amitriptyline as a potential therapy for PD. In rodent models of PD, amitriptyline affects levels of neurotrophic factors including BDNF³⁵ and decreases dopamine cell loss in these models^{36, 37}. It has been suggested to mitigate microglial inflammation³⁸. Moreover, with the suggestion

that amitriptyline may have shorter term symptomatic motor benefit, it may enhance levodopa efficacy³⁹.

When we more closely evaluated trimipramine, we found compelling evidence this could be a potential PD therapeutic. Specifically, the targets of trimipramine make it a potentially strong therapeutic to combat loss of motor function amongst PD patients. This benefit is due to the dual targeting of DRD2 and alpha 2 adrenergic receptors, which is similar to piribedil, an investigational PD medication that was not included within our final CATNIP network due to a lack of available information. In a review of piribedil, it was highlighted that the agonistic D2/D3 activity combined with alpha 2 adrenergic antagonism can lead to preservation of motor function⁴⁰. However, further research must be done to better understand the exact effects that trimipramine has on both dopamine and alpha 2 adrenergic receptors. Further research into trimipramine could quickly lead to a clinical trial for PD patients with specific motor function end points.

We also identified a repurposing opportunity with kinase inhibitors for the treatment of diabetes, due to the strong predicted connection between vandetanib, a thyroid cancer drug, and gliclazide. While there have been some preclinical animal studies investigating the use of kinase inhibitors in diabetes⁴¹,⁴², to our knowledge, there has yet to be an approved kinase inhibitor for the treatment of diabetes. Both vandetanib and gliclazide are known to target VEGFA, which has shown a clear connection to diabetes pathology⁴³ and treatment⁴⁴. Additionally, Hagberg et al. published work suggesting that antagonism of VEGFB, a gene within the same pathway as VEGFA, improves insulin sensitivity and increases skeletal muscle glucose uptake in *db/db* mice⁴⁵. Because vandetanib targets VEGFR1⁴⁶, the receptor VEGFB binds, it could also have insulin sensitizing effects. Further experimental work is required to verify this hypothesis⁴⁷.

Besides the targeting of VEGFA/VEGFR1, vandetanib's target EGFR can also potentially help diabetes pathology. Inflammatory cytokines (including, but not limited to, IL-8 and TNF- α) have been shown to be associated with the progression of diabetic neuropathy⁴⁸. The inhibition of EGFR through the use of a kinase inhibitor in past work has reduced the expression of both to IL-8 and TNF- α in rats⁴⁹. Therefore, we believe vandetanib could be considered as a potential diabetes treatment, due to its ability to target EGFR leading to a possible decrease in inflammatory cytokine production.

In addition to the exciting predicted repurposing opportunities we have chosen to highlight, many other drug classes showed significant repurposing potential for mental disorders, neurological diseases, and several different cancer types. While diving into each of these opportunities is outside the scope of this paper, we hope that researchers take it upon themselves to further investigate these candidate drug class-disease associations.

It is important to acknowledge certain limitations to CATNIP, such as data availability and the application to rare diseases. Although this model does not rely on disease similarity information, it does require known molecular target information to obtain peak predictive power. This target information can frequently be unavailable for early stage compounds. Additionally, this method would have limited use when searching for drugs to be repurposed for diseases with very few or no clinically approved compounds.

To our knowledge, CATNIP is the first method capable of predicting a novel indication for a drug without relying on disease similarities. Many predictions gained from CATNIP have substantial preclinical research or mechanistic support, suggesting that other predictions may also provide valuable information for future investigations. Due to its demonstrated ability to identify large scale drug repurposing opportunities, CATNIP will likely serve as a significant basis towards a bright future in drug repurposing efforts.

Methods

Indication Mapping

Using a custom Python script, we webscraped DrugBank 5.0⁵⁰ for drug compound names and indication information with a total of 3066 drugs being found. Indication information were run through the Unified Medical Language System (UMLS) tool, MetaMap¹³, to match DrugBank assigned indications to MESH IDs and UMLS Concept Unique Identifiers (CUIs). MetaMap is a computational approach that combines linguistic and natural language processing techniques to map biomedical texts to the UMLS Metathesaurus. MetaMap has previously been shown to successfully exceeded human mapping capabilities¹⁴. Using a custom Python script we identified synonym candidate to further improve indication semantics. A random subset of the indications were manually reviewed and found to correctly map to standardized terms with a 95% accuracy (**Figure S1B**). We then filtered our list of drugs to the 2576 drugs that shared at least one indication with another drug.

Similarity Feature Collection

Compound Features

Similarities between drugs were found by creating all possible pairs of the drugs with known indications. Multiple compound similarity features and drug target similarity features were collected. The drug targets listed within DrugBank 5.0⁵⁰ were used as our set of 'known targets' for each drug. Additionally, we collected genomic information about each drug target using MSigDB^{51, 52}. The features, sources and metrics used to measure similarity are listed in **Supplementary Table 1**. The measures of similarity included, but were not limited to, Pearson Correlation, Jaccard Index, and Dice Similarity. In cases where there was insufficient or missing information, features were imputed by using the median value for that feature in drug pairs with complete information.

Network Features

We curated a biological network that contains 22,399 protein-coding genes, 6,679 drugs, and 170 TFs. The protein-protein interactions represent established interaction⁵³⁻⁵⁵, which include both physical (protein-protein) and non-physical (phosphorylation, metabolic, signaling, and regulatory) interactions. The drug-protein interactions were curated from several drug target databases⁵⁵.

Statistical Analysis

For each similarity feature, a Kolmogorov-Smirnov (KS) test was performed between the set of drug pairs that shared an indication and those that did not share an indication. The KS test was chosen to identify non-linear predictive power. In addition, the Pearson correlation between all numeric features was calculated. These tests were performed using custom scripts in R statistical software⁵⁶.

Model Building

We trained a two-class classifier predictive model using the features described above. Our classes were determined as a binary of “shared” or “non-shared” indication. Drugs were only included if they shared an indication with at least one other drug. A 5-fold cross-validation gradient boosting model was used after careful model selection and implemented using the XGBoost package⁵⁷ within the R statistical software. Additional models that were tested and compared using the AUC and AUPRC of 5-fold cross-validation were: Support Vector Machine with a radial kernel model, logistic regression with elastic net and logistic regression with lasso, all using custom R scripts. A custom-made R script was used to implement a grid-search to optimize the hyperparameters of our model. Our model objective was a logistic regression for binary classification and we output a score pre-logistic transformation. The class size of “shared” vs. “non-shared” was imbalanced, therefore we applied downsampling to each fold of training via the R package Caret⁵⁸.

Classification Evaluation

For evaluating the model performance on predicting if two drugs share an indication, receiver operating characteristic (ROC) and precision-recall curve (PRC) curves were created in R using the pROC⁵⁹ and precrec⁶⁰ packages respectively. The raw-logistic values were normalized on a scale from 0-1 to enable easier interpretation and ROC/PRC calculation. Area-under-the-ROC curve (AUC) and area-under-the-PRC (AUPRC) scores were used to evaluate model performance.

Drug Similarity Network

Network Construction

We constructed a drug similarity network based upon our classifier results with drugs as nodes and our raw model output as the edge weight. This network was visualized using the visNetwork package⁶¹ and used in analyses using the iGraph package⁶² within R⁵⁶.

ATC Repurposing Analysis

The Anatomical Therapeutic Chemical (ATC) code for all drugs were found in DrugBank⁵⁰, and the highest level code was assigned. Drugs with multiple ATC codes assigned to them were re-assigned into the category “Various”. A circular repurposing network was created with ATC codes as the nodes using the iGraph⁶² and gGraph⁶³ packages with R⁵⁶. The graph edge weights were based on the mean classifier output between all drugs of each ATC code category. To reduce noise within the repurposing network an initial cut-off of drug pairs with a classifier output of 7.4 and above was implemented, leaving 792 drug pairs to examine. Manual literature searches were used to validate repurposing opportunities.

Drug Class Repurposing Analysis

Drug classes for all drugs were found in DrugBank⁵⁰ and were filtered to include only “inhibitor”, “antagonist,” and “agonist” classes that had at least 20 drugs, to

ensure enough statistical power. Additionally, we identified four main disease areas of interest: “mental disorders”, “neurological diseases”, “diabetes”, and “cancer”. The UMLS¹³ sematic codes “modb” and “neop” were used to identify indications falling within mental disorders and cancer, respectively. Cancer was further refined into different cancer types based on a keyword search in a custom Python script. All UMLS concept IDs containing the word “diabetes” were included within the diabetes category. For “neurological diseases”, we refined our list to only include Parkinson’s Disease, Alzheimer’s, Epilepsy, and Dementia, to balance both specificity in disease type and enough drugs to make statistically sufficient sample size.

Wilcox-Mann-Whitney tests between all drug class-disease associations were performed. The test specifically tested if the mean of the CATNIP scores of drug pairs with one drug being a member of the class of interest and the other being approved for the disease of interest were significantly different than the mean of the CATNIP scores of all drug pairs that included one drug within the class of interest and the other drug not being approved for the diseases of interest. A positive location shift meant that drug class-disease pairs had significantly higher CATNIP scores than drug class-non-disease pairs. A FDR multiple hypothesis correction was applied.

CATNIP Feature Effect Analysis

The effect of each feature on the CATNIP score for specific drug pairs was found by iteratively changing the feature value to the median value of that feature for all drug pairs. Since the clear majority of all drug pairs do not share an indication this is the best approximate for that feature having no contribution to the CATNIP score. The difference in the new CATNIP score and the correct CATNIP score was then measured.

Acknowledgements

C.M.G., J.E., N.S.M., and O.E. conceived, designed and developed the methodology for this work. C.M.G., J.E., and O.E. analyzed and interpreted the data. C.M.G. executed the machine learning analyses. C.M.G. and J.E. wrote the initial draft of the manuscript. M.D.G. provided expertise within diabetes, as well as other clinical aspects of the paper. C.H. provided expertise within Parkinson's Disease, as well as other clinical aspects of the paper. O.E. supervised the study. All the authors reviewed and approved the manuscript.

References

1. Adams, C.P. & Brantner, V.V. Estimating The Cost Of New Drug Development: Is It Really \$802 Million? *Health Affairs* **25**, 420-428 (2006).
2. Ishida, J., Konishi, M., Ebner, N. & Springer, J. Repurposing of approved cardiovascular drugs. *Journal of Translational Medicine* **14**, 269 (2016).
3. Goldstein, S.R. et al. Raloxifene use in clinical practice: efficacy and safety. *Menopause* **16**, 413-421 (2009).
4. Chiang, A.P. & Butte, A.J. Systematic evaluation of drug-disease relationships to identify leads for novel drug uses. *Clinical pharmacology and therapeutics* **86**, 507-510 (2009).
5. Gayvert, K.M., Madhukar, N.S. & Elemento, O. A Data-Driven Approach to Predicting Successes and Failures of Clinical Trials. *Cell chemical biology* **23**, 1294-1301 (2016).
6. Madhukar, N.S. et al. A New Big-Data Paradigm for Target Identification and Drug Discovery. *bioRxiv*, 134973 (2017).
7. Madhukar, N.S., Gayvert, K., Gilvary, C. & Elemento, O. A Machine Learning Approach Predicts Tissue-Specific Drug Adverse Events. *bioRxiv*, 288332 (2018).
8. McCullough, M. (Philly.com, Web; 2018).
9. Dudley, J.T., Deshpande, T. & Butte, A.J. Exploiting drug-disease relationships for computational drug repositioning. *Briefings in bioinformatics* **12**, 303-311 (2011).
10. Gottlieb, A., Stein, G., Rupp, E. & Sharan, R. PREDICT: A method for inferring novel drug indications with application to personalized medicine, Vol. 7. (2011).
11. Luo, H. et al. Drug repositioning based on comprehensive similarity measures and Bi-Random walk algorithm. *Bioinformatics* **32**, 2664-2671 (2016).
12. Wishart, D.S. et al. DrugBank: a knowledgebase for drugs, drug actions and drug targets. *Nucleic acids research* **36**, D901-D906 (2008).
13. Aronson, A.R. Effective mapping of biomedical text to the UMLS Metathesaurus: the MetaMap program. *Proceedings. AMIA Symposium*, 17-21 (2001).
14. Pratt, W. & Yetisgen-Yildiz, M. A study of biomedical concept identification: MetaMap vs. people. *AMIA ... Annual Symposium proceedings. AMIA Symposium* **2003**, 529-533 (2003).
15. Keiser, M.J. et al. Predicting new molecular targets for known drugs. *Nature* **462**, 175-181 (2009).

16. Craig Jordan, V. The role of tamoxifen in the treatment and prevention of breast cancer. *Current Problems in Cancer* **16**, 134-176 (1992).
17. Milani, M., Jha, G. & Potter, D.A. Anastrozole Use in Early Stage Breast Cancer of Post-Menopausal Women. *Clinical medicine. Therapeutics* **1**, 141-156 (2009).
18. Chen, L., Zeng, W.-M., Cai, Y.-D., Feng, K.-Y. & Chou, K.-C. Predicting Anatomical Therapeutic Chemical (ATC) classification of drugs by integrating chemical-chemical interactions and similarities. *PloS one* **7**, e35254-e35254 (2012).
19. Triolo, M., Annema, W., de Boer, J.F., Tietge, U.J. & Dullaart, R.P. Simvastatin and bezafibrate increase cholesterol efflux in men with type 2 diabetes. *European journal of clinical investigation* **44**, 240-248 (2014).
20. Teramoto, T., Shirai, K., Daida, H. & Yamada, N. Effects of bezafibrate on lipid and glucose metabolism in dyslipidemic patients with diabetes: the J-BENEFIT study. *Cardiovascular diabetology* **11**, 29 (2012).
21. Tenenbaum, A. et al. Effect of bezafibrate on incidence of type 2 diabetes mellitus in obese patients. *European heart journal* **26**, 2032-2038 (2005).
22. Pujols, L., Mullol, J. & Picado, C. Alpha and beta glucocorticoid receptors: relevance in airway diseases. *Current allergy and asthma reports* **7**, 93-99 (2007).
23. Cavallari, J.M., Jawaro, T.S., Awad, N.I. & Bridgeman, P.J. Glucagon for refractory asthma exacerbation. *The American Journal of Emergency Medicine* **35**, 144-145 (2017).
24. Insuela, D.B.R. et al. Glucagon induces airway smooth muscle relaxation by nitric oxide and prostaglandin E2. *Journal of Endocrinology* **225**, 205-217 (2015).
25. Langmead, C.J., Watson, J. & Reavill, C. Muscarinic acetylcholine receptors as CNS drug targets. *Pharmacology & therapeutics* **117**, 232-243 (2008).
26. Laruelle, M., Frankle, W.G., Narendran, R., Kegeles, L.S. & Abi-Dargham, A. Mechanism of action of antipsychotic drugs: from dopamine D2 receptor antagonism to glutamate NMDA facilitation. *Clinical therapeutics* **27**, S16-S24 (2005).
27. Zhang, J., Yang, P.L. & Gray, N.S. Targeting cancer with small molecule kinase inhibitors. *Nature reviews cancer* **9**, 28 (2009).
28. Perry, E., Smith, C. & Perry, R. Cholinergic nicotinic and muscarinic receptors in dementia of Alzheimer, Parkinson and Lewy body types. *Journal of Neural Transmission-Parkinson's Disease and Dementia Section* **2**, 149-158 (1990).
29. Xu, Y. et al. Neurotransmitter receptors and cognitive dysfunction in Alzheimer's disease and Parkinson's disease. *Progress in neurobiology* **97**, 1-13 (2012).
30. Roesler, R. & Schwartzmann, G. Gastrin-releasing peptide receptors in the central nervous system: role in brain function and as a drug target. *Frontiers in endocrinology* **3**, 159 (2012).
31. Seppi, K. et al. The Movement Disorder Society evidence-based medicine review update: treatments for the non-motor symptoms of Parkinson's disease. *Movement Disorders* **26**, S42-S80 (2011).

32. Miyasaki, J. et al. Practice Parameter: Evaluation and treatment of depression, psychosis, and dementia in Parkinson disease (an evidence-based review):[RETIRED]: Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* **66**, 996-1002 (2006).
33. Frost, J., Okun, S., Vaughan, T., Heywood, J. & Wicks, P. Patient-reported outcomes as a source of evidence in off-label prescribing: analysis of data from PatientsLikeMe. *Journal of medical Internet research* **13**, e6 (2011).
34. Antonini, A. et al. Randomized study of sertraline and low-dose amitriptyline in patients with Parkinson's disease and depression: effect on quality of life. *Movement disorders: official journal of the Movement Disorder Society* **21**, 1119-1122 (2006).
35. Paumier, K.L. et al. Tricyclic antidepressant treatment evokes regional changes in neurotrophic factors over time within the intact and degenerating nigrostriatal system. *Experimental neurology* **266**, 11-21 (2015).
36. Paumier, K.L. et al. Chronic amitriptyline treatment attenuates nigrostriatal degeneration and significantly alters trophic support in a rat model of parkinsonism. *Neuropsychopharmacology* **40**, 874 (2015).
37. Kandil, E.A., Abdelkader, N.F., El-Sayeh, B.M. & Saleh, S. Imipramine and amitriptyline ameliorate the rotenone model of Parkinson's disease in rats. *Neuroscience* **332**, 26-37 (2016).
38. Lauterbach, E.C. Repurposing psychiatric medicines to target activated microglia in anxious mild cognitive impairment and early Parkinson's disease. *American journal of neurodegenerative disease* **5**, 29 (2016).
39. Kamińska, K., Lenda, T., Konieczny, J., Wardas, J. & Lorenc-Koci, E. Interactions of the tricyclic antidepressant drug amitriptyline with L-DOPA in the striatum and substantia nigra of unilaterally 6-OHDA-lesioned rats. Relevance to motor dysfunction in Parkinson's disease. *Neurochemistry international* **121**, 125-139 (2018).
40. Millan, M.J. From the cell to the clinic: a comparative review of the partial D2/D3 receptor agonist and α 2-adrenoceptor antagonist, pramipexole, in the treatment of Parkinson's disease. *Pharmacology & therapeutics* **128**, 229-273 (2010).
41. Louvet, C. et al. Tyrosine kinase inhibitors reverse type 1 diabetes in nonobese diabetic mice. *Proceedings of the National Academy of Sciences* **105**, 18895-18900 (2008).
42. Kikuchi, Y. et al. A Rho-kinase inhibitor, fasudil, prevents development of diabetes and nephropathy in insulin-resistant diabetic rats. *Journal of Endocrinology* **192**, 595-603 (2007).
43. Aiello, L.P. et al. Vascular endothelial growth factor in ocular fluid of patients with diabetic retinopathy and other retinal disorders. *New England Journal of Medicine* **331**, 1480-1487 (1994).
44. Duh, E. & Aiello, L.P. Vascular endothelial growth factor and diabetes: the agonist versus antagonist paradox. *Diabetes* **48**, 1899-1906 (1999).
45. Hagberg, C.E. et al. Targeting VEGF-B as a novel treatment for insulin resistance and type 2 diabetes. *Nature* **490**, 426 (2012).

- 708 46. Bianco, R. et al. Vascular endothelial growth factor receptor-1 contributes to
709 resistance to anti-epidermal growth factor receptor drugs in human cancer
710 cells. *Clinical Cancer Research* **14**, 5069-5080 (2008).
- 711 47. Robciuc, M.R. et al. VEGFB/VEGFR1-induced expansion of adipose
712 vasculature counteracts obesity and related metabolic complications. *Cell*
713 *metabolism* **23**, 712-724 (2016).
- 714 48. Navarro-Gonzalez, J.F. & Mora-Fernandez, C. The role of inflammatory
715 cytokines in diabetic nephropathy. *Journal of the American Society of*
716 *Nephrology* **19**, 433-442 (2008).
- 717 49. Qu, W.-s. et al. Inhibition of EGFR/MAPK signaling reduces microglial
718 inflammatory response and the associated secondary damage in rats after
719 spinal cord injury. *Journal of neuroinflammation* **9**, 178 (2012).
- 720 50. Wishart, D.S. et al. DrugBank 5.0: a major update to the DrugBank database
721 for 2018. *Nucleic Acids Research* **46**, D1074 (2017).
- 722 51. Subramanian, A. et al. Gene set enrichment analysis: A knowledge-based
723 approach for interpreting genome-wide expression profiles. *Proceedings of*
724 *the National Academy of Sciences* **102**, 15545 (2005).
- 725 52. Liberzon, A. et al. Molecular signatures database (MSigDB) 3.0. *Bioinformatics*
726 **27**, 1739-1740 (2011).
- 727 53. Das, J. & Yu, H. HINT: High-quality protein interactomes and their
728 applications in understanding human disease. *BMC systems biology* **6**, 92
729 (2012).
- 730 54. Khurana, E., Fu, Y., Chen, J. & Gerstein, M. Interpretation of genomic variants
731 using a unified biological network approach. *PLoS computational biology* **9**,
732 e1002886 (2013).
- 733 55. Aksoy, B.A. et al. PiHelper: an open source framework for drug-target and
734 antibody-target data. *Bioinformatics* **29**, 2071-2072 (2013).
- 735 56. R Core Team in R Foundation for Statistical Computing. Retrieved from
736 <http://www.r-project.org/> (Vienna, Austria; 2017).
- 737 57. Chen, T. & Guestrin, C. in Proceedings of the 22nd ACM SIGKDD International
738 Conference on Knowledge Discovery and Data Mining 785-794 (ACM, San
739 Francisco, California, USA; 2016).
- 740 58. Kuhn, M. Building Predictive Models in R Using the caret Package. *2008* **28**,
741 26 (2008).
- 742 59. Robin, X. et al. pROC: an open-source package for R and S+ to analyze and
743 compare ROC curves. *BMC Bioinformatics* **12**, 77-77 (2011).
- 744 60. Saito, T. & Rehmsmeier, M. Precrec: fast and accurate precision-recall and
745 ROC curve calculations in R. *Bioinformatics* **33** (1), 145-147 (2017).
- 746 61. Almende, B.V., Thieurmél, B. & Robert, T. visNetwork: Network Visualization
747 using 'vis.js' Library. *The R Journal* **10**, 251-268 (2018).
- 748 62. Csardi, G. & Nepusz, T. The igraph software package for complex network
749 research. *InterJournal Complex Systems*, 1695-1695 (2006).
- 750 63. Pedersen, T.L. ggraph: An Implementation of Grammar of Graphics for
751 Graphs and Networks. **33** (1), 145-147 (2018).
- 752 64. Carter, N.J. Bilastine. *Drugs* **72**, 1257-1269 (2012).

- 753 65. Krause, K., Spohr, A., Zuberbier, T., Church, M.K. & Maurer, M. Up-dosing with
754 bilastine results in improved effectiveness in cold contact urticaria. *Allergy*
755 **68**, 921-928 (2013).
- 756 66. Greaves, M.W. Antihistamines in Dermatology. *Skin Pharmacology and*
757 *Physiology* **18**, 220-229 (2005).
- 758 67. Kuna, P. et al. The role and choice criteria of antihistamines in allergy
759 management - expert opinion. *Postępy dermatologii i alergologii* **33**, 397-410
760 (2016).
- 761 68. La Rosa, M.e.a. A randomized, double-blind, placebo-controlled, crossover
762 trial of systemic flunisolide in the treatment of children with severe atopic
763 dermatitis. *Current Therapeutic Research* **56**, 720 - 726 (1995).
- 764 69. Ekström, T., Lindgren, B.R. & Tibbling, L. Effects of ranitidine treatment on
765 patients with asthma and a history of gastro-oesophageal reflux: a double
766 blind crossover study. *Thorax* **44**, 19-23 (1989).
- 767 70. Dixon, A.E. et al. A pilot randomized controlled trial of pioglitazone for the
768 treatment of poorly controlled asthma in obesity. *Respiratory Research* **16**,
769 143 (2015).
- 770 71. Moore, M. et al. Amoxicillin for acute lower respiratory tract infection in
771 primary care: subgroup analysis of potential high-risk groups. *The British*
772 *journal of general practice : the journal of the Royal College of General*
773 *Practitioners* **64**, e75-e80 (2014).
- 774 72. Reznikov, L.R. et al. The vagal ganglia transcriptome identifies candidate
775 therapeutics for airway hyperreactivity. *American Journal of Physiology-Lung*
776 *Cellular and Molecular Physiology* **315**, L133-L148 (2018).
- 777 73. Beigelman, A., Chipps, B.E. & Bacharier, L.B. Update on the utility of
778 corticosteroids in acute pediatric respiratory disorders. *Allergy and asthma*
779 *proceedings* **36**, 332-338 (2015).
- 780 74. Hua, F., Wang, X. & Zhu, L. Terlipressin Decreases Vascular Endothelial
781 Growth Factor Expression and Improves Oxygenation in Patients with Acute
782 Respiratory Distress Syndrome and Shock. *The Journal of Emergency*
783 *Medicine* **44**, 434-439 (2013).
- 784 75. Crestani, B. et al. Octreotide treatment of idiopathic pulmonary fibrosis: a
785 proof-of-concept study. *European Respiratory Journal* **39**, 772 (2012).
- 786 76. Abid, S. et al. 17β-estradiol dysregulates innate immune responses to
787 *Pseudomonas aeruginosa* respiratory infection and is modulated by estrogen
788 receptor antagonism. *Infection and immunity* **85**, e00422-00417 (2017).
- 789 77. Kharkevich, D.A., Chizh, B.A. & Kasparov, S.A. Stimulant effect of thyrotropin-
790 releasing hormone and its analog, RGH 2202, on the diaphragm respiratory
791 activity, and their antagonism with morphine: possible involvement of the N-
792 methyl-D-aspartate receptors. *Brain research* **551**, 110-115 (1991).
- 793 78. El-Haggar, S.M., Farrag, W.F. & Kotkata, F.A. Effect of ketotifen in obese
794 patients with type 2 diabetes mellitus. *Journal of Diabetes and its*
795 *Complications* **29**, 427-432 (2015).
- 796 79. Manjunath, S., Kugali, S.N. & Deodurg, P.M. Effect of clonidine on blood
797 glucose levels in euglycemic and alloxan-induced diabetic rats and its

798 interaction with glibenclamide. *Indian journal of pharmacology* **41**, 218-220
799 (2009).
800 80. Paul, S., Wand, M., Emerick, G.T. & Richter, J.M. The role of latanoprost in an
801 inflammatory bowel disease flare. *Gastroenterology report* **2**, 232-234
802 (2014).
803 81. Kern, T.S. et al. Topical Administration of Nepafenac Inhibits Diabetes-
804 Induced Retinal Microvascular Disease and Underlying Abnormalities of
805 Retinal Metabolism and Physiology. *Diabetes* **56**, 373 (2007).
806 82. Pereira Arias, A.M. et al. Indomethacin decreases insulin secretion in patients
807 with type 2 diabetes mellitus. *Metabolism* **49**, 839-844 (2000).
808
809
810
811
812
813
814




Figure 1: Schematic of CATNIP repurposing approach. A) The use of drug similarity properties to predict if two drugs will share an indication using a gradient boosting model, the model is referred to as CATNIP. B) Schematic showing the use of CATNIP output scores to create a network, with the scores used as edge weights. The colors of each drug represent the known disease and this demonstrates how one could identify novel indications for drugs through the network.




Figure 2: CATNIP model accurately predicts drugs that share an indication and can be used for repurposing. A) Receiver-operating characteristic curve for CATNIP, the performance for drug pairs with high and low structural similarity is also shown. B) A network of all drug pairs with a CATNIP score higher than 7.4. Nodes (drugs) are colored based on ATC classification and a specific example of repurposing between ATC classifications is highlighted. C) A graph of all ATC classification and the median CATNIP score between the drugs belonging to each of them (only including drug pairs with > 7.4 CATNIP score). The edges between ATC Classifications with the highest median CATNIP scores are colored red.




Figure 3: CATNIP identifies drug class repurposing opportunities. The location shift, calculated using Wilcoxon-Mann-Whitney for all CATNIP scores of drug class-disease drug pairs vs drug class-non-disease drug pairs for A) “neurological diseases” and B) “diabetes”. Drug classes are colored based on the percent of drugs within the class that are approved for treatment of the specific disease and only significant associations are shown (FDR < 0.1). C) The network of neurological drugs and adrenergic uptake inhibitors drug pairs with the highest CATNIP scores. D) The decrease in the CATNIP score when removing each feature for amitriptyline and select Parkinson’s Disease drugs. E) The network of anti-diabetes and kinase inhibitor drug pairs with the highest CATNIP scores. F) The decrease in the CATNIP score when removing each feature for the drug pair vandetanib and gliclazide.

848 Table 1: Literature Support for ATC Repurposing Predictions

ATC Code 1	ATC Code 2	Reference
Dermatologicals	Respiratory System	64-68
Alimentary Tract and Metabolism	Respiratory System	69-72
Sensory Organs	Respiratory System	73-75
Systemic Hormonal Preparations, Excluding Sex Hormones And Insulins	Respiratory System	76, 77
Sensory Organs	Alimentary Tract and Metabolism	78-82

849

850 **Supplementary Figures**

851 Figure S1: MetaMap performs well in drug indication mapping. A) The number of
852 occurrences of different UMLS semantic types. B) The accuracy of mapping
853 indications using MetaMap for indications categorized a “Structured” and the
854 “Description” section.

855

856 Figure S2: Target ontology similarity data types vary for drug pairs that share an
857 indication and those that do not. The violin plots of similarity distributions for the
858 similarities of targets’ A) biological processes, B) cellular component, C)
859 molecular function, D) chemical perturbation, E) oncological, F) immunogenic, G)
860 micro-RNA, and H) transcription factor. Statistical significance found by
861 Kolmogorov-Smirnov test.

862

863 Figure S3: Target similarity data types vary for drug pairs that share an indication
864 and those that do not. The violin plots of similarity distributions for the similarities
865 of A) targets, B) the Protein-Protein Interaction network distance between targets
866 and the C) correlation of target essential within cancer cell lines. Statistical
867 significance found by Kolmogorov-Smirnov test.

868

869 Figure S4: Target pathway similarity data types vary for drug pairs that share an
870 indication and those that do not. The violin plots of similarity distributions for the
871 similarities of the A) reactome pathways, B) all pathway types and C) KEGG
872 pathways a drug’s target is known to be involved within. Statistical significance
873 found by Kolmogorov-Smirnov test.

874

875 Figure S5: Structure similarity varies for drug pairs that share an indication and
876 those that do not. A) The violin plot of the Dice chemical fingerprint similarity,
877 statistical significance found by Kolmogorov-Smirnov test.

Figure S6: CATNIP performs significantly better than random. A) The Precision – Recall curve for classifying if two drugs share an indication using CATNIP and the random expectation.

Figure S7: CATNIP identifies drug class repurposing opportunities. The location shift, calculated using Wilcoxon-Mann-Whitney for all CATNIP scores of drug class-disease drug pairs vs drug class-non-disease drug pairs for A) Mental Disorder, B) Skin Cancer, C) Lung Cancer, D) Breast Cancer, E) Thyroid Cancer, F) Large Intestine Cancer, G) Upper Aerodigestive Tract Cancer H) Gastric Cancer I) Renal Cancer. Drug classes are colored based on the percent of drugs within the class that are approved for treatment of the specific disease and only significant associations are shown (FDR < 0.1).

Figure S8: CATNIP identifies other drug class repurposing opportunities within cancer. The location shift, calculated using Wilcoxon-Mann-Whitney for all CATNIP scores of drug class-disease drug pairs vs drug class-non-disease drug pairs for A) Urinary Tract Cancer, B) Pleura Cancer, C) Endometrium Cancer, D) Ovarian Cancer, E) Pancreatic Cancer, F) Bone Cancer, G) Oesophagus Cancer, H) Lymphoma/Leukemia, and I) Autonomic Ganglia Cancer. Drug classes are colored based on the percent of drugs within the class that are approved for treatment of the specific disease and only significant associations are shown (FDR < 0.1).

Figure S9: CATNIP scores are statistically higher between drugs of certain drug classes and drugs that treat associated diseases. The distributions of CATNIP score between A) kinase inhibitors and drugs known to treat cancer and those that do not and B) dopamine antagonists and drugs known to treat mental illness and those that do not.

Figure S10: Target features drive the prediction of trimipramine as a Parkinson's Disease treatment. A) The decrease in the CATNIP score when removing each feature for trimipramine and select Parkinson's Disease drugs.

Figure S11: Many pathways or gene ontology groups overlap, fueling CATNIP predictions. The overlap between amitriptyline and select Parkinson's Disease drugs for A) reactome pathways, B) KEGG pathways, and D) molecular function gene ontologies. The overlap between vandetanib and gliclazide for A) reactome pathways, B) KEGG pathways, and D) molecular function gene ontologies.

Supplementary Tables

Table S1: The drug similarity features used within CATNIP.

Table S2: Comparison of model performance using other model types.

Supplementary Data

File 1: All pathways and gene ontologies that amitriptyline's targets and the targets of select Parkinson's Disease drugs' targets are associated with.

File 2: All pathways and gene ontologies that trimipramine's targets and the targets of select Parkinson's Disease drugs' targets are associated with.

File 3: All pathways and gene ontologies that vandetanib's targets and gliclazide's are associated with.