

## Influence of plant genotype and soil on the wheat rhizosphere microbiome: identification of a core microbiome across eight African and European soils

Marie Simonin<sup>1,2</sup>, Cindy Dasilva<sup>1</sup>, Valeria Terzi<sup>3</sup>, Eddy L. M. Ngonkeu<sup>4</sup>, Diégane Diouf<sup>5,6</sup>, Aboubacry Kane<sup>5,6</sup>, Gilles Béna<sup>1</sup>, Lionel Moulin<sup>1,5\*</sup>

1. IRD, Cirad, Univ Montpellier, IPME, Montpellier, France.

2. INRA, UMR1345 Institut de Recherches en Horticulture et Semences, SFR4207 QUASAV, F-49071, Beaucozé, France.

3. CREA-GB, Research Centre for Genomics and Bioinformatics, I-29017 Fiorenzuola d'Arda, Italy

4. Department of Plant Biology, Faculty of Science, University of Yaoundé I, Yaoundé, Cameroon.

5. Laboratoire Mixte International Adaptation des Plantes et Microorganismes Associés aux Stress Environnementaux (LAPSE), IRD, ISRA, UCAD, Dakar, Sénégal.

6. Département de Biologie Végétale, Université Cheikh Anta Diop (UCAD), Dakar, Sénégal.

\*Corresponding author:

Lionel Moulin, [lionel.moulin@ird.fr](mailto:lionel.moulin@ird.fr)

## Abstract

An avenue to improve cereal yields is to harness and manipulate the plant microbiome to increase crop nutrition and resistance to pathogens and abiotic stressors. To develop microbiome engineering solutions, more research is needed to characterize plant microbiomes across contrasted geographical locations, agricultural practices and plant genotypes. Previous work on the wheat microbiome focused on bacterial diversity and more marginally on archaeal and fungal diversity. Hence, no integrative assessment of the wheat rhizospheric microbiome including protist diversity (i.e. amoeba, ciliates, stramenopiles) is currently available. Here, we characterized the rhizosphere microbiome of wheat by considering both prokaryotic (archaea and bacteria) and eukaryotic (fungi and protists) communities in soils from four different countries (Cameroon, France, Italy, Senegal). The goals of this study were to determine the influence of wheat genotype, agricultural practices (conventional vs organic) and soil type on the rhizosphere microbiome. An additional goal was to determine if a rhizosphere core microbiome existed across these different countries.

We found a limited effect of the wheat genotype on the rhizosphere microbiome (explain 2% of variance) and observed that the majority of the microbial taxa were consistently associated to multiple wheat genotypes grown in the same soil. Large differences in taxa richness and in microbiome structure were observed between the eight soils studied (57% of variance) and the two agricultural practices (10% of variance). However, despite these differences in microbiome composition across soils, we observed that 179 taxa (2 archaea, 104 bacteria, 41 fungi, 32 protists) were consistently associated to wheat roots, constituting a core microbiome. In addition to being prevalent, these few core taxa were highly abundant and collectively represented 50% of the relative abundance of our entire wheat microbiome dataset.

We provide evidence of a wheat core microbiome and propose a “most wanted” taxa list, including 9 hubs (potential keystone taxa), that should be targets for future culturomics, metagenomics and for the creation of wheat synthetic microbiomes. Additionally, we confirm that protists are an integral part of the plant holobiont and we encourage more research to study their roles for plant nutrition and in controlling plant diseases.

## Keywords

Phytobiome; Winter Wheat; Rhizosphere; Protists; Fungi; Archaea; Microbial network; Keystone taxa

## Introduction

Wheat, with a worldwide production of more than 730 million metric tons in 2018-19 (<http://www.fao.org/statistics/en/>) is now the second most important grain crop (behind corn) that is mainly used for food, animal fodder, and industrial raw materials. To face the challenges of climate change and of the nutritional needs of a growing world population, there is a necessity of a 70% increase in cereal production by 2050 [1]. This increase in agricultural production needs to be sustainable and have limited environmental consequences on biodiversity, human health and climate [2]. An avenue to improve cereal yields is to harness and manipulate the plant microbiome to improve its nutrition and resistance to pathogens and abiotic stressors [3]. The plant microbiome is comprised of complex communities of bacteria, archaea, fungi and protists that interact with the host plant in different compartments (e.g. rhizosphere, endosphere, phyllosphere) [4]. In particular, the root-rhizosphere interface is the nexus of a variety of interactions from which the plant can benefit to acquire mineral nutrients or water and thus is a compartment that is key for future microbiome engineering solutions in agriculture [5].

Following the advent of next generation sequencing, several studies characterized the rhizospheric wheat microbiome and investigated the influence of the compartment (rhizoplane vs endosphere), crop management or wheat genotypes on the diversity and structure of these complex microbial communities [6–9]. However, a great majority of these studies focused only on bacterial diversity (16S rRNA gene) and more rarely on fungal diversity (ITS) [10–12]. To our knowledge, no integrative assessment of the wheat rhizospheric microbiome including also the diversity of protists (i.e. amoeba, ciliates, stramenopiles) is currently available. Protists as predators, saprotrophs or phototrophs influence nutrient cycles in the rhizosphere and exert a strong top-down control on microbial (mainly bacterial) biomass and composition [13]. Still, despite protists high diversity and biomass in the rhizosphere, they represent an overlooked component of the plant holobiont.

Moreover, a limited number of studies investigated the effect of the soil used for growing wheat on the rhizosphere microbiome [14,15]. For other crops, the culture soil has been demonstrated to be the most important factor structuring the root microbiome before crop management and plant genotype [16,17]. More work is thus required to study the influence of soil type and geographical location on the wheat microbiome and hence determine which microbial taxa are specific to a soil/location or are common across multiple wheat production systems (i.e. wheat core microbiome). Determining if a wheat core microbiome exists is crucial to help orientate future strain cultivation-based efforts and design microbiome engineering efforts through modifications of agricultural practices or microbial inoculations.

These analyses to determine the presence of a wheat core microbiome need to be conducted at the finest taxonomic level possible (microbial species or strain), as the ecologies and metabolisms of

closely related microbial taxa can be extremely different. Previous analyses of wheat microbiome diversity were conducted by lumping amplicon sequences at a 97% identity threshold to create microbial operational taxonomic units (OTUs) that were assumed to represent “species”. New analyses including thousands of genomes indicate that optimal thresholds to represent bacterial species and discriminate phenotypes is at 100% of identity (i.e. exact sequence variant) for the V4 region of the 16S rRNA gene [18]. New denoising algorithms for amplicon sequences now enable to eliminate sequencing errors and resolve exact sequence variants (ESVs) that vary only by one nucleotide [19]. These new bioinformatic methods open new possibilities to characterize plant microbiomes and their thousands of microbial taxa at the finest taxonomic level achievable by short-read amplicon sequencing.


Here, we characterized the rhizosphere microbiome of wheat by considering both prokaryotic (archaea and bacteria) and eukaryotic (fungi and protists) communities of different wheat genotypes grown in soils from four different countries. The goals of this study were to determine the influence of wheat genotype, agricultural practices (conventional vs organic) and soil type on the diversity, structure and taxonomic composition of the rhizosphere microbiome. An additional goal was to determine if a rhizospheric core microbiome existed by identifying microbial taxa present on wheat roots grown in very contrasting soils from different countries. In a growth chamber experiment, we first characterized the rhizosphere microbiome (here soil tightly bound to roots) of eight different genotypes of winter bread wheat (*Triticum aestivum* L.) grown in one soil (FR2) to assess specifically the genotype effect. Second, we characterized the rhizosphere microbiome of three wheat genotypes grown in eight contrasted soils (Table 1) collected from different countries: in Central Africa, Cameroon (CAM1 and CAM2 soils) and West Africa, Senegal (SEN1 and SEN2 soils) and in Europe, France (FR1 and FR2 soils) and Italy (IT1 and IT2 soils). The total microbiome diversity was characterized using amplicon sequencing of the marker genes 16S rRNA (prokaryotic diversity: archaea and bacteria) and 18S rRNA (eukaryotic diversity: fungi and protists). At the taxon level (exact sequence variant), we identified the core microbiome across all soils and genotypes and determined the relative abundance of these core taxa and their potential role as hub taxa in the wheat microbiome using network analyses.

## Results

### **Limited effects of wheat genotype on the rhizosphere microbiome in one soil**

We observed limited effects of the wheat genotype on the rhizosphere microbiome after 30 days of growth in experimental chambers under controlled conditions. There were no statistical differences on the total ESV richness (272 to 368 total ESVs on average) and phylogenetic diversity (15.2 to 18.7 total PD) between 7 genotypes for the total microbial diversity (Fig. 1A, 1B). Only the genotype BM presented a lower diversity than some genotypes. Interestingly the genotypes recruited different proportions of prokaryotes and eukaryotes on their root surfaces. Some genotypes with the highest prokaryotic diversity (AP, RP, TM) presented a lower eukaryotic diversity, while the genotypes that presented the highest eukaryotic diversity (CT, CH, CY) had low to intermediate levels of prokaryotic diversity (Fig. 1A). As a consequence, no significant correlation was observed between prokaryotic and eukaryotic ESV richness because of the contrasted patterns observed between the different genotypes (Fig. 1C).

No significant effect of wheat genotype was either observed on total microbiome structure (Fig. 1D,  $P=0.13$ ). This limited genotype effect was also observed when analyzing the patterns separately on prokaryotic (Fig. S1,  $P=0.04$ ) and eukaryotic community structure (Fig. S1,  $P=0.22$ ). The phylogenetic trees show that hundreds of microbial taxa are associated to the roots of multiple wheat genotypes. We observed that 117 microbial taxa were reliably associated (i.e. present in multiple replicates) to 3 or 4 wheat genotypes, 93 taxa were associated to 5-6 genotypes and 85 taxa to 7-8 genotypes (Fig. 1 bar graph). These taxa associated to multiple genotypes were especially affiliated to the bacterial families Burkholderiaceae, Chitinophagaceae, Caulobacteraceae or fungal Ascomycota and cercozoan Filosa for the eukaryotes (list of 85 taxa associated to 7-8 genotypes is available in Additional file 1). Additionally, we observed that the bacterial phyla or class Actinobacteria, Deltaproteobacteria and Alphaproteobacteria were more specifically associated to the genotypes with a high bacterial richness (AP, RP, TM, Fig. 1).




**Fig. 1: In the soil FR2, influence of 8 wheat genotypes on the rhizosphere microbiome. A) Observed Exact Sequence Variants (ESVs) richness and B) Faith's phylogenetic diversity of the rhizosphere microbiome for the eight genotypes for the prokaryotic, eukaryotic and total (prokaryotic + eukaryotic) microbial community. Note that on the x axis, the genotypes are ordered from low to high total microbiome diversity. The statistical differences between the eight genotypes are represented by different letters: two genotypes sharing the same letter are not statistically different. For clarity, only the statistical results for the total**

diversity are presented, detailed results for prokaryotic and eukaryotic diversity are presented in Fig S1. C) Absence of linear correlation between prokaryotic and eukaryotic diversity (ESV richness) across the 8 wheat genotypes. D) NMDS ordination showing the absence of significant differences in the structure of the rhizosphere microbiome (prokaryotes and eukaryotes combined) between the eight wheat genotypes. Detailed results for prokaryotic and eukaryotic community structure in Fig S1. Phylogenetic trees of the E) prokaryotic and F) eukaryotic communities showing that many microbial taxa are present on the root of multiple wheat genotypes. On the eight outer rings (E) or vertical bands (F), the presence of a colored circle indicates that an ESV was found in the rhizosphere microbiome of a specific wheat genotype. For legibility purposes on the trees and the bar graph, only ESVs that were observed in minimum two replicates of the same genotype condition are represented (n=448 ESVs for prokaryotes and n=179 ESVs for eukaryotes).

## Characterization of wheat rhizospheric microbiome across eight African and European soils

### *Strong effect of soil on the structure and diversity of the rhizosphere microbiome*

Cultivation of wheat in different soils led to large differences in the ESV richness (100 to 380 total ESVs) and phylogenetic diversity (8 to 22 total PD) of the rhizosphere microbiome (Fig. 2A, 2B). SEN1 soil had the lowest diversity and SEN2, FR1 and IT2 presented the highest microbiome diversity. The two Cameroonian soils (CAM1 and CAM2) and the FR2 soil presented intermediate levels of diversity (300-350 ESVs). The proportion of prokaryotes and eukaryotes varied between soils but across all soils a positive relationship between prokaryote and eukaryote diversity was found (Fig. 2C,  $R^2=0.4$ ,  $P<0.001$ ). A very strong effect of the soil was also observed on the structure of the microbial community explaining 57% of the variance, while the effects of wheat genotype (3 genotypes tested, 2% of the variance) and the interactive effects between genotype and soil were not significant (Fig. 2D). The root microbiome samples separated along the NMDS axis 1, mainly according to the continent where the soil was collected (African soils on the right and European soils on the left). A clear separation between the Cameroonian and Senegalese soils was also observed on the NMDS axis 2, while the differences in community structure between the two European soils were not as strong.




**Fig. 2: Comparisons of the rhizosphere microbiome across eight agricultural soils from Cameroon (CAM1, CAM2), France (FR1, FR2), Italy (IT1, IT2) and Senegal (SEN1, SEN2). A) Observed Exact Sequence Variants (ESVs) richness and B) Faith's phylogenetic diversity of the rhizosphere microbiome in the eight soils for the prokaryotic, eukaryotic and total (prokaryotic + eukaryotic) community. Note that on the x axis, the soils are ordered from low to high total microbiome diversity. The statistical differences between the eight soils are represented by different letters: two soils sharing the same letter are not statistically different. For clarity, only the statistical results for the total diversity are presented, detailed results for prokaryotic and eukaryotic diversity are presented in Fig S2. C) Significant positive linear correlation between prokaryotic and eukaryotic diversity (ESV richness) across the eight soils. D) NMDS ordination showing a significant soil effect on the structure of the rhizosphere microbiome (prokaryotes and eukaryotes combined). Detailed results for prokaryotic and eukaryotic community structure in Fig S2.**

### *Agricultural practices affect the structure and diversity of the rhizosphere microbiome*

To assess the influence of agricultural practices, we took advantage of the fact that the Italian and French soils were collected in long-term experimental plots under two types of farming practices at the same location (conventional farming with chemical fertilization, and organic farming with culture rotation with legumes or gramineous plants, Table 1). We observed a significant difference in total taxa richness between the two agricultural practices, only for the Italian soils (Fig. 3A). Taxa richness was 56% higher

in the organic Italian soil than in the conventional Italian soil. However, we observed a significant difference in microbiome structure between conventional and organic farming in both Italian and French soils (Fig. 3B). The separations between Conventional and Organic microbiome samples were present on the NMDS axis 2 and the agricultural practices explained globally 10% of the variance in community structure (Fig. 3B). Using differential abundance testing, we identified 51 taxa that were significantly enriched in conventional farming (in Italian or French soils) and 16 taxa in Organic farming (Fig. S3). For the taxa enriched in Conventional farming, half were bacteria and the other half eukaryotes (Cercozoans, Fungi and Oomycetes) including taxa affiliated to *Chitinophaga*, *Flavobacterium*, *Bacillus*, *Pseudomonas*, *Glissomonadida*, *Cordyceps* or *Phytophtora*. For the taxa enriched in organic farming, most bacteria were affiliated to *Flavobacterium* and *Cellvibrio*, while the eukaryotes were two uncultured fungi, two uncultured Stramenopiles and the oomycetes *Pythium*.




**Fig. 3: Effect of agricultural practices (conventional vs organic farming) on the root microbiome diversity and structure in the French and Italian soils. A) Observed Exact Sequence Variants (ESVs) richness of the rhizosphere microbiome for the prokaryotic, eukaryotic and total (bacterial + eukaryotic) community. B) NMDS ordination showing a significant effect of agricultural practices on the structure of the rhizosphere microbiome (prokaryotes and eukaryotes). The list of the 67 taxa significantly affected by farming practices is available in Additional file 1.**

#### *Phylogenetic distribution of rhizosphere microbial taxa across the eight soils*

Across the eight soils studied, we observed that the wheat rhizosphere microbiome was extremely diverse with a total of 4760 ESVs identified. For the Prokaryotes, Archaea represented only 0.9% of the reads (29 ESVs, Phylum Thaumarcheota only) that were affiliated to two archaeal families of ammonia oxidizers (Nitrosophaeraceae and Nitrosotaleaceae) and the remaining 99.1% of the reads belonged to 3652 bacterial ESVs in the final dataset. Across all soils, the most abundant and diverse bacterial families were Burkholderiaceae (10% relative abundance, 251 ESVs, phylum Proteobacteria) followed by Chitinophagaceae (4.2% relative abundance, 296 ESVs, phylum Bacteroidetes) and

Flavobacteriaceae (4% relative abundance, 79 ESVs, phylum Bacteroidetes) (bubble plot, Fig. 4). For the Eukaryotes (1125 ESVs), fungal taxa were the most abundant and diverse (61% relative abundance, 459 ESVs), followed by Cercozoa (17%, 415 ESVs) and Oomycota taxa (14%, 47 ESVs). The eukaryotic dominant clades were Sordariomycetes (7.3%, 90 ESVs, Ascomycota), Peronosporales (8.6%, 36 ESVs, Oomycota) and Chytridiomycetes (6.5%, 110 ESVs, Chytridiomycota) (bubble plot, Fig. 4). As seen on the phylogenetic trees, these abundant and diverse prokaryotic or eukaryotic taxa were present in multiple soils (Fig. 4). More specifically, when considering all the prokaryotic and eukaryotic taxa associated to wheat roots (n=4760 EVs), we found that 402 taxa (8.5%) were observed in both African (Cameroon and Senegal) and European (France and Italy) soils and that more than 2000 taxa were specific of the 2 continents (Venn diagram, Fig. 4). The phyla in which few taxa were shared between continents (clear continental phylogenetic signals) were principally in the Acidobacteria, Chloroflexi, and Bacteroidetes phyla for Bacteria and the phylogenetic signals were not as clear for eukaryotes (Fig. 4). However, it is important to note that the large majority of prokaryotic and eukaryotic ESVs were observed in multiple countries (Fig. 4).



**Fig. 4: Phylogenetic trees of the prokaryotic (top) and eukaryotic (bottom) taxa present in the eight African and European soils. On the eight outer rings represent the eight soils and the presence of a colored circle indicates that an ESV was found in this soil. Core taxa are indicated by a red triangle on the outer ring. For legibility purposes on the trees, only ESVs that were observed in minimum two replicates of the same genotype-soil condition are represented (n=1478 ESVs for prokaryotes and n=537 ESVs for eukaryotes). In the top-left corner, a Venn diagram represent the number of taxa specific of African or European soils and**


**the number taxa observed in both continents. A bubble plot presenting the relative abundance of the most abundant clades (relative abundance >0.5%) across all samples and the size of the points represent the number of taxa (ESVs) in each clade.**

## **Evidences for a root core microbiome shared across African and European soils**


### ***2.9% of the microbiome diversity represent 50% of the rhizosphere microbiome abundance***

Core taxa (ESV level) of the wheat rhizosphere microbiome were identified based on a criterion of presence of the ESV in 25% of the samples with no relative abundance threshold (core taxa can be rare). This prevalence criterion enabled the identification of the wheat core microbiome specific of Africa, of Europe and common to both continents. The 179 ESVs that responded to the criterion defining a core taxon in our study are indicated by a red triangle on the phylogenetic trees in Fig. 4. These core taxa were distributed across all domains (2 Archaea, 104 Bacteria, 73 Eukaryotes) and spanned 65 families (Fig. 5A, B) and 84 genera. We observed that 118 taxa out of the 179 taxa were observed in soils from both continents (66%), while only 20 core taxa were specific of African soils and 41 of European soils (Fig. 5A). Surprisingly, this list of only 179 ESVs on the total 4760 ESVs (prokaryote + eukaryote) represented 51% of all the sequences in the dataset (Fig. 5D, E) indicating that collectively these 179 core taxa represent half of the relative abundance of the rhizosphere microbiome. From one soil to another, the number of observed taxa belonging to this list of core taxa varied (Fig. 5C) with the SEN1 soil presenting only 22 core taxa on average (total relative abundance of 52%, Fig. 5D) and the FR2 soil presenting 90 core taxa on average (total relative abundance of 65%, Fig. 5D). Interestingly, we did not observe a positive correlation between the number of core taxa observed and their relative abundance in a sample, indicating that across very different soils, these taxa always represented a high relative abundance even when a low number of them were present (Fig. S4).

We also studied the relative abundance of each core taxon in the different soils and found that the core taxa clustered in two main groups based on their distribution patterns across all samples (Fig. 6). These results show that even if most core taxa are present in both African and European soils, they are generally very abundant only in soils from one continent and not the other. The 98 taxa affiliated to the cluster 1 were predominantly very abundant in European soils and the 81 ESVs of the cluster 2 were most abundant in African soils. Interestingly, the proportion of prokaryotes and eukaryotes were different between the 2 clusters. The cluster 1 (dominant in European soils) was principally composed of bacterial taxa (65%) and fungal taxa (19%) or protists were less represented, while the cluster 2 (dominant in African soils) was dominated by eukaryotes (28% Fungi, 16% Rhizaria, 5% Stramenopiles) and presented 48% of bacterial taxa (Fig. 6).



**Fig. 5:** A) Venn diagram showing that among the 179 taxa identified as core taxa, 118 taxa were present in both African and European soils and 20 taxa were specific of African soils and 41 of European soils. B) A bubble plot presenting the number of core taxa and their cumulative relative abundance by clade. C) Number and D) cumulative relative abundance of the core taxa present in each sample in the eight soils. Note that on the x axis the soils are ordered from low to high y-axis value. The green line represents the average across all samples and the associated confidence interval (95%). E) Pie charts showing that while the core taxa represent only 2.9% of the total diversity, their cumulative relative abundance is 50.3% across all samples (half of the reads).




**Fig. 6: Heatmap representing the relative abundance of the 179 core taxa in the different soils. Based on the distribution of the taxa in the samples, the taxa clustered in two main groups principally associated to the continent (Africa vs Europe) where the soil was collected (but see soil SENb2). The cluster 1 regroups 98 ESVs (65% Bacteria, 19% Fungi) and the cluster 2 regroups 81 ESVs (48% Bacteria, 28% Fungi, 16% Rhizaria).**

## Co-occurrence network of the core wheat microbiome and identification of hub taxa

The cross-domain network of the 179 core taxa computed with the SParse InversE Covariance estimation for Ecological Association Inference (SPIEC-EASI) was comprised of two main components, one being dominated by bacterial taxa and the other dominated by fungal and cercozoan (protists) taxa (Fig. 7A). The two main components broadly corresponded to the two clusters identified earlier based on the taxa relative abundance across all samples, with the cluster 1 gathering taxa mainly abundant in European soils and the cluster 2 in African soils (Fig. 7B). In the network, the predicted interactions were predominantly positive (440 positive edges vs 56 negative edges) and for each ESV, the average number of associations with other taxa was 5.54 (node degree). Eight taxa were identified

as potential “hub” taxa based on their centrality and number of associations in the network. These hub taxa are hypothesized to be keystone taxa or key connector taxa in a community because of their central position in the network and their high connectivity. Three fungal ESVs that were extremely prevalent across all soils (60 to 85% of samples) were identified as hub taxa: *Mortierella* (Mucoromycota), *Exophiala* (Ascomycota) and an uncultured Chytridiomycetes (Chytridiomycota). The other five hub taxa were bacterial ESVs in the Gammaproteobacteria phylum, one ESVs affiliated to the genus *Thermomonas* and two ESVs from the Burkholderiaceae family: *Massilia* and an uncultured taxon; in the Deltaproteobacteria class, one ESVs affiliated to the genus *Anaeromyxobacter*; in the class Bacteroidia, one ESV affiliated to the species *Niastella koreensis* and finally an ESV from the class Blastocatellia from an uncultured Blastocatellaceae. Interestingly, five of these potential hub taxa were connectors between the two main components of the network described above, with these connections being often negative edges.



**Fig. 7: Cross domain network of the rhizosphere core taxa including both prokaryotes and eukaryotes (179 ESVs, n=60 samples).** The lines (edges) between the taxa (nodes) represent the predicted interactions either positive (light blue) or negative (red). The same network is presented twice but the nodes representing the different taxa are colored by their domain or eukaryotic supergroup in the left figure (A) and are colored by their cluster affiliation in the right figure (B) as defined in the heatmap in Fig. 5. The core taxa affiliated to the cluster 1 generally had higher relative abundance in European soils, while the cluster 2 taxa were most abundant in African soils. The “Hub” taxa represented with a triangle symbol in the networks were identified as the ones presenting the highest centrality (betweenness and closeness centrality) and connectivity (node degree) in the network (panels C and D). These hubs are potential keystone taxa or key connectors in the community. The taxonomic affiliations of the hub taxa are indicated in the bottom plots and they are highlighted with triangle symbols.

A list of the most prevalent core taxa across all samples (>40% of samples) and of the ones identified as hub taxa are presented in Table 1 (complete list in Additional file 1 with sequence and taxonomy information). The most prevalent taxa across all samples were three eukaryotic ESVs affiliated to the fungal genera *Mortierella* (85% of samples) and *Fusarium* (82%) and a cercozoan zooflagellate belonging to the Allapsidae family (77%). The most prevalent bacterial taxa were *Bradyrhizobium japonicum* (72% of samples) and an *Arthrobacter* (70%). We observed that the relative abundance across all samples of the core taxa varied greatly (min=0.02%, max=5.7%) and that a majority of the core taxa were relatively rare in terms of abundance in the community (median=0.1%). This was also true for the eight taxa identified as hubs for which only two fungal taxa had a high relative abundance (4.24% and 1.71%) and the six others had relative abundance inferior to 0.3%. Using a complementary dataset not presented here, we determined that at least 33% of the core taxa were also observed in the wheat endosphere in the same plant samples (Table 1).

**Table 1: Summary information on the most prevalent core taxa (>40% of samples) and hub taxa**

Group	Phylum/Class	Family	Final affiliation	Prevalence	Relative Abundance	Hub Taxa	Present in Endosphere	Cluster
Fungi	Mucoromycota	Mucoromycotina	<i>Mortierella</i>	85%	4.24%	Hub	Endo	1
Fungi	Ascomycota	Eurotiomycetes	<i>Exophiala</i>	62%	0.16%	Hub	Endo	1
Fungi	Chytridiomycota	Chytridiomycetes	Uncultured Chytridiomycetes	60%	1.71%	Hub	Endo	2
Bacteria	Gammaproteobacteria	Burkholderiaceae	<i>Massilia</i>	47%	0.25%	Hub	Endo	2
Bacteria	Blastocatellia	Blastocatellaceae	Uncultured Blastocatellaceae	43%	0.30%	Hub	Endo	1
Bacteria	Gammaproteobacteria	Burkholderiaceae	Uncultured Burkholderiaceae	42%	0.20%	Hub	Endo	1
Bacteria	Delta proteobacteria	Archangiaeae	<i>Anaeromyxobacter</i>	38%	0.05%	Hub	Endo	2
Bacteria	Gammaproteobacteria	Xanthomonadaceae	<i>Thermomonas</i>	38%	0.25%	Hub	Endo	1
Bacteria	Bacteroidia	Chitophagaceae	<i>Niastella koreensis</i>	32%	0.09%	Hub	Endo	2
Fungi	Ascomycota	Sordariomycetes	<i>Fusarium</i>	82%	1.69%	Endo	1	2
Rhizaria	Filosa-Sarcomonadea	Allapsidae	Allapsidae (Group-Te)	77%	0.91%			2
Bacteria	Alphaproteobacteria	Xanthobacteraceae	<i>Bradyrhizobium japonicum</i>	72%	0.14%	Endo	2	2
Bacteria	Actinobacteria	Micrococcaceae	<i>Arthrobacter</i>	70%	0.19%			2
Rhizaria	Filosa-Granofilosea	Limnofilidae	<i>Limnofila borokensis</i>	70%	0.20%			2
Bacteria	Nitrospira	Nitrospiraceae	<i>Nitrospira</i>	67%	0.14%			1
Bacteria	Gammaproteobacteria	Enterobacteriaceae	<i>Pantoea agglomerans</i>	67%	0.39%	Endo	2	2
Rhizaria	Filosa-Sarcomonadea	Paracercomonadidae	<i>Paracercomonas</i>	63%	0.15%	Endo	1	1
Bacteria	Gammaproteobacteria	Burkholderiaceae	<i>Paucibacter</i>	62%	0.09%			2
Fungi	Ascomycota	Sordariomycetes	<i>Chaetomium</i>	60%	0.59%	Endo	1	1
Bacteria	Gammaproteobacteria	Burkholderiaceae	<i>Ramlibacter</i>	60%	0.09%			2
Bacteria	Actinobacteria	Geodermatophilaceae	<i>Blastococcus</i>	57%	0.13%			1
Bacteria	Gammaproteobacteria	Burkholderiaceae	<i>Noviherbipirillum</i>	57%	0.21%			2
Bacteria	Alphaproteobacteria	Caulobacteraceae	<i>Phenylobacterium</i>	57%	0.10%			2
Fungi	Ascomycota	Sordariomycetes	<i>Bionectria</i>	55%	0.31%			2
Fungi	Ascomycota	Dothideomycetes	Uncultured Dothideomycetes	55%	0.31%	Endo	1	1
Rhizaria	Filosa-Sarcomonadea	Cercomonadidae	<i>Eocercomonas</i>	53%	0.05%			2
Bacteria	Bacteroidia	Chitophagaceae	<i>Flavatilea</i>	50%	0.08%			2
Fungi	Ascomycota	Sordariomycetes	Uncultured Sordariomycetes	50%	0.25%	Endo	2	2
Bacteria	Gammaproteobacteria	Burkholderiaceae	<i>Massilia</i>	48%	0.31%	Endo	1	1
Bacteria	Bacteroidia	Chitophagaceae	Uncultured Chitophagaceae	48%	0.42%	Endo	1	1
Fungi	Ascomycota	Dothideomycetes	Uncultured Pleosporales	48%	0.18%	Endo	2	2
Bacteria	Gammaproteobacteria	Burkholderiaceae	<i>Ramlibacter</i>	47%	0.08%			2
Rhizaria	Filosa-Imbricatea	Thaumatomonadidae	<i>Thaumatomonas</i>	47%	0.22%			2
Fungi	Ascomycota	Dothideomycetes	Uncultured Dothideomycetes	47%	0.22%			2
Fungi	Chytridiomycota	Chytridiomycetes	Spizellomycetales-Rhizophlyctidales	45%	0.45%			2
Fungi	Chytridiomycota	Chytridiomycetes	Spizellomycetales-Rhizophlyctidales	45%	0.19%			2
Fungi	Ascomycota	Sordariomycetes	Uncultured Sordariomycetes	45%	0.37%			1
Bacteria	Alphaproteobacteria	Caulobacteraceae	<i>Asticcacaulis</i>	43%	0.12%			1
Fungi	Basidiomycota	Tremellomycetes	<i>Cryptococcus</i>	43%	1.40%	Endo	1	1
Stramenopile	Oomycota	Peronosporales	<i>Pythium</i>	43%	0.46%	Endo	2	2
Fungi	Chytridiomycota	Chytridiomycetes	<i>Rhizophlyctis</i>	43%	0.25%			1
Bacteria	Gammaproteobacteria	Burkholderiaceae	Uncultured Burkholderiaceae	43%	0.39%			1
Bacteria	Bacteroidia	Microscillaceae	Uncultured Microscillaceae	43%	0.19%			1
Bacteria	Gammaproteobacteria	Burkholderiaceae	<i>Acidovorax</i>	42%	0.79%	Endo	1	1
Rhizaria	Filosa-Sarcomonadea	Allapsidae	Allapsidae (Group-Te)	42%	0.66%			2
Bacteria	Bacilli	Bacillaceae	<i>Bacillus</i>	42%	0.23%			2
Bacteria	Bacteroidia	Chitophagaceae	<i>Flavisolibacter</i>	42%	0.07%			2
Bacteria	Bacteroidia	Chitophagaceae	<i>Flavisolibacter</i>	42%	0.11%			2
Bacteria	Gammaproteobacteria	Methylphilaceae	<i>Methylotenera</i>	42%	0.04%			2
Bacteria	Alphaproteobacteria	Sphingomonadaceae	<i>Sphingomonas</i>	42%	0.07%			2
Fungi	Chytridiomycota	Chytridiomycetes	Uncultured Chytridiomycetes	42%	0.76%			2
Stramenopile	MAST	MAST-12C	Uncultured MAST-12C	42%	0.08%			2
Rhizaria	Filosa-Sarcomonadea	Sandonidae	Uncultured Sandonidae	42%	0.81%	Endo	1	1
Fungi	Ascomycota	Sordariomycetes	Uncultured Sordariomycetes	42%	0.13%	Endo	2	2

## Discussion

In this study, we took an integrative approach to characterize the rhizosphere microbiome (i.e. soil tightly bound to the roots) of multiple wheat genotypes by considering the total microbial diversity, including archaea, bacteria, fungi and protists in soils from different countries. We show that the main drivers of the wheat root microbiome are the culture soil and agricultural practices, while wheat genotypes had very limited effects. Across eight contrasted soils from two continents and three wheat genotypes, we found that 179 prokaryotic and eukaryotic taxa were consistently associated to wheat roots, constituting a core microbiome. In addition to being prevalent, these few core taxa were highly abundant and collectively represented 50% of the relative abundance of the wheat microbiome on average. This work enabled the identification of a “most wanted” taxa list for future wheat microbiome research that will be targets for culturomic, genomic and synthetic community studies to develop microbiome engineering in agriculture.

### **The wheat root microbiome is shaped by the soil and agricultural practices but not by the plant genotype**

By growing eight genotypes in the same soil, we observed that wheat varieties had very limited effects on rhizosphere microbiome diversity (richness and phylogenetic diversity) and structure. The variability among replicates of a same genotype was often larger than between genotypes. These results were confirmed in our second sub-experiment where we did not find a significant effect of the three genotypes selected in the eight test soils (only 2% of variance explained). These results are consistent with previous work showing a small effect of the wheat genotype explaining only 1 to 4% of the total variance in the community or effects appearing only after two years of cultivation on the field [6,14,20]. Hence, the lack of genotype effect in our study could be explained by the short duration of our experiment with the harvest taking place at the vegetative stage when root microbiome is not fully “mature” [6,21].

Consistent with the literature, the main factor shaping the wheat root microbiome diversity and structure was the test soil (57% of the variance). A clear separation between African and European soils was observed indicating potential continental effects on plant microbiome structure. It should be noted that the two European soils were closely clustered on the NMDS, while the Cameroonian and Senegalese soils were clearly separated on the second axis. We also observed distinct community structure between the two soils sampled from each country, that were the strongest for the two Senegalese soils. These separations could be explained by differences in soil parameters, especially pH (lower pH in African soils in our study) that has been found to be the main soil parameter influencing wheat microbiomes at a regional scale in China [15].

Large differences in taxa richness and phylogenetic diversity were present between the eight soils (100 to 380 total taxa), indicating that not only the microbiome composition was different but also the number of taxa selected by the plant, as already shown in the wheat rhizosphere [7,14,20]. We did not observe clear diversity patterns associated to countries, for instance the two Senegalese soils had respectively the lowest and highest phylogenetic diversity of the study and large differences in ESV richness were also observed between the two Italian soils. In contrast, the diversity indices of the two Cameroonian or French soils were not statistically different from each other despite large differences in soil properties and agricultural practices. Interestingly across all soils, prokaryotic diversity was positively correlated to eukaryotic diversity signifying that the environmental conditions favourable to high diversity are similar across all microorganisms. These results show in the contrasted eight soils studied that prokaryotic diversity is 2 to 3 times higher than eukaryotic diversity on wheat roots.

Agricultural practices were also a significant driver of microbiome structure (10% of the variance) in Italian and French soils in which we compared root microbiomes between conventional and organic farming. However, the taxa richness was significantly affected by agricultural practices only in the Italian soils. These results are very similar to the findings of Hartman et al. (2018)[9] that observed more effects of conventional vs organic practices on wheat microbiome structure (9-10% of the variance) than on diversity but also with other studies in various cropping systems [11,21]. The IT2 soil sampled from an organic farming field presented a higher taxa richness (56% higher) than IT1. It is noteworthy that this IT1 soil has been sampled from a field uninterruptedly sowed for 15 years with small grain cereals (wheat and barley) that can be considered a long-term monoculture that were cultivated adopting conventional farming procedures. The lower microbial diversity of IT1 soil in comparison with the other European soils can be explained by the different hypotheses. First, the higher diversity in the IT2 organic field could be associated to the addition of new microorganisms with amendments in organic farming. Moreover, crop rotation compared to long-term monocropping is leading on average to a ~15.1% increase in microbial richness [22].

The root-associated microbiome was dominated by the bacterial family Burkholderiaceae (10% of all sequences), oomycetes from the order Peronosporales (8.6%) and fungi from the Sordariomycetes (7.3%) and Chytridiomycetes (6.5%) classes. The diversity of protists, especially Cercozoans, was extremely high, representing half of the eukaryotic diversity in the microbiome, with many abundant predator taxa affiliated to *Rhogostoma* (testate amoebae), *Allapsidae-Group Te* (glissomonads) and *Sandonidae* (glissomonads). This study shows that non-fungal eukaryotes are diverse and abundant on wheat roots and they deserve more attention to determine their functional roles in nutrient cycling and in controlling microbiome structure through microbial predation.

While many taxa were found to be shared between different genotypes and soils (see core microbiome section below), we also observed microbial taxa that were specific of some soils or agricultural practices

and that were responsible for the significant shifts in community structure and diversity presented above. In particular, ESVs affiliated to the Acidobacteria, Chloroflexi and Bacteroidetes phyla were often found only in African or European soils but not in both, suggesting a continental phylogenetic signal or full clades sensitive to differences in soil pH. Differentially abundant taxa between conventional and organic farming were identified in almost all groups (bacteria, oomycetes, fungi, cercozoans) indicating a clear restructuring of the microbiome as a whole. Interestingly, potential plant pathogenic taxa were found to be enriched in conventional (*Phytophthora*) or organic farming (*Pythium*).

Using an integrative approach of the plant microbiome, contrary to most studies that separate bacteria and fungi in their analyses, we intentionally grouped all taxa in our structure and diversity analyses together to present a global view of the wheat microbiome as it occurs in nature. With this approach, we show that the soil and agricultural practices are the key drivers of wheat root microbiome and that a large part its micro-eukaryotic diversity is constituted of protists that has been so far overlooked in the plant holobiont.

### **Evidences for a wheat core microbiome and identification of hub taxa**

Many prokaryotic and eukaryotic taxa were found to be consistently associated to wheat roots from multiple genotypes and soils, with the exact same 16S or 18S rRNA sequence variants detected in soils sampled thousands of kilometers apart. More specifically, we identified 179 taxa that constitute the core microbiome of wheat roots in eight African and European soils. These core taxa were highly prevalent (present in multiple soils) but also very abundant which was not necessarily expected as we did not impose a relative abundance threshold to identify core taxa. Collectively, these 179 taxa (2.9% of the total diversity) represented 50% of the reads in the dataset, suggesting an important biomass of these microorganisms on root surfaces. These findings show that under very contrasted conditions (continents, soil types, farming practices, wheat genotypes), wheat associates with the same microbial species or strains (ESV level) which indicate a potential co-evolution between wheat and these microorganisms. These results are consistent with large-scale microbiome studies that identified the core microbiome of different plants like *Arabidopsis thaliana* [16] or sugarcane [23] and a global soil microbiome study that highlighted that only 500 OTUs account for half of soil microbial communities worldwide [24].

The wheat root microbiome can be diverse and variable (100 to 380 taxa) but these findings show that it can be decomposed in two parts: a core microbiome presenting a low diversity that is stable and abundant across conditions; and an “accessory” microbiome that is extremely diverse and condition-specific [25]. This observation is consistent with a previous wheat microbiome study that showed that highly co-occurring taxa (i.e. potential core taxa) were not affected by cropping practices and that the “accessory” wheat microbiome could be manipulated by changes in farming practices [9]. The high prevalence and abundance of the core taxa identified suggest an ecological significance in the root habitat and warrants further research to isolate and phenotype these organisms that are mainly uncultured

(i.e. 62 core taxa without genus affiliations). Based only on amplicon sequencing data, we cannot determine the role of these microorganisms for plant fitness or if the absence of some of these core taxa could impact the wheat microbiome or plant health. Thus, here we provide a “most-wanted” list composed of 2 archaea, 103 bacteria, 41 fungi and 32 protists that should be the target of future culturomics, metagenomics and for the creation of wheat synthetic microbiomes.

Most of these core taxa were present in both African and European soils (118 taxa) but based on their relative abundance they grouped into two clusters, with taxa having a higher abundance in either European soils (Cluster 1) or African soils (Cluster 2). These results were confirmed in the cross-domain network of all core taxa that separated into two main components that broadly corresponded to the two clusters. Future synthetic community studies should consider and evaluate the effects of contrasted relative abundances of the core taxa to represent realistic structuration of the wheat microbiomes.

Among all core taxa, a fungal ESV affiliated to the genus *Mortierella* (phylum Mucoromycota) occupied a crucial place in the wheat microbiome. This taxon had the highest prevalence (85% of samples) and relative abundance in the dataset (4.2%) and was identified as a hub taxon in the cross-domain network. This *Mortierella* ESV was detected on the wheat rhizoplane but also in the endosphere in a complementary dataset not presented in this article. This fungus is described as a saprotroph-symbiotroph [26] capable of solubilizing phosphate in the rhizosphere [27]. Our results are consistent with other studies that identified *Mortierella* as a keystone taxa of the wheat rhizosphere unsensitive to cropping practices in Switzerland or the USA [9,28] with potential positive effects on wheat yield in Canada [29]. Altogether, these findings encourage future research to develop targeted cultivation approaches of the wheat-associated *Mortierella* fungi that seem to play a central role in the wheat microbiome across the world. The other most prevalent fungal core taxa were affiliated to the genera *Fusarium* (Ascomycota, 82% of samples), *Exophiala* (Ascomycota, 62% of samples, hub) and *Chaetomium* (Ascomycota, 60% of samples) described as pathotrophs or saprotrophs that also have been observed on wheat roots in different cropping systems and described as hub taxa [21,28,30].

The taxonomic group that exhibited the highest number of core taxa was bacteria (103 taxa) with the dominant family Burkholderiaceae presenting the highest number of core taxa (28 taxa) including two hub taxa. In particular, an ESV affiliated to the genus *Massilia* was identified as a hub taxon in the wheat microbiome of our study. This taxon is known as a copiotrophic root colonizer [31] found to be dominant in the wheat and maize rhizosphere [11,32]. The most prevalent bacterial core taxon was *Bradyrhizobium japonicum* (Alphaproteobacteria, 72% of samples), a species frequently found in root-associated microbiomes, and which can fix nitrogen in symbiosis with legumes (especially soybean).

Here, we show also that archaea and protists that are microbial groups generally ignored in most plant microbiome studies were characterized as wheat core taxa. The two archaea identified as core taxa were both affiliated to the genus *Nitrososphaera* (Phylum Thaumarcheota) that are involved in the nitrogen

cycle as ammonia-oxidizers in the nitrification process (oxidation of ammonia in nitrate) [33] and are very abundant in soils [34,35]. Interestingly, another nitrifier taxa affiliated to the bacterial genus *Nitrospira* (Nitrospira phylum, 67% of all samples) was also identified as core taxon which suggests that microorganisms involved in nitrification could play an important role for nitrogen availability on roots. *Nitrososphaera* taxa were described as hubs in the wheat rhizosphere [30] and as highly prevalent in the rhizosphere and rhizoplane of *Arabidopsis thaliana* and maize [36,37], indicating that root surfaces represent a common habitat for these archaea.

Among protists, the most prevalent core taxa were all cercozoans from the phyla Filosa-Sarcomonadae and Filosa-Granofilosea affiliated to *Allapsidae – Group Te* (77% of samples), *Limonofila borokensis* (70%) and *Paracercomonas* (63%). These three taxa have also been identified as extremely common and abundant in the rhizosphere of *Arabidopsis thaliana* [38] and *Allapsidae – Group Te* as a hub taxon in the maize rhizosphere [39]. Five Oomycetes were also categorized as core taxa with three of them affiliated to the *Pythium* genus and one to *Phytophthora* that are both described as potential plant pathogens. These observations are consistent with previous plant microbiome studies considering oomycetes diversity, that always described *Pythium* as the most dominant taxa among oomycetes [38,40,41]. These results confirm that protists are an integral part of the plant holobiont and their roles in controlling microbial populations through predation, disease incidence or contribution to nutrient cycles through the microbial loop deserve more attention [13].

## Conclusions

This work presents a detailed characterization of the wheat root-associated microbiome under contrasted environmental conditions and propose a “most-wanted” list of microorganisms identified as core and hub taxa. In this list of 179 prokaryotic and eukaryotic taxa, 62 are uncultured taxa (no genus affiliation) and only 11 taxa have a species affiliation, suggesting that almost nothing is known about the ecology of the key microorganisms associated to a major crop like wheat. Future research efforts are needed to try to cultivate these microorganisms associated to wheat as they could unlock new knowledge and biotechnological resources to improve crop yields and resistance to diseases in a sustainable way. These efforts should be conducted with an integrative approach of the plant holobiont by not only focusing on bacteria and fungi but also considering archaea, oomycetes and cercozoans.

## Methods

### Soil description, sampling and physico-chemical analyses

Eight soils primarily cultivated with wheat which have been previously treated with chemical or organic fertilization were collected from four countries: Senegal and Cameroon (West and Central Africa);

France and Italy (Europe). They were chosen for their contrasting physico-chemical and land-use characteristics to represent a diversity of agricultural practices (conventional, organic, agroforestry), texture, pH, carbon and nutrient contents (Table 1). The soils from France and Italy were collected from long-term experimental plots comparing conventional and organic cropping practices and were used to test the influence of agricultural practices on the wheat microbiome. In each field, we collected 12 microsites distant of 20 m from each other and from 0 to 15 cm depth (500 g each). After passing through a 4 mm sieve, samples from the same field were mixed together, a sub-sample was used for physico-chemical analyses, and the remaining was used for pot experiments. The soil samples were shipped to France to the lab IPME to perform the experiments described below. All soil analyses were performed at the Laboratoire des Moyens Analytiques (LAMA) using published protocols: pH and electrical conductivity (EC) in an aqueous extract [42], soil particle size distribution [43], total organic carbon [44], total and available concentrations of nitrogen and phosphorus.

**Table 1: Main soil characteristics and localization of the collection sites**

Soil	Texture	pH	N total (%)	Organic C total (%)	P total (mg/kg)	CaCO <sub>3</sub> total (%)	Agricultural Practices	Lat,Long	Country / region
CAM1	Silty-Clay-Loam	4.5	0.92	9.3	1227	0	Conventional	5.383333, 10.116667	Cameroon, Tubah
CAM2	Silt-Loam	5.7	0.90	10.2	1479	0	Conventional	5.966667, 10.300000	Cameroon, Tubah
FR1	Silt-Loam	8.1	0.24	2.3	1352	4.5	Conventional	45.777144, 3.142915	France, Clermont-Ferrand
FR2	Silt-Loam	8.2	0.21	4.3	1112	23.8	Organic (legume rotation)	45.768014, 3.157844	France, Clermont-Ferrand
IT1	Silt	8.3	0.16	2.6	777	12.8	Conventional	44.933333, 9.900000	Italy, Fiorenzuola D'arda PC
IT2	Silt-Loam	8.1	0.26	4.4	656	22.5	Organic ( <i>Lolium perenne</i> rotation)	44.933333, 9.900000	Italy, Fiorenzuola D'arda PC
SEN1	Clay	6.3	0.12	1.5	355	0	Conventional	16.533333, -15.183333	Senegal, Fanaye (Saint Louis)
SEN2	Loamy Sand	6.7	0.06	0.7	83	0	Agroforestry	12.816667, -14.883333	Senegal, Saré Yéro Bana (Kolda Region)

### Experimental design: 2 sub-experiments

During the same experiment, we performed two sub-experiments in a growth-chamber under controlled conditions in Montpellier (France) with the eight soils collected from African and European countries. The sub-experiment 1 looked at the effects of tender wheat genotype alone on the root microbiome by

growing eight genotypes in one soil (FR2). The eight winter bread wheats grown in this sub-experiment were: Apache (AP), Bermude (BM), Champlein (CH), Carstens (CT), Cheyenne (CY), Rubisko (RB), Soissons (SS) and Terminillo (TM).

The Sub-experiment 2 looked at the interactive effects of wheat genotype and soil on the wheat root microbiome. Three genotypes from the eight presented above were selected (AP, BM, RB) and were grown in the eight soils collected in Africa and Europe. These three genotypes were selected because they are the most recent genotypes (in terms of date of registration) and are highly cultivated.

### **Plant growth and harvest**

Seeds were surface disinfected by washing for 40 min in a 9,6% sodium hypochlorite solution [45], rinsed 5 times for 3 min in sterile water, then chlorine traces were removed by washing 3 times for 7 min in 2% (w/v) sodium thiosulfate [46] and then rinsed 5 times again for 3 min in sterile water, and left in sterile water for another 45 min. Sterilized seeds were then incubated for germination on sterile agar plates (8 g.L<sup>-1</sup>) for 2 days in the dark at 27°C. Five germinated seeds of each genotype were grown on each sampled soil mixed with 50% of sterile sand (size 0.4 to 1.4 mm) in pots of 400 mL with 3 replicates per condition. All pots were incubated in a growth chamber (Temperature day/night: 22°C/18°C, duration day/night: 16 h/8 h, humidity: 60-65%) for a month corresponding to the 3-4 leaf stage at sampling time. The plants were covered with a transparent plastic bag to avoid cross contaminations, especially during watering. Overall, for the sub-experiment 1, eight varieties of wheat were grown in one soil (8 conditions) in triplicates (24 pots). For the sub-experiment 2, three varieties of wheat were grown in eight soils (24 conditions) in triplicates (72 pots). At the end of the experiment, the three tallest plants from each pot were selected. Their roots were collected and shaked to remove non-adhering soil, and placed in sterile falcon conical tubes (50 ml) containing 20 mL of NaCl 0.9% to collect rhizosphere soil (tightly bound to roots). These tubes were vortexed for 5 min, roots were removed from the tubes and the tubes were centrifuged for 30 min at 4000 rpm (Eppendorf). After centrifugation, the pellet constituting the rhizosphere microbiome was flash frozen in liquid nitrogen and stored at -80°C before DNA extraction.

### **DNA extraction and high-throughput sequencing**

Total DNA was extracted using the PowerSoil Extraction Kit (Mo Bio Laboratories, Carlsbad, CA, USA) according to the manufacturer's instructions. The quality of the DNA was checked by gel electrophoresis and quantified using NanoDrop spectrophotometer. DNA samples were stored at -80°C. The total DNA samples were used as templates using primers F479 and R888 for V4-V5 region of 16S rRNA amplicons and primers FF390 and FR1 for V7-V8 region of 18S rRNA amplicons [47]. For 16S rRNA gene fragment, 5 ng of template were used for a 25µl of PCR conducted under the following conditions: 94°C for 2 min, 35 cycles of 30 s at 94°C, 52°C for 30 s and 72°C for 1min, following by 7

min at 72°C. Similarly, 18S rRNA gene fragment was amplified under the following PCR conditions: 94°C for 3 min, 35 cycles of 1 min at 94°C, 52°C for 1 min and 72°C for 1 min, following by 5 min at 72°C. For both, PCR was performed using Phusion High-Fidelity DNA Polymerase (BioLabs). The PCR products were purified using QIAquick PCR Purification Kit (QIAGEN) and quantified using the PicoGreen staining Kit. Next steps were performed by Genome Québec platform: a second PCR of nine cycles was conducted twice for each sample under similar PCR conditions with purified PCR products and 10 base pair multiplex identifiers added to the primers at 5' position to specifically identify each sample and avoid PCR bias. Finally, PCR products were purified and quantified. The high-throughput sequencing was performed on a MiSeq platform (Illumina, San Diego, CA, USA) by Genome Québec platform. All Illumina sequence data from this study were submitted to the European Nucleotide Archive (ENA) under accession number PRJEB34506.

### Bioinformatic analysis of 16S and 18S rRNA gene sequences

The raw 16S rRNA gene and 18S rRNA gene sequences were processed using Qiime 2 (version 2019.1) [48]. After performing a quality screening, we used DADA2 [49] to process the raw sequences into exact sequence variants (ESVs). DADA2 resolves biological sequences at the highest resolution (level of single-nucleotide differences) that correspond to the best proxy to identify microbial species [18]. The DADA2 workflow performs filtering, dereplication, chimera identification, and the merging of paired-end reads. The taxonomic affiliations were assigned using the SILVA 132 database [50] (Quast et al. 2012) for the 16S rRNA gene sequences and the Protist Ribosomal Reference database (PR<sup>2</sup>) [51] for the 18S rRNA gene sequences. ESVs present in only one sample and with less than 10 observations in the entire dataset were excluded. ESVs affiliated to chloroplasts, mitochondria (16S rRNA gene dataset), to Chloroplastida or to Animalia (18S rRNA gene dataset) were removed in order to keep only microbial ESVs. We used negative controls and the *Decontam* package [52] in R 3.5.2 [53] to identify contaminant sequences from reagents or introduced during the manipulation of the samples. Four 16S rRNA gene ESVs and seventeen 18S rRNA gene ESVs were identified as contaminant and removed from the dataset. For the 16S rRNA gene dataset, a total of 5145 ESVs and 1455888 reads were present in the final dataset. This prokaryotic dataset (bacteria and archaea) was rarefied to 1018 sequences per sample and 8 samples had to be excluded due to low read numbers after removing non-microbial sequences. For the 18S rRNA gene dataset, a total of 1409 ESVs and 1281880 reads were present in the final dataset. This eukaryotic dataset (micro-eukaryotes) was rarefied to 1127 sequences per sample and 6 samples had to be excluded due to low read numbers after removing non-microbial sequences. Phylogenetic trees of the 16S and 18S rRNA gene ESVs were prepared in Qiime2 (function *qiime phylogeny align-to-tree-mafft-fasttree*) and their visualizations were performed using iTOL [54].

## Diversity and core microbiome analyses

We explored the effects of the wheat genotype, agricultural practices and soil on the alpha diversity of the rhizosphere microbiome by calculating the observed ESV richness and Faith's phylogenetic diversity (R package *picante*) [55]. The effects on microbiome community structure were investigated based on a Bray-Curtis distance matrix visualized using Non-metric Multi-Dimensional Scaling (NMDS, *metaMDS* function) and associated to a permutational multivariate analysis of variance (*adonis* function, 999 permutations) in the R package *vegan* [56]. The core taxa representative of the eight African and European soils used in this study were identified based on a criterion of prevalence in at least 25% of the samples (i.e. presence in a minimum of 16 of the 64 samples) with no criterion related to the relative abundance of the taxa, to consider rare but prevalent microbial taxa (function *core\_members* in *microbiome* package) [57]. Based on this criterion, a list of core taxa was identified and the number and cumulative relative abundance of these core taxa in each sample were calculated. This prevalence criterion enabled the identification of the wheat core microbiome specific of Africa, of Europe and common to both continents. The taxonomic affiliation of the core taxa was verified and refined using nucleotide basic local alignment search tool (BLASTN) analyses.

We inferred cross-domain interaction networks of the wheat core taxa using the SParse InversE Covariance estimation for Ecological Association Inference (*SPIEC-EASI*, version 1.0.2) package in R [58,59]. We identified hub taxa that may act as potential keystone taxon in an ecological network, following the approach used in Tipton et al. (2018) [59] and Agler et al. (2016) [60]. Hub taxa were identified based on their high connectivity and centrality within the network compare to the other taxa using three node parameters: node degree (number of correlations with other taxa), betweenness centrality and closeness centrality. The node parameters were determined using the R package *igraph* (version 1.2.4) [61] and the visualization of the network was performed with the package *ggnetwork* (version 0.1) [62].

The statistical effects of the wheat genotype, agricultural practices and/or soil (and their interaction, when applicable) on the univariate endpoints (e.g. ESV richness, Faith's phylogenetic diversity, number of core taxa) were assessed using generalized linear models (*glm* function in package *lme4*) and *post hoc* comparisons were performed using the Tukey method (*warp.emm* function in package *emmeans*). Linear regressions using Spearman correlation test were conducted to investigate relationships between the prokaryotic (16S rRNA gene dataset) and eukaryotic (18S rRNA gene dataset) ESV richness or between the number of core taxa and the relative abundance of core taxa per. Heatmaps were performed using the R package *pheatmap* and scatter plots were prepared using the package *ggplot2* (version 3.1.1) [63].

## Acknowledgements

We thank Dr. Jacques Le Gouis from INRA Clermont-Ferrand (UMR GDEC) for help in sampling French soils and for providing several wheat seeds cultivars. We thank Dr Honore Tekeu, M. Yves Tchietchoua and Ms Adrienne Ngo Ngom for soil sampling at Tubah area in Cameroon and Institute of Agricultural Research for Development (IRAD) that hosted all activities of the wheat research project in Cameroon. We also thank Dr. Fatou NDOYE and Ms Achaitou DJIBO WAZIRI for Senegalese soils sampling. For lab assistance, we thank Isabelle Rimbault and Nathalia Arias Rojas.

## Funding

This work was supported by the MIC-CERES (“Microbial eco-compatible strategies for improving wheat quality traits and rhizospheric soil sustainability”) Project (FC Project ID 2013–1888; AF Project ID 1301–003) jointly supported by Agropolis Fondation (through the “Investissements d’avenir” programme with reference number ANR-10-LABX-0001-01) and Fondazione Cariplo. Marie Simonin was supported by an IRD postdoctoral fellowship. Italian group was partially funded by BIOPRIME Mipaaf project.

## Availability of data and materials

The raw amplicon sequencing data are available on the European Nuclotide Archive (ENA) with the accession number PRJEB34506: <http://www.ebi.ac.uk/ena/data/view/PRJEB34506>. The code used for the bioinformatic analyses to process the amplicon sequencing data and for generating the figures in R is available in Additional files 2 and 3 as .Rmd files. All input files to generate the figures are available in Additional file 3.

## Additional files

**Additional file 1:** An XLSX table containing a tab with the full list of the 179 core taxa with taxonomic, prevalence, relative abundance and sequence information. A second tab presents the list of taxa found to be associated to 7-8 wheat genotypes in the FR2 soil and the third tab provides the list of taxa significantly affected by farming practices.

**Additional file 2:** A zip archive containing the command line code (.Rmd file) and metadata files needed to perform bioinformatic analysis.

**Additional file 3:** A zip archive with the R script (.Rmd file), a html version of the script with the figures associated to each code chunk and all input files needed to reproduce the figures in R.

## Competing interests

The authors declare that they have no competing interests

## References


1. FAO. How to Feed the World in 2050. Rome, Italy: Food and Agriculture Organization; 2009.
2. Schröder P, Sauvêtre A, Gnädinger F, Pesaresi P, Chmeliková L, Doğan N, et al. Discussion paper: Sustainable increase of crop production through improved technical strategies, breeding and adapted management—A European perspective. *Sci Total Environ.* 2019;678:146–161.
3. Schlaeppli K, Bulgarelli D. The plant microbiome at work. *Mol Plant Microbe Interact.* 2015;28:212–217.
4. Turner TR, James EK, Poole PS. The plant microbiome. *Genome Biol.* 2013;14:209.
5. Bender SF, Wagg C, van der Heijden MGA. An Underground Revolution: Biodiversity and Soil Ecological Engineering for Agricultural Sustainability. *Trends Ecol Evol.* 2016;31:440–52.
6. Donn S, Kirkegaard JA, Perera G, Richardson AE, Watt M. Evolution of bacterial communities in the wheat crop rhizosphere. *Environ Microbiol.* 2015;17:610–21.
7. Yin C, Mueth N, Hulbert S, Schlatter D, Paulitz TC, Schroeder K, et al. Bacterial Communities on Wheat Grown Under Long-Term Conventional Tillage and No-Till in the Pacific Northwest of the United States. *Phytobiomes J.* 2017;1:83–90.
8. Mavrodi DV, Mavrodi OV, Elbourne LD, Tetu S, Bonsall RF, Parejko J, et al. Long-term irrigation affects the dynamics and activity of the wheat rhizosphere microbiome. *Front Plant Sci.* 2018;9:345.
9. Hartman K, van der Heijden MGA, Wittwer RA, Banerjee S, Walser J-C, Schlaeppli K. Cropping practices manipulate abundance patterns of root and soil microbiome members paving the way to smart farming. *Microbiome.* 2018;6:14.
10. Sapkota R, Knorr K, Jørgensen LN, O'Hanlon KA, Nicolaisen M. Host genotype is an important determinant of the cereal phyllosphere mycobiome. *New Phytol.* 2015;207:1134–44.
11. Granzow S, Kaiser K, Wemheuer B, Pfeiffer B, Daniel R, Vidal S, et al. The effects of cropping regimes on fungal and bacterial communities of wheat and faba bean in a greenhouse pot experiment differ between plant species and compartment. *Front Microbiol.* 2017;8:902.
12. Lu T, Ke M, Peijnenburg WJGM, Zhu Y, Zhang M, Sun L, et al. Investigation of Rhizospheric Microbial Communities in Wheat, Barley, and Two Rice Varieties at the Seedling Stage. *J Agric Food Chem.* 2018;66:2645–53.
13. Gao Z, Karlsson I, Geisen S, Kowalchuk G, Jousset A. Protists: Puppet Masters of the Rhizosphere Microbiome. *Trends Plant Sci.* 2019;24:165–76.
14. Mahoney AK, Yin C, Hulbert SH. Community structure, species variation, and potential functions of rhizosphere-associated bacteria of different winter wheat (*Triticum aestivum*) cultivars. *Front Plant Sci.* 2017;8:132.
15. Fan K, Cardona C, Li Y, Shi Y, Xiang X, Shen C, et al. Rhizosphere-associated bacterial network structure and spatial distribution differ significantly from bulk soil in wheat crop fields. *Soil Biol Biochem.* 2017;113:275–84.
16. Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, et al. Defining the core *Arabidopsis thaliana* root microbiome. *Nature.* 2012;488:86–90.

17. Edwards J, Johnson C, Santos-Medellín C, Lurie E, Podishetty NK, Bhatnagar S, et al. Structure, variation, and assembly of the root-associated microbiomes of rice. *Proc Natl Acad Sci.* 2015;112:E911–20.
18. Edgar RC. Updating the 97% identity threshold for 16S ribosomal RNA OTUs. *Bioinformatics.* 2018;34:2371–5.
19. Callahan BJ, McMurdie PJ, Holmes SP. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME J.* 2017;11:2639.
20. Corneo PE, Suenaga H, Kertesz MA, Dijkstra FA. Effect of twenty four wheat genotypes on soil biochemical and microbial properties. *Plant Soil.* 2016;404:141–55.
21. Gdanetz K, Trail F. The Wheat Microbiome Under Four Management Strategies, and Potential for Endophytes in Disease Protection. *Phytobiomes J.* 2017;1:158–68.
22. Venter ZS, Jacobs K, Hawkins H-J. The impact of crop rotation on soil microbial diversity: A meta-analysis. *Pedobiologia.* 2016;59:215–223.
23. Hamonts K, Trivedi P, Garg A, Janitz C, Grinyer J, Holford P, et al. Field study reveals core plant microbiota and relative importance of their drivers. *Environ Microbiol.* 2018;20:124–40.
24. Delgado-Baquerizo M, Oliverio AM, Brewer TE, Benavent-González A, Eldridge DJ, Bardgett RD, et al. A global atlas of the dominant bacteria found in soil. *Science.* 2018;359:320–5.
25. Vandenkoornhuyse P, Quaiser A, Duhamel M, Van AL, Dufresne A. The importance of the microbiome of the plant holobiont. *New Phytol.* 2015;206:1196–206.
26. Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J, et al. FUNGuild: An open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecol.* 2016;20:241–8.
27. Zhang H, Wu X, Li G, Qin P. Interactions between arbuscular mycorrhizal fungi and phosphate-solubilizing fungus (*Mortierella* sp.) and their effects on *Kosteletzkya virginica* growth and enzyme activities of rhizosphere and bulk soils at different salinities. *Biol Fertil Soils.* 2011;47:543.
28. Schlatter DC, Yin C, Burke I, Hulbert S, Paulitz T. Location, Root Proximity, and Glyphosate-Use History Modulate the Effects of Glyphosate on Fungal Community Networks of Wheat. *Microb Ecol.* 2018;76:240–57.
29. Borrell AN, Shi Y, Gan Y, Bainard LD, Germida JJ, Hamel C. Fungal diversity associated with pulses and its influence on the subsequent wheat crop in the Canadian prairies. *Plant Soil.* 2017;414:13–31.
30. Fan K, Weisenhorn P, Gilbert JA, Chu H. Wheat rhizosphere harbors a less complex and more stable microbial co-occurrence pattern than bulk soil. *Soil Biol Biochem.* 2018;125:251–60.
31. Ofek M, Hadar Y, Minz D. Ecology of Root Colonizing Massilia (Oxalobacteraceae). *PLOS ONE.* 2012;7:e40117.
32. Li X, Rui J, Mao Y, Yannarell A, Mackie R. Dynamics of the bacterial community structure in the rhizosphere of a maize cultivar. *Soil Biol Biochem.* 2014;68:392–401.
33. Tourna M, Stieglmeier M, Spang A, Könneke M, Schintlmeister A, Urich T, et al. Nitrososphaera viennensis, an ammonia oxidizing archaeon from soil. *Proc Natl Acad Sci.* 2011;108:8420–5.


34. Pester M, Rattei T, Flechl S, Gröngröft A, Richter A, Overmann J, et al. amoA-based consensus phylogeny of ammonia-oxidizing archaea and deep sequencing of amoA genes from soils of four different geographic regions. *Environ Microbiol*. 2012;14:525–39.
35. Simonin M, Richaume A, Guyonnet JP, Dubost A, Martins JMF, Pommier T. Titanium dioxide nanoparticles strongly impact soil microbial function by affecting archaeal nitrifiers. *Sci Rep*. 2016;6:33643.
36. Carvalhais LC, Dennis PG, Badri DV, Kidd BN, Vivanco JM, Schenk PM. Linking Jasmonic Acid Signaling, Root Exudates, and Rhizosphere Microbiomes. *Mol Plant Microbe Interact*. 2015;28:1049–58.
37. Walters WA, Jin Z, Youngblut N, Wallace JG, Sutter J, Zhang W, et al. Large-scale replicated field study of maize rhizosphere identifies heritable microbes. *Proc Natl Acad Sci*. 2018;115:7368–73.
38. Sapp M, Ploch S, Fiore-Donno AM, Bonkowski M, Rose LE. Protists are an integral part of the *Arabidopsis thaliana* microbiome. *Environ Microbiol*. 2018;20:30–43.
39. Zhao Z-B, He J-Z, Geisen S, Han L-L, Wang J-T, Shen J-P, et al. Protist communities are more sensitive to nitrogen fertilization than other microorganisms in diverse agricultural soils. *Microbiome*. 2019;7:33.
40. Durán P, Thiergart T, Garrido-Oter R, Agler M, Kemen E, Schulze-Lefert P, et al. Microbial Interkingdom Interactions in Roots Promote *Arabidopsis* Survival. *Cell*. 2018;175:973–983.e14.
41. Hassani MA, Durán P, Hacquard S. Microbial interactions within the plant holobiont. *Microbiome*. 2018;6:58.
42. Richards LA. Diagnosis and Improvement of Saline and Alkali Soils. *Soil Sci*. 1954;78:154.
43. Bouyoucos GJ. A Recalibration of the Hydrometer Method for Making Mechanical Analysis of Soils 1. *Agron J*. 1951;43:434–8.
44. Pétard J. Les méthodes d'analyse. *Anal Sols*. 1993;
45. Hurek T, Reinhold-Hurek B, Montagu MV, Kellenberger E. Root colonization and systemic spreading of *Azoarcus* sp. strain BH72 in grasses. *J Bacteriol*. 1994;176:1913–23.
46. Miché L, Balandreau J. Effects of Rice Seed Surface Sterilization with Hypochlorite on Inoculated *Burkholderia vietnamensis*. *Appl Environ Microbiol*. 2001;67:3046–52.
47. Terrat S, Plassart P, Bourgeois E, Ferreira S, Dequiedt S, Adele-Dit-De-Renseville N, et al. Meta-barcoded evaluation of the ISO standard 11063 DNA extraction procedure to characterize soil bacterial and fungal community diversity and composition. *Microb Biotechnol*. 2015;8:131–42.
48. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol*. 2019;1.
49. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods*. 2016;13:581–3.
50. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res*. 2012;41:D590–D596.

51. Guillou L, Bachar D, Audic S, Bass D, Berney C, Bittner L, et al. The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with curated taxonomy. *Nucleic Acids Res.* 2013;41:D597–604.
52. Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data. *Microbiome.* 2018;6:226.
53. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria; 2015; Available from: <http://www.R-project.org/>
54. Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Res.* 2016;44:W242–5.
55. Kembel SW, Cowan PD, Helmus MR, Cornwell WK, Morlon H, Ackerly DD, et al. Picante: R tools for integrating phylogenies and ecology. *Bioinformatics.* 2010;26:1463–4.
56. Oksanen J, Kindt R, Legendre P, O’Hara B, Stevens MHH, Oksanen MJ, et al. The vegan package. *Community Ecol Package.* 2007;
57. Lahti L, Shetty S, Blake T. Tools for microbiome analysis in R. *Microbiome Package Version 099.* 2017;88:2012–2017.
58. Kurtz ZD, Müller CL, Miraldi ER, Littman DR, Blaser MJ, Bonneau RA. Sparse and Compositionally Robust Inference of Microbial Ecological Networks. *PLOS Comput Biol.* 2015;11:e1004226.
59. Tipton L, Müller CL, Kurtz ZD, Huang L, Kleerup E, Morris A, et al. Fungi stabilize connectivity in the lung and skin microbial ecosystems. *Microbiome.* 2018;6:12.
60. Agler MT, Ruhe J, Kroll S, Morhenn C, Kim S-T, Weigel D, et al. Microbial Hub Taxa Link Host and Abiotic Factors to Plant Microbiome Variation. *PLOS Biol.* 2016;14:e1002352.
61. Csardi G, Nepusz T. The igraph software package for complex network research. *InterJournal Complex Syst.* 2006;1695:1–9.
62. Tyner SC, Briatte F, Hofmann H. Network Visualization with ggplot2. *R J [Internet].* 2017 [cited 2019 Aug 2]; Available from: <https://hal.archives-ouvertes.fr/hal-01722543>
63. Wickham H. *ggplot2: Elegant Graphics for Data Analysis.* Springer; 2016.


## Supplementary figures




**Fig S1: Results of the Sub-experiment 1: Effects of the wheat genotypes on the prokaryotic and eukaryotic microbial communities. Top: Observed ESVs richness, Middle: Faith's phylogenetic diversity, Bottom: Community Structure represented in a NMDS**



**Fig S2: Results of the Sub-experiment 2: Effects of the soil and wheat genotypes on the prokaryotic and eukaryotic microbial communities. Top: Observed ESV richness, Middle: Faith's phylogenetic diversity, Bottom: Community Structure represented in a NMDS**



**Fig S3: Taxa significantly enriched in Conventional or Organic agricultural practices in the French and Italian soil. Each taxa Log2 fold change relative to the other agricultural condition is indicated on the x-axis.**



**Fig S4: Correlation between the number of core taxa and the cumulative relative abundance of the core taxa per sample in the eight soils. The size of the points indicates the total ESV richness in the sample (prokaryotes + eukaryotes).**