

## **Heterozygous ATP-binding Cassette Transporter G5 Gene Deficiency and Risk of Coronary Artery Disease**

Short title: Heterozygous *ABCG5* deficiency and risk of CAD

Akihiro Nomura\*, MD PhD, Connor A. Emdin\*, DPhil, Hong Hee Won, PhD, Gina M. Peloso, PhD, Pradeep Natarajan, MD, Diego Ardissino, MD, John Danesh, FRCP DPhil, Heribert Schunkert, MD, Adolfo Correa, MD PhD, Matthew J. Bown, MD FRCS, Nilesh J. Samani, MD FRCP, Jeanette Erdmann, PhD, Ruth McPherson, MD, Hugh Watkins, MD PhD, Danish Saleheen, MD, Roberto Elosua, MD PhD, Masa-aki Kawashiri, MD PhD, Hayato Tada, MD PhD, Namrata Gupta, PhD, Svat H. Shah, MD MHS, Daniel J. Rader, MD, Stacey Gabriel, PhD, Amit V. Khera\*, MD, Sekar Kathiresan\*, MD

\*: These authors contributed equally

**Address for correspondence:**  
Sekar Kathiresan, MD  
Verve Therapeutics  
26 Landsdowne Street, 1<sup>st</sup> Floor  
Cambridge, MA 02139  
Email: skathiresan1@mgh.harvard.edu  
Phone: 617 603 0070

## Abstract

**Background:** Familial sitosterolemia is a rare, recessive Mendelian disorder characterized by hyperabsorption and decreased biliary excretion of dietary sterols. Affected individuals typically have complete genetic deficiency – homozygous loss-of-function (LoF) variants — in the ATP-binding cassette transporter G5 (*ABCG5*) or G8 (*ABCG8*) genes, and have substantially elevated plasma sitosterol and low-density lipoprotein cholesterol (LDL-C) levels. The impact of partial genetic deficiency of *ABCG5* or *ABCG8*, as occurs in heterozygous carriers of LoF variants, on LDL-C and risk of coronary artery disease (CAD) has remained uncertain.

**Methods:** We first recruited nine sitosterolemia families, identified causative LoF variants in *ABCG5* or *ABCG8*, and evaluated the associations of these *ABCG5* or *ABCG8* LoF variants with plasma phytosterols and lipid levels. We next assessed for LoF variants in *ABCG5* or *ABCG8* in CAD cases (n=29,361) versus controls (n=357,326). We tested the association of rare LoF variants in *ABCG5* or *ABCG8* with blood lipids and risk for CAD. Rare LoF variants were defined as protein-truncating variants with minor allele frequency less than 0.1% in *ABCG5* or *ABCG8*.

**Results:** In sitosterolemia families, seven pedigrees harbored causative LoF variants in *ABCG5* and two pedigrees in *ABCG8*. Homozygous LoF variants in either *ABCG5* or *ABCG8* led to marked elevations in sitosterol and LDL-C. Of those sitosterolemia families, heterozygous carriers of *ABCG5* LoF variants exhibited increased sitosterol and LDL-C levels compared to non-carriers. Within the large-scale CAD case-control cohorts, prevalence of rare LoF variants in *ABCG5* and in *ABCG8* were approximately 0.1% each. *ABCG5* heterozygous LoF variant carriers had significantly elevated LDL-C levels (24.7 mg/dL; 95% confidence interval [CI] 14 to 35; P=1.1x10<sup>-6</sup>), and were at two-fold increased risk of CAD (odds ratio 2.06, 95% CI 1.27 to 3.35; P=0.004). By contrast, *ABCG8* heterozygous LoF carrier status was not associated with increased LDL-C or risk of CAD.

**Conclusions:** Although familial sitosterolemia is traditionally considered as a recessive disorder, we observed that heterozygous carriers of a LoF variant in *ABCG5* had significantly increased sitosterol and LDL-C levels and a two-fold increase in risk of CAD.

## Introduction

Familial sitosterolemia (OMIM #210250) is a rare Mendelian recessive disorder characterized by tendonous xanthomas<sup>1</sup>, high plasma plant sterols and cholesterol levels, and increased risk of premature myocardial infarction.<sup>2-4</sup> The ATP-binding cassette transporters G5 (*ABCG5*) and G8 (*ABCG8*) are primary causal genes of familial sitosterolemia. *ABCG5*, *ABCG8*, and N-terminal Niemann-Pick C1 Like 1 (*NPC1L1*)<sup>5, 6</sup> determine the efflux and absorption of sterols on the surface of intestine and bile duct.<sup>7, 8</sup> *NPC1L1* regulates sterol absorption whereas *ABCG5* and *ABCG8* form obligate heterodimers<sup>9</sup> and coordinately control the excretion at both the brush border membrane of enterocyte and the apical membrane of hepatocytes.<sup>5, 10-12</sup>

Complete deficiency due to homozygous or compound heterozygous loss-of-function (LoF) variants in *ABCG5* and/or *ABCG8* causes markedly increased sitosterolemia and cholesterol levels, and potentially accelerated atherosclerotic disease as well.<sup>1-4</sup> Genome-wide association studies also demonstrated that loci at the *ABCG5-ABCG8* gene region were associated with phytosterols, low-density lipoprotein (LDL) cholesterol,<sup>13</sup> and risk of coronary artery disease (CAD).<sup>14</sup> However, it is uncertain whether partial deficiency of *ABCG5* or *ABCG8* as conferred by LoF variants are also associated with higher cholesterol levels and an increased risk of CAD.

Here, we explored the metabolic consequences of *ABCG5* or *ABCG8* deficiencies. We recruited probands and relatives in sitosterolemia families and assessed whether observed *ABCG5* or *ABCG8* causative LoF variants were associated with increased plasma phytosterols and LDL cholesterol. We then analyzed exome sequences from 112,700 participants and genotype data from an additional 293,134 individuals to test whether carriers of rare heterozygous LoF variants in *ABCG5* or *ABCG8* had elevated blood lipids and risk of CAD.

## Methods

### Study Participants

Nine sitosterolemia pedigrees with phytosterol and lipid profiles derived from the Kanazawa University Mendelian Disease (KUMD) Registry were evaluated for *ABCG5* and *ABCG8* variants (**Supplemental Figure 1**).<sup>3</sup> Sitosterolemia was diagnosed by 1) plasma sitosterol concentration  $\geq 10$   $\mu\text{g/mL}$  and 2) presence of tendon or tuberous xanthomas and/or history of premature CAD.<sup>15</sup> Causative LoF variants in the pedigrees were defined as pathogenic or likely pathogenic by the American College of Medical Genetics (ACMG) standard criteria,<sup>16</sup> and were identified by Sanger sequencing or whole exome sequencing (WES). Controls were unaffected relatives without any causative LoF variants in the sitosterolemia pedigrees.

Next, *ABCG5* and *ABCG8* were sequenced in the Myocardial Infarction Genetics consortium (MIGen), UK Biobank and the TruSeq Custom Amplicon target resequencing (TSCA) studies. MIGen case control studies included the Italian Atherosclerosis Thrombosis and Vascular Biology (ATVB) study,<sup>17</sup> the Deutsches Herzzentrum München Myocardial Infarction Study (DHM),<sup>18</sup> the Exome Sequencing Project Early-Onset Myocardial Infarction study (ESP-EOMI),<sup>19</sup> the Jackson Heart Study (JHS),<sup>20</sup> the Leicester Acute Myocardial Infarction Study (Leicester),<sup>21</sup> the Lubeck Myocardial Infarction Study (Lubeck),<sup>22</sup> the Ottawa Heart Study (OHS),<sup>23</sup> the Precocious Coronary Artery Disease (PROCARDIS) study,<sup>24</sup> the Pakistan Risk of Myocardial Infarction Study (PROMIS),<sup>25</sup> and the Registre Gironi del COR (Gerona Heart Registry or REGICOR) study.<sup>26</sup> TSCA included the Duke CATHGEN study (DUKE),<sup>27</sup> the MedStar study (MedStar),<sup>28</sup> and the PennCath study (PennCath).<sup>28</sup> An additional 293,134 individuals in UK Biobank underwent array based genotyping for the *ABCG5* stop variant rs199689137 (p.R446Ter) and were included in the analysis.

All participants in each study provided written informed consent for genetic studies. The institutional review board at Partners HealthCare (Boston, MA, USA) and each participating institution approved the study protocol.

### Lipid measurement and coronary artery disease ascertainment

In sitosterolemia pedigree-based analysis, all blood samples were obtained after a 12-hour overnight fast, either before initiation of lipid-lowering treatment or after discontinuation of medication for at least 4 weeks. Plasma levels of non-cholesterol sterols were determined using gas-liquid chromatography–mass spectrometry.<sup>4</sup> In addition to absolute non-cholesterol sterol levels, cholesterol adjusted ratios (each non-cholesterol sterol level to total cholesterol [TC] level ratio) were also evaluated. Plasma concentrations of TC, triglyceride, and HDL cholesterol were determined in MIGen, TSCA and UK Biobank using enzymatic assays. In MIGen and TSCA, LDL

cholesterol level was calculated using the Friedewald equation for those with triglycerides <400 mg/dL. In the UK Biobank, LDL cholesterol levels were directly measured using an antibody-based assay.

In MiGen, CAD cases were identified as early-onset (premature) myocardial infarction (defined as  $\leq$  50 years old in male and  $\leq$  60 years old in female). In the UK Biobank, CAD cases were defined as myocardial infarction at any age. In TSCA, coronary artery disease was defined as stenosis on angiography ( $\geq$  1 coronary vessel with  $>$  50% stenosis) at ages  $\leq$  50 years old for men and  $\leq$  60 years old for women.

### Exome sequencing

WES or targeted sequencing for KUMD, MiGen, and TSCA was performed at the Broad Institute as previously described.<sup>5,6</sup> In brief, genomic deoxyribonucleic acid was captured on protein-coding regions using the NimbleGen Sequencing Capture Array or the Illumina TrueSeq Custom Amplicon. Sequencing reads were aligned to a human reference genome (build 37) using the Burrows–Wheeler Aligner-Maximal Exact Match algorithm. Aligned non-duplicate reads were locally realigned, and base qualities were recalibrated using the Genome Analysis ToolKit (GATK) software.<sup>29</sup> Variants were jointly called using the GATK HaplotypeCaller software. WES in UK Biobank was performed by the Regeneron Genetics Center as previously described.<sup>7</sup>

In large cohort studies, rare LoF variants were defined as those with minor allele frequency less than 0.1% and which caused: (1) insertions or deletions of DNA that modified the reading frame of protein translation (frameshift); (2) point mutations at conserved splice site regions that altered the splicing process (splice-site); or (3) point mutations that changed an amino acid codon to a stop codon, leading to the truncation of a protein (nonsense). LoF variants were identified using the LOFTEE plugin of the Variant Effect Predictor software (version 82).<sup>8,9</sup>

### Statistical Analysis

In sitosterolemia family-based analyses, the differences in lipid levels, lipoproteins and non-cholesterol sterols by *ABCG5* or *ABCG8* LoF variant carrier status were assessed using the *Mann-Whitney U* test. The effects of *ABCG5* and *ABCG8* rare LoF variants on lipid profiles in MiGen, UK Biobank and TSCA was evaluated using linear regression, adjusting for age, gender, study, and first five principal components of ancestry. A Cochran–Mantel–Haenszel statistic meta-analysis for stratified 2-by-2 tables was used to associate *ABCG5* and *ABCG8* rare LoF variants with risk of CAD. As a sensitivity analysis, we performed an inverse variance weighted fixed effects meta-analysis of the adjusted odds ratio, derived in each cohort using logistic regression adjusted for age, sex, study and five principal components of ancestry. P values of less than 0.025 were considered to indicate statistical significance (i.e., Bonferroni correction for the testing of two genes). Statistical

analyses were performed using R software version 3.4.3 (The R Project for Statistical Computing, Vienna, Austria).


## Results

### ***ABCG5 or ABCG8 causative LoF variants, blood phytosterol and cholesterol levels in sitosterolemia families***

We recruited nine Japanese families with sitosterolemia and sequenced the exons of the *ABCG5* and *ABCG8* genes in 47 individuals from these families (**Supplemental Figure 1**). Among the individuals within these families, 9 carried a homozygous or compound heterozygous *ABCG5* or *ABCG8* causative LoF variants while 28 carried a heterozygous *ABCG5* or *ABCG8* LoF causative variants. Most of those LoF variants were considered as pathogenic protein truncating or missense variants by the ACMG standard criteria (**Supplemental Table 1**). As expected, *ABCG5* and *ABCG8* homozygote or compound heterozygous LoF variant carriers showed very high sitosterol / TC ratio and LDL cholesterol levels compared to non-carriers. Regarding heterozygous state, carriers of *ABCG5* or *ABCG8* heterozygous LoF variant carriers exhibited increased sitosterol / TC ratio compared with non-carriers. Moreover, *ABCG5* heterozygous LoF variant carrier status was associated with an increased LDL cholesterol level. (**Table 1 and Figure 1**).

**Figure 1. Sitosterol to total cholesterol ratio and LDL cholesterol levels among individuals with homozygous and heterozygous sitosterolemia, and unaffected controls in sitosterolemia families.**

Each dot indicates an individual's value. Each horizontal line represents a mean value for each carrier status. \*:  $P < 0.025$  by *Mann-Whitney U* test.



## ***ABCG5 or ABCG8 rare heterozygous LoF variation, blood lipids and risk for CAD in large cohorts***


Next, we examined whether rare heterozygous LoF variant carrier status in *ABCG5* or *ABCG8* increased blood lipids and were at elevated risk of CAD using large-scale genetics cohorts. We sequenced the protein coding regions of *ABCG5* and *ABCG8* in 112,700 individuals from three datasets: 58,791 participants from MIGen, 52,195 participants in UK Biobank and 1,714 participants from TSCA (**Table 2**). We detected 108 individuals harboring rare *ABCG5* LoF alleles and the prevalence of *ABCG5* heterozygous carrier status was 0.1%. (**Supplemental Table 2**). We also discovered 154 individuals who harbored rare *ABCG8* LoF alleles, a heterozygous carrier prevalence also around 0.1%.

Individuals carrying *ABCG5* LoF variants had significantly increased total cholesterol levels (17 mg/dl; 95% confidence interval [CI], 13 to 32;  $P = 6.9 \times 10^{-6}$ ) and LDL cholesterol levels (25 mg/dl, 95% CI, 13 to 32;  $P = 1.1 \times 10^{-6}$ , **Figure 2**). Then, we investigated the association between rare *ABCG5* heterozygous LoF variant carrier status and CAD risk using more than 380,000 participants from the three WES cohorts and additional UK biobank genotyping array-based cohort. We identified 34 carriers of *ABCG5* heterozygous LoF variants among 29,321 coronary artery disease cases (0.1%) and 63 among 357,326 controls (0.02%). In a Cochran-Mantel-Haenszel fixed-effect meta-analysis, individuals carrying *ABCG5* heterozygous LoF variants were at two-fold risk of CAD (Odds ratio [OR], 2.06; 95% CI, 1.27 to 3.35;  $P$  value = 0.004) (**Figure 3**). A similar effect estimate was noted in a meta-analysis of adjusted odds ratios derived using logistic regression (OR 2.04; 95% CI 1.28 to 3.26;  $P$  = 0.003).

In contrast to *ABCG5*, carriers of rare *ABCG8* heterozygous LoF variants did not exhibit significant increase in any of blood lipids including LDL cholesterol (beta, 0.08; 95% CI, -0.06 to 0.23;  $P = 0.25$ ) (**Supplemental Table 3**). Moreover, *ABCG8* heterozygous LoF variant carrier status was not at elevated risk for CAD (OR, 0.79; 95% CI, 0.47 to 1.35;  $P = 0.39$ ) (**Supplemental Table 3**).


**Figure 2. Effects of loss-of-function variants in *ABCG5* on blood lipid profiles from MiGen and UK Biobank.**

Effect sizes were calculated using linear regression adjusted by age, gender, study, case-control status, and first five principal components of ancestry. Triglyceride was natural log-transformed before analysis. Fixed-effects meta-analysis was applied to combine results.



**Figure 3. Effect of loss-of-function variants in *ABCG5* on myocardial infarction risk.**


A meta-analysis across studies was performed using the Cochran–Mantel–Haenszel statistics for stratified 2-by-2 tables. MIGen, myocardial infarction genetics consortium.



We explored whether the effect size of *ABCG5* LoF variants on CAD risk was consistent with their effect on LDL cholesterol. We observed a linear dose-response relationship between CAD risk and LDL cholesterol change conferred by DNA sequence variants in *LDLR*, *PCSK9*, *ABCG5* and *ABCG8*. The effect of *ABCG5* LoF variants on CAD risk (106% increase in risk) was consistent with the estimate based on the change in LDL cholesterol (25 mg/dl) (**Figure 4**).

**Figure 4. Relationship between effect on LDL cholesterol levels and coronary artery disease risk for *ABCG5*, *ABCG8*, *PCSK9* and *LDLR* loss-of-function variants.**

CAD, coronary artery disease; LDL, low-density lipoprotein



## Discussion

In this study, we evaluated whether rare heterozygous LoF variations in *ABCG5* or *ABCG8* were associated with blood lipid levels and CAD risk. We used two different approaches — sitosterolemia family-based analysis and population-based analyses from over 380,000 individuals — to test whether rare heterozygous LoF variants in *ABCG5* or *ABCG8* associated with phytosterols, lipids and CAD. We found that carriers of heterozygous LoF variants in *ABCG5* had higher sitosterol and ~25 mg/dl higher LDL cholesterol and were at two-fold risk of CAD.

These results permit several conclusions. First, individuals who carried rare heterozygous LoF variants in *ABCG5* had significantly elevated LDL cholesterol levels and were at elevated risk for CAD. Although there have been reports of premature atherosclerosis among patients with homozygous sitosterolemia,<sup>10</sup> it was unclear if *ABCG5* partial deficiency by heterozygous LoF variants also increases blood lipid levels and CAD risk. The findings of this study clearly demonstrated that *ABCG5* partial deficiency increased both LDL cholesterol level and CAD risk. Considering the higher burden of CAD at a population level, these findings suggests that *ABCG5* LoF variant carriers may derive clinical benefit from LDL cholesterol lowering therapy. Importantly, the *NPC1L1* inhibitor ezetimibe is known to reduce intestinal cholesterol and phytosterol absorption in patients with sitosterolemia, and could have increased efficacy in individuals with partial *ABCG5* deficiency.<sup>30</sup>

Second, it has been unclear whether elevated plant sterol levels or elevated blood cholesterol levels cause atherosclerosis among patients with sitosterolemia.<sup>14</sup> The proportional relationship between CAD risk and elevation in LDL cholesterol levels observed among *ABCG5* heterozygous LoF carriers suggests that the substantial increase in blood LDL cholesterol levels among individuals with sitosterolemia causes atherosclerosis (**Figure 3**). These findings are also consistent with a recent meta-analysis that did not observe a significant association between circulating sitosterol levels and risk of cardiovascular disease.<sup>31</sup> Moreover, the effect size of *ABCG5* heterozygous LoF variant carrier status on both blood lipids and CAD risk was consistent with predictions based on known familial hypercholesterolemia and hypobetalipoproteinemia variants (**Figure 4**). This result is consistent with recent studies, and it might support the hypothesis that the main driver of CHD risk in *ABCG5* LoF variant carriers was LDL cholesterol rather than plant sterols.

This study has several limitations. First, functional analyses of each LoF variant were not performed. Consequently, some annotated LoF variants included in this study may not actually cause deficiency in the *ABCG5/G8* proteins to increase plant sterols and blood lipids. This would underestimate of the effect of *ABCG5* and especially *ABCG8* deficiency on blood lipid levels and CAD risk. Second, CAD definition was different among study cohorts. However, the effect direction

among studies was largely consistent and we observed little heterogeneity for the meta-analysis ( $I^2$  squared of 0%).

In conclusion, 0.1% of population carried rare LoF variants in *ABCG5* and these heterozygous carriers of had an elevated sitosterol and LDL cholesterol levels and were at two-fold risk for CAD.

## Author List

Akihiro Nomura\*, MD PhD, Connor A. Emdin\*, DPhil, Hong Hee Won, PhD, Gina M. Peloso, PhD, Pradeep Natarajan, MD, Diego Ardissino, MD, John Danesh, FRCP DPhil, Heribert Schunkert, MD, Adolfo Correa, MD, PhD, Matthew J Bown, FRCS, Nilesh J Samani, FRCP, Jeanette Erdmann, PhD, Ruth McPherson, MD, Hugh Watkins, Danish Saleheen, MD, Roberto Elosua, MD PhD, Masa-aki Kawashiri, MD PhD, Hayato Tada, MD PhD, Namrata Gupta, PhD, Svat H. Shah, MD MHS, Daniel J. Rader, MD, Stacey Gabriel, PhD, Amit V. Khera\*, MD, Sekar Kathiresan\*, MD

\*: These authors contributed equally

## Acknowledgements

We would like to express our gratitude to all the participants and staff of KUMD, MiGen consortium, TSCA, and UK biobank for their outstanding contributions.

## Source of Funding

This study was funded by the National Institutes of Health (R01 HL127564). Exome sequencing in ATVB, PROCARDIS, OHS, PROMIS, Leicester, Lubeck was supported by 5U54HG003067 to SG. The ATVB Study was supported by a grant from RFPS-2007-3-644382 and Programma di ricerca Regione-Università 2010-2012 Area 1–Strategic Programmes–Regione Emilia-Romagna. Funding for the ESP-EOMI was provided by RC2 HL103010 (HeartGO), RC2 HL102923 (LungGO), and RC2 HL102924 (WHISP). Exome sequencing was performed through RC2 HL102925 (BroadGO) and RC2 HL102926 (SeattleGO). The JHS is supported and conducted in collaboration with Jackson State University (HHSN268201800013I), Tougaloo College (HHSN268201800014I), the Mississippi State Department of Health (HHSN268201800015I) and the University of Mississippi Medical Center (HHSN268201800010I, HHSN268201800011I and HHSN268201800012I) contracts

from the National Heart, Lung, and Blood Institute (NHLBI) and the National Institute on Minority Health and Health Disparities (NIMHD). The authors also wish to thank the staffs and participants of the JHS. REGICOR study was supported by the Spanish Ministry of Economy and Innovation through the Carlos III Health Institute (Red Investigación Cardiovascular RD12/0042, PI09/90506), European Funds for Development (ERDF-FEDER), and by the Catalan Research and Technology Innovation Interdepartmental Commission (2014SGR240). Samples for the Leicester cohort were collected as part of projects funded by the British Heart Foundation (British Heart Foundation Family Heart Study, RG2000010; UK Aneurysm Growth Study, CS/14/2/30841) and the National Institute for Health Research (NIHR Leicester Cardiovascular Biomedical Research Unit Biomedical Research Informatics Centre for Cardiovascular Science, IS\_BRU\_0211\_20033). GMP is supported by K01HL125751 and R03HL141439.

## Disclosures

A.N. received consulting fees from CureApp Inc. and speaker fees from Daiichi Sankyo, Kowa, and Otsuka pharmaceutical. He is a co-founder of CureApp Institute. C.A.E. reports personal fees from Navitor Pharma and Novartis. H.T. received honoraria from Astellas Pharma, Amgen Astellas BioPharma, Bayer Japan, Boehringer Ingelheim, Daiichi Sankyo, Kowa, Mitsubishi Tanabe Pharma Corporation, MSD, Sanofi, Sanwa Kagaku Kenkyusho, Sumitomo Dainippon Pharma, and Takeda Pharmaceutical. M.K. received honoraria from Amgen Astellas Biopharma, Astellas Pharma, Daiichi Sankyo, Kowa Pharmaceutical, MSD, Pfizer Japan and Sanofi. A.V.K. has served as a consultant or received honoraria from Color Genomics, Illumina, and Navitor Pharmaceuticals, received grant support from the Novartis Institute for Biomedical Research and IBM Research, and reports a patent related to a genetic risk predictor (20190017119). S.K. is an employee of Verve Therapeutics. He is a founder of Maze Therapeutics, Verve Therapeutics, and San Therapeutics. He holds equity in Catabasis and San Therapeutics. He is a member of the scientific advisory boards for Regeneron Genetics Center and Corvidia Therapeutics; served as a consultant for Acceleron, Eli Lilly, Novartis, Merck, Novo Nordisk, Novo Ventures, Ionis, Alnylam, Aegerion, Huag Partners, Noble Insights, Leerink Partners, Bayer Healthcare, Illumina, Color Genomics, MedGenome, Quest, and Medscape; and reports patents related to a method of identifying and treating a person having a predisposition to or afflicted with cardiometabolic disease (20180010185) and a genetic risk predictor (20190017119). The remaining authors have nothing to disclose.

## Affiliations

Department of Cardiovascular Medicine, Kanazawa University Graduate School of Medical Sciences, Kanazawa, Japan. (A.N., M.K., H.T.); Innovative Clinical Research Center, Kanazawa University (iCREK), Kanazawa, Japan. (A.N.); Center for Genomic Medicine, Massachusetts General Hospital, Boston, Massachusetts, USA. (C.A.E., P.N., N.G., S.G., A.V.K., S.K.); Cardiovascular Disease Initiative, Broad Institute, Cambridge, Massachusetts, USA. (C.A.E., P.N., A.V.K., S.K.); Department of Medicine, Massachusetts General Hospital, Boston, Massachusetts, USA. (C.A.E., P.N., A.V.K., S.K.); Samsung Advanced Institute for Health Sciences and Technology, Sungkyunkwan University, Samsung Medical Center, Seoul, Republic of Korea. (H.H.W.); Department of Biostatistics, Boston University School of Public Health, Boston, Massachusetts, USA. (G.M.P.); Cardiology, Azienda Ospedaliero-Universitaria di Parma, University of Parma, Parma, Italy. (D.A.); ASTC: Associazione per lo Studio Della Trombosi in Cardiologia, Pavia, Italy. (D.A.); MRC/BHF Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK. (J.D.); The National Institute for Health Research Blood and Transplant Research Unit (NIHR BTRU) in Donor Health and Genomics at the University of Cambridge, Cambridge, UK. (J.D.); Wellcome Trust Sanger Institute, Genome Campus, Hinxton, UK. (J.D.); Department of Cardiology, Deutsches Herzzentrum München; Technische Universität München; Deutsches Zentrum für Herz-Kreislauf-Forschung (DZHK), München, Germany. (H.S.); Department of Medicine and Pediatrics, University of Mississippi Medical Center, Jackson, Mississippi, USA. (A.C.); Department of Cardiovascular Sciences, University of Leicester, Leicester, UK. (M.J.B., N.J.S.); NIHR Leicester Biomedical Research Centre, Glenfield Hospital, Leicester UK. (M.J.B., N.J.S.); Institute for Cardiogenetics, University of Lübeck, DZHK (German Research Centre for Cardiovascular Research), partner site Hamburg/Lübeck/Kiel, and University Heart Center Lübeck, Lübeck, Germany. (J.E.); University of Ottawa Heart Institute, Ottawa, Canada. (R.M.); Cardiovascular Medicine, Radcliffe Department of Medicine and the Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, UK. (H.W.); Department of Biostatistics and Epidemiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA. (D.S.); Cardiovascular Epidemiology and Genetics, Hospital del Mar Research Institute, Barcelona, Spain. (R.E.); CIBER Enfermedades Cardiovasculares (CIBERCV), Barcelona, Spain. (R.E.); Facultat de Medicina, Universitat de Vic-Central de Cataluña, Vic, Spain. (R.E.); Division of Cardiology, Department of Medicine, Duke University, Durham, North Carolina, USA. (S.H.S.); Departments of Genetics, University of

Pennsylvania, Philadelphia, Pennsylvania, USA. (D.J.R.); Verve Therapeutics, Cambridge, Massachusetts, USA. (S.K.)

## References

1. Bhattacharyya AK and Connor WE. Beta-sitosterolemia and xanthomatosis. A newly described lipid storage disease in two sisters. *J Clin Invest.* 1974;53:1033-43.
2. Salen G, Horak I, Rothkopf M, Cohen JL, Speck J, Tint GS, Shore V, Dayal B, Chen T and Shefer S. Lethal atherosclerosis associated with abnormal plasma and tissue sterol composition in sitosterolemia with xanthomatosis. *J Lipid Res.* 1985;26:1126-33.
3. Katayama T, Satoh T, Yagi T, Hirose N, Kurita Y, Anzai T, Asakura Y, Yoshikawa T, Mitamura H and Ogawa S. A 19-year-old man with myocardial infarction and sitosterolemia. *Intern Med.* 2003;42:591-4.
4. Tada H, Nohara A, Inazu A, Sakuma N, Mabuchi H and Kawashiri MA. Sitosterolemia, Hypercholesterolemia, and Coronary Artery Disease. *J Atheroscler Thromb.* 2018;25:783-789.
5. Altmann SW, Davis HR, Jr., Zhu LJ, Yao X, Hoos LM, Tetzloff G, Iyer SP, Maguire M, Golovko A, Zeng M, Wang L, Murgolo N and Graziano MP. Niemann-Pick C1 Like 1 protein is critical for intestinal cholesterol absorption. *Science.* 2004;303:1201-4.
6. Davis HR, Jr., Zhu LJ, Hoos LM, Tetzloff G, Maguire M, Liu J, Yao X, Iyer SP, Lam MH, Lund EG, Detmers PA, Graziano MP and Altmann SW. Niemann-Pick C1 Like 1 (NPC1L1) is the intestinal phytosterol and cholesterol transporter and a key modulator of whole-body cholesterol homeostasis. *J Biol Chem.* 2004;279:33586-92.
7. Berge KE, Tian H, Graf GA, Yu L, Grishin NV, Schultz J, Kwiterovich P, Shan B, Barnes R and Hobbs HH. Accumulation of dietary cholesterol in sitosterolemia caused by mutations in adjacent ABC transporters. *Science.* 2000;290:1771-5.
8. Lee MH, Lu K, Hazard S, Yu H, Shulnenin S, Hidaka H, Kojima H, Allikmets R, Sakuma N, Pegoraro R, Srivastava AK, Salen G, Dean M and Patel SB. Identification of a gene, ABCG5, important in the regulation of dietary cholesterol absorption. *Nat Genet.* 2001;27:79-83.
9. Graf GA, Yu L, Li WP, Gerard R, Tuma PL, Cohen JC and Hobbs HH. ABCG5 and ABCG8 are obligate heterodimers for protein trafficking and biliary cholesterol excretion. *J Biol Chem.* 2003;278:48275-82.
10. Temel RE, Tang W, Ma Y, Rudel LL, Willingham MC, Ioannou YA, Davies JP, Nilsson LM and Yu L. Hepatic Niemann-Pick C1-like 1 regulates biliary cholesterol concentration and is a target of ezetimibe. *J Clin Invest.* 2007;117:1968-78.
11. Klett EL, Lee MH, Adams DB, Chavin KD and Patel SB. Localization of ABCG5 and ABCG8 proteins in human liver, gall bladder and intestine. *BMC Gastroenterol.* 2004;4:21.

12. Kidambi S and Patel SB. Sitosterolaemia: pathophysiology, clinical presentation and laboratory diagnosis. *J Clin Pathol*. 2008;61:588-94.
13. Webb TR, Erdmann J, Stirrups KE, Stitzel NO, Masca NG, Jansen H, Kanoni S, Nelson CP, Ferrario PG, Konig IR, Eicher JD, Johnson AD, Hamby SE, Betsholtz C, Ruusalepp A, Franzen O, Schadt EE, Bjorkegren JL, Weeke PE, Auer PL, Schick UM, Lu Y, Zhang H, Dube MP, Goel A, Farrall M, Peloso GM, Won HH, Do R, van Iperen E, Kruppa J, Mahajan A, Scott RA, Willenborg C, Braund PS, van Capelleveen JC, Doney AS, Donnelly LA, Asselta R, Merlini PA, Duga S, Marziliano N, Denny JC, Shaffer C, El-Mokhtari NE, Franke A, Heilmann S, Hengstenberg C, Hoffmann P, Holmen OL, Hveem K, Jansson JH, Jockel KH, Kessler T, Kriebel J, Laugwitz KL, Marouli E, Martinelli N, McCarthy MI, Van Zuydam NR, Meisinger C, Esko T, Mihailov E, Escher SA, Alver M, Moebus S, Morris AD, Virtamo J, Nikpay M, Olivieri O, Provost S, AlQarawi A, Robertson NR, Akinsansya KO, Reilly DF, Vogt TF, Yin W, Asselbergs FW, Kooperberg C, Jackson RD, Stahl E, Muller-Nurasyid M, Strauch K, Varga TV, Waldenberger M, Wellcome Trust Case Control C, Zeng L, Chowdhury R, Salomaa V, Ford I, Jukema JW, Amouyel P, Kontto J, Investigators M, Nordestgaard BG, Ferrieres J, Saleheen D, Sattar N, Surendran P, Wagner A, Young R, Howson JM, Butterworth AS, Danesh J, Ardissono D, Bottinger EP, Erbel R, Franks PW, Girelli D, Hall AS, Hovingh GK, Kastrati A, Lieb W, Meitinger T, Kraus WE, Shah SH, McPherson R, Orho-Melander M, Melander O, Metspalu A, Palmer CN, Peters A, Rader DJ, Reilly MP, Loos RJ, Reiner AP, Roden DM, Tardif JC, Thompson JR, Wareham NJ, Watkins H, Willer CJ, Samani NJ, Schunkert H, Deloukas P, Kathiresan S, Myocardial Infarction G and Investigators CAEC. Systematic Evaluation of Pleiotropy Identifies 6 Further Loci Associated With Coronary Artery Disease. *J Am Coll Cardiol*. 2017;69:823-836.
14. Teupser D, Baber R, Ceglarek U, Scholz M, Illig T, Gieger C, Holdt LM, Leichtle A, Greiser KH, Huster D, Linsel-Nitschke P, Schafer A, Braund PS, Tiret L, Stark K, Raaz-Schrauder D, Fiedler GM, Wilfert W, Beutner F, Gielen S, Grosshennig A, Konig IR, Lichtner P, Heid IM, Klutigg A, El Mokhtari NE, Rubin D, Ekici AB, Reis A, Garlichs CD, Hall AS, Matthes G, Wittekind C, Hengstenberg C, Cambien F, Schreiber S, Werdan K, Meitinger T, Loeffler M, Samani NJ, Erdmann J, Wichmann HE, Schunkert H and Thiery J. Genetic regulation of serum phytosterol levels and risk of coronary artery disease. *Circ Cardiovasc Genet*. 2010;3:331-9.
15. Merkens LS, Myrie SB, Steiner RD and Mymin D. Sitosterolemia. *GeneReviews*®. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK131810/> (Updated 2018 May 17; last accessed 2019 Sep 18).
16. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, Grody WW, Hegde M, Lyon E, Spector E, Voelkerding K, Rehm HA-O and Committee ALQA. Standards and guidelines for the

interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17:405-24.

17. Atherosclerosis, Thrombosis, and Vascular Biology Italian Study Group. No evidence of association between prothrombotic gene polymorphisms and the development of acute myocardial infarction at a young age. *Circulation*. 2003;107:1117-22.
18. Myocardial Infarction Genetics Consortium Investigators. Inactivating mutations in NPC1L1 and protection from coronary heart disease. *N Engl J Med*. 2014;371:2072-82.
19. Do R, Stitzel NO, Won HH, Jorgensen AB, Duga S, Angelica Merlini P, Kiezun A, Farrall M, Goel A, Zuk O, Guella I, Asselta R, Lange LA, Peloso GM, Auer PL, Project NES, Girelli D, Martinelli N, Farlow DN, DePristo MA, Roberts R, Stewart AF, Saleheen D, Danesh J, Epstein SE, Sivapalaratnam S, Hovingh GK, Kastelein JJ, Samani NJ, Schunkert H, Erdmann J, Shah SH, Kraus WE, Davies R, Nikpay M, Johansen CT, Wang J, Hegele RA, Hechter E, Marz W, Kleber ME, Huang J, Johnson AD, Li M, Burke GL, Gross M, Liu Y, Assimes TL, Heiss G, Lange EM, Folsom AR, Taylor HA, Olivieri O, Hamsten A, Clarke R, Reilly DF, Yin W, Rivas MA, Donnelly P, Rossouw JE, Psaty BM, Herrington DM, Wilson JG, Rich SS, Bamshad MJ, Tracy RP, Cupples LA, Rader DJ, Reilly MP, Spertus JA, Cresci S, Hartiala J, Tang WH, Hazen SL, Allayee H, Reiner AP, Carlson CS, Kooperberg C, Jackson RD, Boerwinkle E, Lander ES, Schwartz SM, Siscovick DS, McPherson R, Tybjaerg-Hansen A, Abecasis GR, Watkins H, Nickerson DA, Ardissono D, Sunyaev SR, O'Donnell CJ, Altshuler D, Gabriel S and Kathiresan S. Exome sequencing identifies rare LDLR and APOA5 alleles conferring risk for myocardial infarction. *Nature*. 2015;518:102-6.
20. Taylor HA, Jr. The Jackson Heart Study: an overview. *Ethn Dis*. 2005;15:S6-1-3.
21. Dhillon OS, Khan SQ, Narayan HK, Ng KH, Struck J, Quinn PA, Morgenthaler NG, Squire IB, Davies JE, Bergmann A and Ng LL. Prognostic value of mid-regional pro-adrenomedullin levels taken on admission and discharge in non-ST-elevation myocardial infarction: the LAMP (Leicester Acute Myocardial Infarction Peptide) II study. *J Am Coll Cardiol*. 2010;56:125-33.
22. Myocardial Infarction Genetics and CARDIoGRAM Exome Consortia Investigators. Coding Variation in ANGPTL4, LPL, and SVEP1 and the Risk of Coronary Disease. *N Engl J Med*. 2016;374:1134-44.
23. McPherson R, Pertsemlidis A, Kavaslar N, Stewart A, Roberts R, Cox DR, Hinds DA, Pennacchio LA, Tybjaerg-Hansen A, Folsom AR, Boerwinkle E, Hobbs HH and Cohen JC. A common allele on chromosome 9 associated with coronary heart disease. *Science*. 2007;316:1488-91.

24. Clarke R, Peden JF, Hopewell JC, Kyriakou T, Goel A, Heath SC, Parish S, Barlera S, Franzosi MG, Rust S, Bennett D, Silveira A, Malarstig A, Green FR, Lathrop M, Gigante B, Leander K, de Faire U, Seedorf U, Hamsten A, Collins R, Watkins H, Farrall M and Consortium P. Genetic variants associated with Lp(a) lipoprotein level and coronary disease. *N Engl J Med.* 2009;361:2518-28.
25. Saleheen D, Zaidi M, Rasheed A, Ahmad U, Hakeem A, Murtaza M, Kayani W, Faruqui A, Kundi A, Zaman KS, Yaqoob Z, Cheema LA, Samad A, Rasheed SZ, Mallick NH, Azhar M, Jooma R, Gardezi AR, Memon N, Ghaffar A, Fazal ur R, Khan N, Shah N, Ali Shah A, Samuel M, Hanif F, Yameen M, Naz S, Sultana A, Nazir A, Raza S, Shazad M, Nasim S, Javed MA, Ali SS, Jafree M, Nisar MI, Daood MS, Hussain A, Sarwar N, Kamal A, Deloukas P, Ishaq M, Frossard P and Danesh J. The Pakistan Risk of Myocardial Infarction Study: a resource for the study of genetic, lifestyle and other determinants of myocardial infarction in South Asia. *Eur J Epidemiol.* 2009;24:329-38.
26. Senti M, Tomas M, Marrugat J, Elosua R and Investigators R. Paraoxonase1-192 polymorphism modulates the nonfatal myocardial infarction risk associated with decreased HDLs. *Arterioscler Thromb Vasc Biol.* 2001;21:415-20.
27. Davies RW, Wells GA, Stewart AF, Erdmann J, Shah SH, Ferguson JF, Hall AS, Anand SS, Burnett MS, Epstein SE, Dandona S, Chen L, Nahrstaedt J, Loley C, Konig IR, Kraus WE, Granger CB, Engert JC, Hengstenberg C, Wichmann HE, Schreiber S, Tang WH, Ellis SG, Rader DJ, Hazen SL, Reilly MP, Samani NJ, Schunkert H, Roberts R and McPherson R. A genome-wide association study for coronary artery disease identifies a novel susceptibility locus in the major histocompatibility complex. *Circ Cardiovasc Genet.* 2012;5:217-25.
28. Reilly MP, Li M, He J, Ferguson JF, Stylianou IM, Mehta NN, Burnett MS, Devaney JM, Knouff CW, Thompson JR, Horne BD, Stewart AF, Assimes TL, Wild PS, Allayee H, Nitschke PL, Patel RS, Myocardial Infarction Genetics C, Wellcome Trust Case Control C, Martinelli N, Girelli D, Quyyumi AA, Anderson JL, Erdmann J, Hall AS, Schunkert H, Quertermous T, Blankenberg S, Hazen SL, Roberts R, Kathiresan S, Samani NJ, Epstein SE and Rader DJ. Identification of ADAMTS7 as a novel locus for coronary atherosclerosis and association of ABO with myocardial infarction in the presence of coronary atherosclerosis: two genome-wide association studies. *Lancet.* 2011;377:383-92.
29. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, del Angel G, Rivas MA, Hanna M, McKenna A, Fennell TJ, Kurnytsky AM, Sivachenko AY, Cibulskis K, Gabriel SB, Altshuler D and Daly MJ. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet.* 2011;43:491-8.

30. Salen G, von Bergmann K, Lutjohann D, Kwiterovich P, Kane J, Patel SB, Musliner T, Stein P, Musser B and Multicenter Sitosterolemia Study G. Ezetimibe effectively reduces plasma plant sterols in patients with sitosterolemia. *Circulation*. 2004;109:966-71.
31. Genser B, Silbernagel G, De Backer G, Bruckert E, Carmena R, Chapman MJ, Deanfield J, Descamps OS, Rietzschel ER, Dias KC and Marz W. Plant sterols and cardiovascular disease: a systematic review and meta-analysis. *Eur Heart J*. 2012;33:444-51.

**Table 1. Clinical characteristics by *ABCG5* and *ABCG8* variant carrier status in sitosterolemia families.**

	Non-carrier	<i>ABCG5</i>		<i>ABCG8</i>	
		Hetero	Homo	Hetero	Homo
N	11	22	8	5	1
Age, mean ± SD	40.5 ± 19	41.1 ± 21	12.9 ± 20	45.2 ± 19	1
Male sex, n (%)	3 (27)	12 (55)	3 (38)	2 (40)	0
Lipid profile					
Total cholesterol, mg/dL, mean ± SD	180 ± 28	236 ± 70*	512 ± 250*	258 ± 67	968
LDL cholesterol, mg/dL, mean ± SD	105 ± 30	158 ± 59*	412 ± 240*	150 ± 52	832
HDL cholesterol, mg/dL, mean ± SD	61.1 ± 15	56.1 ± 19	47.4 ± 13	57.0 ± 13	46
Triglyceride, mg/dL, median (IQR)	87 (74–96)	76 (55–150)	188 (140–248)*	154 (73–154)	71
Lipoproteins					
Apolipoprotein A1, mg/dL, median (IQR)	150 (137–165)	140 (128–150)	106 (97–129)	NA	NA
Apolipoprotein B, mg/dL, median (IQR)	67 (63–93)	106 (91–117)	262 (198–303)*	NA	NA
Non-cholesterol sterols					
Sitosterol, µg/mL, median (IQR)	2.2 (1.8–2.7)	7.6 (5.5–10)*	102 (74–125)*	9.9 (8.2–12)*	36.5
Campesterol, µg/mL, median (IQR)	3.6 (3.2–4.5)	13 (10–14)*	70 (65–95)*	NA	NA
Sitosterol / TC, µg/mg, median (IQR)	1.2 (1.1–1.4)	3.5 (2.5–4.6)*	22 (14–30)*	4.4 (3.9–5.3)*	3.8
Campesterol / TC, µg/mg, median (IQR)	2.2 (1.7–2.6)	5.2 (4.6–6.5)*	13 (7.5–18)*	NA	NA

\*: P value < 0.025 compared to non-carrier controls. P values were calculated by *Mann-Whitney U* test. Abbreviations: IQR, interquartile range; SD, standard deviation; TC, total cholesterol.

**Table 2. Clinical characteristics of participants in MiGen, UK Biobank and targeted resequencing study.**

<b>Myocardial Infarction Genetics Consortium</b>	<b>UK Biobank</b>	<b>Target Resequencing Study</b>	
<b>N = 58,791</b>	<b>N = 336,391</b>	<b>N = 1,714</b>	
Age, years (SD)	54 (10)	57 (8)	57.2 (13)
Male gender, n (%)	41,203 (71)	156,112 (46)	1,140 (67)
BMI (SD), kg/m <sup>2</sup>	27 (5)	27 (5)	29 (6)
Current smoker, n (%)	18,173 (33)	25,802 (8)	639 (37)
<b>Medical history</b>			
Coronary artery disease, n (%)	16,106 (33)	12,073 (3)	1,184 (69)
Hypertension, n (%)	16,839 (35)	153,535 (46)	798 (47)
Type 2 diabetes, n (%)	11,245 (22)	15,770 (5)	277 (16)
<b>Lipid profile</b>			
Total cholesterol, mg/dl (SD)	178 (60)	222 (41)	182 (51)
LDL cholesterol, mg/dl <sup>†</sup> (SD)	109 (45)	138 (34)	108 (41)
HDL cholesterol, mg/dl (SD)	36 (14)	56 (15)	43 (7)
Triglycerides, mg/dl (SD)	154 (127)	155 (90)	135 (82)

Abbreviations: SD, standard deviation.