

Structural basis of cowpox evasion of NKG2D immunosurveillance

Eric Lazear^{1,9,12}, Michel M. Sun^{3,11,12}, Xiaoli Wang^{1,2}, Theresa L. Geurs³, Christopher A. Nelson¹, Jessica A. Campbell^{4,8}, Danna Lippold³, Alexander S. Krupnick^{5,10}, Randall S. Davis⁶, Leonidas N. Carayannopoulos^{4,7}, Anthony R. French^{1,3*} and Daved H. Fremont^{1,2*}

Departments of Pathology & Immunology¹, Biochemistry & Molecular Biophysics², Pediatrics³, Internal Medicine⁴, and Surgery⁵ Washington University in St. Louis, St. Louis, Missouri, USA; Departments of Medicine, Microbiology, and Biochemistry and Molecular Genetics⁶, University of Alabama at Birmingham, Birmingham, AL 35294

⁷Current address: Celgene Pharmaceutical Company, Summit, NJ.

⁸Current address: Pfizer Inc., St. Louis, MO.

⁹Current address: Courier Therapeutics, Houston, TX.

¹⁰Current address: Department of Surgery, University of Virginia, Charlottesville, VA.

¹¹Current address: Department of Ophthalmology, University of California, Los Angeles, CA.

¹²These authors contributed equally to this work

* Co-corresponding authors:

fremont@wustl.edu, phone: 314-255-6171, fax: 314-362-8888

french_a@kids.wustl.edu, phone: 314-286-2885; fax: 314-286-2895

Running title: Structural basis of NKG2D viral subversion

Abstract

NKG2D is a key component of cytotoxic antitumor and antiviral responses. Multiple viruses evade NKG2D recognition by blocking NKG2D ligand expression on infected cells. In contrast, cowpox virus targets NKG2D directly by encoding a secreted antagonist, Orthopoxvirus MHC Class I-like Protein (OMCP). We have previously reported that OMCP also binds to the orphan receptor FcRL5 on innate B cells. Here, we demonstrate that mammalian-derived, glycosylated OMCP binds NKG2D but not FcRL5. Cowpox viruses either lacking OMCP, or expressing an NKG2D-binding deficient mutant, are significantly attenuated in wild type and FcRL5-deficient mice but not NKG2D-deficient mice, demonstrating that OMCP is critical in subverting NKG2D-mediated immunity *in vivo*. Next we determined the structure of OMCP bound to human NKG2D. Despite a structure similar to that of host NKG2D ligands, OMCP uses a drastically different orientation for NKG2D binding. The re-orientation of OMCP is associated with dramatically higher affinity for human NKG2D and the targeted interface is highly conserved in mammalian NKG2Ds, increasing the zoonotic potential of cowpox virus. We also show that cell surface presented OMCP can trigger NKG2D effector functions equivalently to host NKG2D ligands, demonstrating that NKG2D-mediated signaling requires clustering but is insensitive to binding orientation. Thus, in contrast to TCR/MHC interactions, the docking topology of NKG2D with its ligands does not appear to regulate its activation.

Author Summary

Virally infected or tumor-transformed cells display NKG2D ligands (NKG2DLs) on their cell surface, which activates NKG2D-bearing lymphocytes to kill the transformed cell. Pathogens are known to counter this by blocking NKG2DL expression and/or surface display. In contrast, some tumor cells cleave endogenous NKG2DLs creating soluble NKG2D antagonists. Unlike other viral pathogens, cowpox virus uses a strategy analogous to cancer cells by targeting NKG2D directly with a soluble, high affinity NKG2D-antagonist named OMCP. We determined that OMCP's virulence *in vivo* is attributed to blocking NKG2D-mediated NK cell responses with no apparent effect due to binding to other receptors or cell types. We have also determined the crystal structure of cowpox OMCP bound to human NKG2D, revealing that despite conservation of the ligand scaffolding with host NKG2DLs, the viral protein is engaged with a radically altered orientation compared to all host NKG2DLs. Our structure provides key insight into how OMCP binds with an ~5,000-fold increased affinity compared to human NKG2DLs and show that the OMCP binding site is exceptionally conserved among primates and rodents, suggesting that the ability of OMCP to recognize this conserved interface contributes to the broad zoonotic potential of cowpox virus. Finally, we show that cell membrane-anchored OMCP can trigger equivalent NKG2D-mediated killing as host NKG2DLs, demonstrating that NKG2D signaling is insensitive to ligand binding orientation.

Introduction

Intracellular surveillance mediated by MHC class I (MHCI) is a critical host immune function and as such MHCI molecules are frequently targeted for destruction or intracellular retention by viruses [1]. Many herpesviruses encode at least one protein that prevents the cell surface expression of MHCI [1, 2]. However, this immune evasion strategy renders the infected cell susceptible to NK cell-mediated lysis due to loss of inhibitory signals [3]. Viral infection also leads to cell surface display of NKG2D ligands (NKG2DLs) recognized by the activating receptor NKG2D, further predisposing the infected cell towards NK cell-mediated lysis. Therefore, viruses that target MHCI expression often also sabotage NKG2D-mediated cell responses by targeting NKG2DLs on infected cells [4-7].

NKG2DLs are not normally expressed on the cell surface but can be induced by cellular stress [8]. The specific trigger for NKG2DL expression is not known, but NKG2DLs are upregulated in response to several viral infections [9-12]. NKG2DLs comprise a large group of proteins all recognized by NKG2D, despite having low sequence identity. NKG2DLs include the MIC (A and B) and ULBP (1-6) families in humans as well as MULT1 and the RAE-1 (α - ϵ) and H60 (α -c) families in mice [13]. The redundancy in NKG2DLs is likely due to a combination of tissue specific expression patterns of the ligands and the need to counter viral NKG2D evasion strategies [14]. Many viruses have evolved mechanisms to inhibit the cell surface expression of NKG2DLs as a means of interfering with NKG2D surveillance of viral infection. This strategy is most apparent among β - and γ -herpesviruses, in which four murine cytomegalovirus proteins (m138, m145, m152, m155) [15-18], two human cytomegalovirus proteins (UL16, UL142) [19, 20] and one Kaposi's sarcoma-associated herpesvirus protein (K5) [21] have been demonstrated to block NKG2DL surface expression. This evasion strategy is also found in RNA viruses, as hepatitis C virus NS3/4a and human immunodeficiency virus Nef proteins also block the expression of a subset of NKG2DLs [22, 23]. Additionally, human cytomegalovirus, herpes

simplex virus type 1 and Epstein-Barr virus each also encode at least one miRNA that prevents translation of MICB [24, 25]. Similarly, JCV and BKV polyoma viruses target ULBP3 with miRNAs [26]. However, blocking NKG2DL expression on the infected cell is an imperfect evasion strategy, since no single viral protein or miRNA has been shown to block the expression of all NKG2DLs.

Like several herpesviruses, cowpoxvirus (CPXV) also sabotages MHCI expression. CPXV expresses CPXV012 and CPXV203, two proteins that prevent TAP-mediated peptide transport and MHCI trafficking to the cell surface, respectively [27-34]. Ectromelia virus, a related orthopoxvirus, induces NKG2DL expression, and NKG2D is critical for the control of ectromelia virus pathogenesis [35]. Infection with another orthopoxvirus, monkeypox virus, leads to dramatic expansion of NK cells but impaired NK cell function [36]. Together this suggests that CPXV infected cells would be sensitive to NK cell-mediated lysis.

Unlike herpesviruses, CPXV does not target NKG2DLs. Instead this virus targets NKG2D directly by encoding a competitive inhibitor of NKG2DLs, orthopoxvirus MHC class I-like protein (OMCP) [37, 38]. OMCP is a 152 residue protein that is secreted from infected cells and antagonizes the NKG2D-mediated killing of NKG2DL-expressing target cells [37]. OMCP also was reported to bind to B cells via Fc-receptor like protein 5 (FcRL5) and to monocyte/macrophages [39]. OMCP binds to murine NKG2D with an affinity equal or greater than all tested murine NKG2DLs, and to human NKG2D with an affinity ~5,000-fold higher than human NKG2DLs [37, 38, 40, 41].

Despite their divergence in sequence identity, all known host NKG2DLs share common structural features [42, 43]. NKG2DLs contain an MHCI-like platform domain composed of an eight-stranded beta sheet with two helices [44-48]. The platform domain is subdivided into $\alpha 1$ and $\alpha 2$ domains, with each domain containing four beta strands and an alpha helix. Unlike MHCI, the groove between the helices of the NKG2DL platform domain is closed and therefore NKG2DLs do not bind peptides.

Like host NKG2DLs, OMCP also adopts an MHCI-like platform domain [38]. However, the platform domain of OMCP has been trimmed to have only a six-stranded beta sheet with shorter flanking helices. In this paper, we term the helix of the $\alpha 1$ domain H1 and the discontinuous helix of the $\alpha 2$ domain is termed H2a and H2b. The H2a and H2b helices of OMCP are also rearranged to be flatter against the beta sheet and to be splayed apart from each other. These differences in the OMCP structure were hypothesized to be important for the high affinity binding of OMCP to NKG2D. However, OMCP was still expected to bind to NKG2D in the same orientation as host NKG2DLs, i.e. with the alpha helices oriented diagonally within the symmetric NKG2D binding groove.

Here we report the 2.0 Å-resolution structure of human NKG2D bound to OMCP of the Brighton Red strain of cowpoxvirus. The structure reveals a significant reorientation of OMCP in the NKG2D binding groove relative to host NKG2DLs. The interface of OMCP with NKG2D is highly complementary, buries a significantly larger surface area than host NKG2DLs, and remains continuous across the entire NKG2D binding groove. This novel binding adaptation and high affinity allows OMCP to compete with the high local concentration of membrane-associated host NKG2DLs. We further show that the mechanism of NKG2D antagonism requires OMCP to be secreted, lest it lead to NKG2D signaling. Finally, we show that viruses expressing an NKG2D-binding deficient OMCP mutant are equivalently attenuated to viruses that do not express OMCP using mouse models of cowpox infection. Thus demonstrating that while FcRL5 plays no role in cowpox pathogenesis, OMCP is critical in attenuating NKG2D-mediated immunity.

Results

CPXV lacking OMCP is significantly attenuated *in vivo*.

While OMCP has been shown to bind to NKG2D and compete with NKG2D ligands *in vitro*, the effect of OMCP on CPXV virulence *in vivo* has not been determined. We generated an OMCP-

deficient ($\Delta V018$) CPXV using a transient dominant selection strategy. Multi-step growth curves of the WT CPXV, OMCP-deficient CPXV, and a revertant CPXV demonstrated nearly identical replication growth kinetics (Fig. S1), indicating that OMCP is dispensable for viral growth or replication *in vitro*.

Next we investigated the impact of OMCP during systemic CPXV infection (via the intraperitoneal (i.p.) route) in WT B6 mice. Survival of WT mice infected with WT CPXV demonstrated dose dependence with an LD₅₀ value of 0.91×10^6 pfu/mouse (Fig. 1A). Comparison of the survival of WT mice infected i.p. with OMCP-deficient CPXV (Fig. 1B) revealed substantially greater survival in the mice infected with OMCP-deficient CPXV than with WT CPX. Indeed, the LD₅₀ value of 1.64×10^6 pfu/mouse for mice infected with OMCP-deficient CPXV demonstrated that CPXV lacking OMCP is attenuated *in vivo* compared to WT CPXV. Kaplan Meier survival curves of WT mice infected with the OMCP revertant CPXV correlate well with the survival observed following infection with WT CPXV (Fig. 1C), demonstrating that the decreased virulence observed during infection with OMCP-deficient CPXV reflects the absence of OMCP. Together, these results establish for the first time that OMCP facilitates CPXV virulence *in vivo*.

Structure determination of OMCP-NKG2D. We had previously solved the structure of OMCP alone and shown that, similar to host NKG2DLs, OMCP adopts an MHCI-like platform domain [38]. Despite the overall similarity of the domain structure of OMCP to host NKG2DLs, OMCP had several notable deviations in the putative NKG2D-binding site that were hypothesized to be important for the high affinity binding of OMCP to NKG2D. To further understand the unusually high affinity of OMCP for NKG2D, we crystallized and solved the structure of OMCP bound to human NKG2D.

Initial crystallization trials with OMCP and NKG2D yielded ~30 different crystallization conditions. Subsequent data collection and molecular replacement of multiple low-resolution

crystal forms all yielded similar partial solutions, with alternating sheets of OMCP-NKG2D complexes separated by undefined density. In the original structure of OMCP alone, the beta sheets packed to form a trimer with the alpha helices oriented away from the center [38]. An identical OMCP trimer formed in the OMCP-NKG2D partial solutions, with NKG2D now bound to the outward facing helices (data not shown). In an attempt to change the lattice packing, we introduced mutations into the beta sheet of OMCP that were designed to break the trimeric interface. These mutations were on the opposite face of OMCP from the NKG2D binding site to avoid disrupting OMCP-NKG2D binding. A mutant form of OMCP (Y23D, F95D) crystallized with NKG2D in a new space group and the crystals diffracted to 2.0 Å (Table 1)(Figure 2A).

The electron density map was continuous and unambiguous throughout all chains of the structure, with the exception of Q108 in OMCP. This residue was situated in the center of the largest loop of OMCP and unambiguous density for this residue was also absent from the structure of OMCP alone [38]. The structure of OMCP bound to NKG2D showed no major differences from our previous structure of OMCP alone, with an RMSD for all atoms of 0.8 Å. Likewise, NKG2D was also similar to previous NKG2D structures with RMSDs ranging from 0.5-0.9 Å. The β3-β4 loop of NKG2D is the only region of either OMCP or NKG2D that displayed above-average B factors. This loop is thought to be flexible and has had above average B factors in all previous NKG2D structures [49]. Interestingly, the peptide bond between S193-S194 in our NKG2D structure had a *cis* conformation not described in other NKG2D structures (Figure S2).

The interface between OMCP and NKG2D. OMCP was hypothesized to bind to the same surface of NKG2D used by host NKG2DLs because (i) OMCP competed with host NKG2DLs for NKG2D and (ii) mutations within the NKG2DL-binding pocket of NKG2D altered OMCP binding affinity [38]. OMCP does bind NKG2D using the same concave binding pocket as host NKG2DLs (Figure 2A). OMCP binds primarily using the discontinuous helices of its α2 domain,

H2a and H2b. The position of the H2a and H2b helices is such that every surface exposed side chain of both helices within the binding site directly contacts NKG2D (Figure 2B). Only two contacts are found outside of H2a and H2b, Ile49 and Arg66. Both of these residues are within the $\alpha 1$ domain but lie outside of the H1 helix.

Twelve OMCP residues contact eighteen NKG2D residues to form a mixture of bond types (Table 2). Three residues in each NKG2D half-site are known as core binding residues because they make contacts with all known host NKG2DLs. The core residues of NKG2D subunit A (NKG2D^A) (Tyr152, Tyr199, Met184) form two hydrogen bonds and make extensive hydrophobic contacts with OMCP residues. The core residues of NKG2D^A contact four OMCP residues and the most critical of these residues is Phe122. Phe122 makes multiple hydrophobic contacts with all three NKG2D^A core residues, including *pi*-stacking with Tyr152. Phe122 also forms a backbone-to-sidechain hydrogen bond with Tyr152. Interestingly, OMCP is the first NKG2D ligand not to utilize all six NKG2D core-binding residues, with only Met184 and Tyr152 of NKG2D subunit B (NKG2D^B) contacting OMCP. NKG2D^B Met184 and Tyr152 each make a single hydrogen bond and hydrophobic contacts with OMCP residues. Two OMCP residues, Trp127 and Asp132, make contacts with both NKG2D protomers. OMCP Trp127 forms a hydrogen bond to Lys150 of NKG2D^A and makes several hydrophobic contacts with Leu148 of NKG2D^B, Lys150 and Ser151 of NKG2D^A. OMCP Asp132 forms a hydrogen bond with Tyr152 of NKG2D^B and a salt bridge with Lys150 of NKG2D^A (Figure 3A).

Due to the high affinity of the OMCP-NKG2D interaction we harnessed a high throughput *in vitro* selection approach to find NKG2D-binding null mutants (Table S1). The results of the screen identified D132 as an important residue for disrupting NKG2D binding. We then generated OMCP with the D132R mutation in an attempt to completely ablate NKG2D binding. The D132R mutant protein was unable to bind to NKG2D-expressing cells, but retained binding to FcRL5-expressing cells (Figure 3B). This mutation is likely to cause significant steric clashes,

as well as disrupting both interactions made by Asp132 to NKG2D^A Lys150 and NKG2D^B Tyr152 (Figure 3A).

The role of OMCP receptors in CPXV infection. We generated a recombinant CPXV that expressed the D132R mutant OMCP to isolate the effect of OMCP binding to NKG2D from binding to FcRL5. A multi-step growth curve of (D132R) OMCP CPXV demonstrated identical replication growth kinetics to WT CPXV (Fig. S1). Cowpox and its close relatives variola virus, vaccinia virus, and monekeypox can be spread via contact and aerosol transmission [50]; therefore, we choose to study the role of OMCP during acute intranasal (i.n.) infection. Using our series of WT OMCP expressing CPXV, OMCP-deficient CPXV (Δ V018), and (D132R) OMCP mutant expressing CPXV in combination with WT, NKG2D-deficient, and FcRL5-deficient mice, we were able to delineate the impact of OMCP on NKG2D- and FcRL5-expressing cells during CPXV infection *in vivo*.

Mice were infected i.n. with 2.5×10^4 pfu/mouse of WT, OMCP-deficient, or D132R CPXV virus. Infection with WT CPXV resulted in similar mortality in WT, NKG2D-deficient, and FcRL5-deficient mice (Fig. 4 solid lines). However, the complete loss of OMCP (Fig. 4, dashed lines) rendered CPXV severely attenuated in WT and FcRL5-deficient mice, while OMCP-deficient CPXV infection in NKG2D-deficient mice remained similarly lethal as WT CPXV. These results illustrated that under normal circumstances (*i.e.*, WT CPXV infection in WT mice), OMCP is able to effectively abrogate activation through the NKG2D receptor, rendering WT CPXV as lethal in WT mice as it is in NKG2D-deficient mice. Mice infected with the (D132R) OMCP CPXV (Fig. 4, dotted lines) phenocopied the mortality of mice infected with OMCP-deficient CPXV. The survival observed in the FcRL5-deficient mice following infection with each of the strains of CPXV closely mimicked that seen in WT mice (Figs. 4C vs 4A), indicating that there is no significant role of the FcRL5 receptor on survival following this route of infection. The significant attenuation of OMCP-deficient CPXV and the (D132R) OMCP CPXV infection in WT and

FcRL5-deficient mice demonstrated the critical importance of NKG2D blockade during WT CPXV infection, and strongly implicates OMCP blockade of NKG2D as the primary virulence effect.

Glycosylated OMCP does not bind FcRL5. Compared to previous *in vitro* experiments with bacterial-derived OMCP [37-39], OMCP produced in mammalian cells would be modified by N-linked glycosylation. OMCP is glycosylated at N12, one of its two putative N-glycan sites, when produced in mammalian cells (Fig S3A). The absence of a phenotype for FcRL5 in mice led us to confirm whether mammalian-derived, glycosylated OMCP binds to FcRL5-expressing cells. To test whether glycosylation would affect OMCP receptor binding, we compared binding of bacterially-expressed or mammalian-expressed OMCP to Ba/f3 cells expressing NKG2D or FcRL5 (Fig. 5A). Regardless of source, OMCP bound to NKG2D-expressing cells. In contrast to bacterially-expressed OMCP, mammalian-expressed OMCP did not bind to FcRL5-expressing cells. To confirm whether the difference in FcRL5 binding was due to glycosylation, mammalian-expressed OMCP was treated with EndoF to remove the N-linked glycan. The deglycosylated protein gained the ability to interact with FcRL5. Next we tested the binding of mammalian-expressed OMCP to primary human PBMCs (Fig 5B). As a positive control for NKG2D binding, we included the human NKG2DL ULBP3. WT OMCP and ULBP3 stained both NK and a subset of T cells, but not B cells, consistent with NKG2D expression on these cells. In contrast, (D132R) OMCP failed to significantly stain NK and T cells, supporting that WT OMCP cell staining is due to binding NKG2D. Together the *in vivo* virulence (Fig 4) and cell binding experiments (Fig 5) demonstrate that mammalian-derived OMCP functions only as a high affinity NKG2D antagonist and is not a bona fide FcRL5 ligand. Importantly, OMCP blockade of NKG2D plays a significant role in pathogenesis in mice, and potentially an even larger role with higher affinity NKG2D orthologs, like human NKG2D.

Structural basis for species-specific affinity. Previously, the 14-fold higher affinity of OMCP for human vs murine NKG2D was mapped to three amino acid substitutions in the $\beta 5'$ - $\beta 5$ loop of NKG2D, abbreviated L2 [38]. In addition to the substitutions themselves (I182V, M184I and Q185P), the position of the loop between NKG2D orthologs differs. L2 in human NKG2D is bent towards the center of the concave binding cavity compared to L2 of murine NKG2D. Superimposition of murine NKG2D onto the human NKG2D-OMCP structure reveals that the contacts between OMCP and Met184 (mNKG2D residue I200) in NKG2D^B and between Met184 (I200) and Glu185 (P201) in NKG2D^A would be altered due to the different position of the murine $\beta 5'$ - $\beta 5$ loop (Figure 6A-B). This alteration would disrupt contacts with three residues in OMCP H2a, three residues in H2b and Arg66 within the $\alpha 1$ domain. Of the contact residues of L2, Met184 makes the most significant contacts in both NKG2Ds (Table 2)(Figure 6C). Critically, of the 58 NKG2D sequences available in GenBank, 54 conserve the Met184 and Glu185 found in the high affinity human NKG2D (Figure 6D).

Eighteen OMCP variants have been described between different CPXV and MPXV strains [51]. In this study, we have crystallized OMCP from the Brighton Red strain of CPXV which has >60% sequence identity with the highly conserved sequence of the other 17 OMCP variants, collectively termed OMCP_{mpx}. Of the 12 OMCP contact residues observed, 9 are identical to OMCP_{mpx}. Of the remaining contacts, all three are conservative hydrophobic substitutions (I49L, T118I and M135I) (Figure 7). OMCP_{mpx} binds to NKG2D and the substitutions in the NKG2D contact residues are unlikely to grossly affect the affinity of OMCP_{mpx} for NKG2D [37].

A novel NKG2D-binding adaptation. Host NKG2DLs have low sequence identity but overall similar structures, with MHCI-like platform domains binding diagonally across the symmetric binding groove created by the NKG2D homodimer [13, 42, 52]. Host ligands contact one NKG2D half site with H1 and the S1-S2 loop, and contact the second NKG2D half site with H2b.

Despite the similar MHCI-like fold, OMCP binds the NKG2D binding groove in a novel orientation, rotating $\sim 45^\circ$ relative to host NKG2DLs (Figure 7). Instead of using H1 and S1-S2 loop like host ligands, OMCP has replaced these contacts with H2a. This rotation leads to the helices of OMCP being perpendicular to the NKG2D binding groove, instead of lying diagonally across it.

Two unique rearrangements of H2a and H2b make the OMCP orientation possible. The $\alpha 2$ helices of OMCP and host NKG2DLs are discontinuous, with the two shorter helices hinged relative to each other. For host ligands, the angle between H2a and H2b is $\sim 90^\circ$, positioning H2a away from the NKG2D interface. In contrast, OMCP has increased the hinge angle between the helices by $\sim 20^\circ$, leading to a $\alpha 2$ helix that is flatter relative to the beta sheet of OMCP. The flattening of the $\alpha 2$ helix allows H2a and H2b to closely complement the concave binding groove of the NKG2D homodimer (Figure 2B). The tight fit of the $\alpha 2$ helix for NKG2D is reflected in the high shape complementarity (0.77) and buried surface area ($2,612 \text{ \AA}^2$). In contrast, host NKG2DLs have shape complementarity ranging from 0.63-0.72 and buried surface areas ranging from $1,700$ - $2,180 \text{ \AA}^2$ [44, 45, 47].

The second unique feature of the $\alpha 2$ helix is the separation of H2a and H2b relative to each other. This region also contains a translation that completely separates H2a and H2b into two distinct helices. This translation is critical for NKG2D binding because it allows each helix to be directly centered on the core binding sites of each NKG2D monomer (Figure 7). This creates a symmetric binding site on OMCP that recognizes the symmetric binding groove created by the NKG2D dimer. The symmetry between OMCP and NKG2D binding is in stark contrast to the canonical binding of an asymmetric host ligand to the symmetric NKG2D binding groove [52]. However, one element of asymmetry remains in the OMCP-NKG2D interaction because each NKG2D half-site recognizes an OMCP helix in a different N- to C-terminal orientation, demonstrating again the flexibility of NKG2Ds rigid adaptation recognition [42, 53].

The contact sites between NKG2D and host NKG2DLs are made up of two patches centered on the core binding sites of NKG2D and H1/S1-S2 loop and H2b of NKG2DLs [42]. As a result, the interface of NKG2D with NKG2DLs is discontinuous, particularly in the center of the NKG2D binding groove (Figure 7). Due to the unique orientation of OMCP, H2a and H2b make continuous contacts along the entire NKG2D binding groove (Figure 7). The sidechains of OMCP Lys126, Trp127, Glu131 and Asp132 make contacts with residues in the center of the NKG2D binding groove and bridge the core binding sites on each NKG2D monomer (Figure 2B). In particular, OMCP Trp127 is directed towards the center of the NKG2D dimer and makes hydrophobic contacts with residues on both NKG2D monomers, effectively closing any gaps in the binding interface.

Signaling of NKG2D upon ligand engagement. CPXV and MPXV-infected cells secrete OMCP, which can act as an NKG2D-antagonist and block *in vitro* NKG2D-mediated NK cell killing of target cells [37]. This immune evasion strategy is reminiscent of cancer induced-NKG2DL shedding. Some cancer cells proteolytically cleave NKG2DLs from the cell surface using matrix metalloproteinases (MMPs), simultaneously preventing NKG2D-bearing lymphocytes from targeting the cancer cell, as well as creating soluble NKG2DLs to inhibit NKG2D in *trans*. Cell-associated NKG2DLs trigger NKG2D effector functions (Figure 8A), while cancer-induced, soluble NKG2DLs block NKG2D function (Figure 8B). Like shed NKG2DLs, OMCP is soluble and blocks NKG2D function in *trans* [37] (Figure 8C). Unlike host NKG2DLs, OMCP binds NKG2D with a novel orientation. We therefore asked whether OMCP could serve as a NKG2D agonist in the context of the cell membrane, analogously to host NKG2D ligands. Since OMCP is a secreted protein, an artificially cell-associated OMCP was constructed by using a heterologous GPI anchor from Thy1.1 [37] (Figure 5D). To measure NKG2D-mediated cell killing, we stably transduced Ba/F3 cells with retroviral vectors expressing either the OMCP-Thy1.1 construct or host NKG2DLs. OMCP-Thy1.1-expressing target cells were killed

equivalently to host NKG2DL-transduced target cells, indicating that despite its altered binding orientation, cell-associated OMCP was able to activate NKG2D signaling (Figure 8E). Thus, OMCP must be secreted lest it active NKG2D-effector functions itself, despite potential loss of efficacy due to diffusion.

Discussion

Pathogens deploy multiple mechanisms to subvert the host immune response. We have previously reported that the poxvirus protein OMCP binds to two different murine immune receptors, NKG2D and FcRL5. Here, we demonstrate for the first time that OMCP facilitates cowpox virus virulence *in vivo* thorough antagonism of NKG2D. Surprisingly, we found no impact of FCRL5 *in vivo* during infections with OMCP-deficient CPXV or D132R CPVX (expressing OMCP with a mutation selectively abrogating binding with NKG2D). Investigations into the absence of a phenotype in the FCRL5-deficient mice revealed that mammalian-derived, glycosylated OMCP did not bind to FCRL5 in contrast to bacterially-expressed OMCP. Treatment of the mammalian-derived OMCP with EndoF removed the N-linked glycan and exposed a portion of OMCP that would not be present during *in vivo* infections. Surprisingly, this normally hidden surface of OMCP was able to interact with FcRL5-expressing cells. This observation has implications more broadly for the use of bacterially-expressed proteins in receptor-ligand studies. However, these data highlight the importance of OMCP in subverting NKG2D-mediated control of cowpox virus infection. In the absence of OMCP antagonism of NKG2D, cowpox virus was significantly attenuated. Cowpox virus inhibits the cell surface expression of MHCI to avoid recognition by effector T cells[30, 31]. The loss of MHCI expression on the cell surface induces “missing self” and enhanced recognition by NK cells. In the absence of OMCP antagonism of the activating NK cell receptor NKG2D, infected cells become more susceptible to NK cell-mediated killing.

While many viruses have adopted a general mechanism of NKG2D-sabotage by trying to retain multiple host-encoded NKG2D ligands within the infected cell, CPXV and MPXV take the very different approach of targeting NKG2D directly. Since NKG2D is monomorphic, this mechanism has the significant advantage of requiring a single protein to prevent NKG2D recognition of the infected cell. The large number of sequence-divergent host NKG2DLs and their associated polymorphisms are thought to be driven by selection from pathogen-encoded NKG2DL antagonists [14]. Likewise, viral NKG2L antagonists are under selective pressure from the diverse host NKG2DLs in a continual cycle of adaptation. Due to the need to recognize multiple NKG2DLs, NKG2D has a limited mutational space to adapt. The limited ability of NKG2D to mutate is yet another advantage of OMCP directly targeting NKG2D, instead of NKG2DLs.

Similarly to OMCP, some cancer cells shed host NKG2DLs to create their own soluble NKG2D antagonists. However, this strategy has the additional benefit of removing host NKG2DL from the surface of cancer cells. In contrast, CPXV and MPXV lack a known mechanism of blocking host NKG2DL surface expression. Secreted OMCP must then be able to compete efficiently against the high local concentration of multiple host NKG2DLs on the infected cell, as well as against diffusion away from the infected cell. One possible way to increase OMCP's ability to compete with host ligands would be to increase the avidity of OMCP by having multiple NKG2D-binding domains. However, a multimeric OMCP could crosslink NKG2D and potentially trigger NKG2D-mediated killing. Therefore, secreted OMCP must be monomeric to prevent aberrant NKG2D signaling. Thus to compensate for these deficiencies, OMCP must have the highest possible affinity to effectively compete against cell-associated host NKG2DLs [37, 38]. The half-life of ligand-receptor interactions correlate well with physiological competitiveness [54]. OMCP binds human and murine NKG2D with half-lives of 348 and 54 seconds, respectively, compared to half-lives of 1.5-18 seconds for most NKG2DLs [38, 45, 55]. Indeed, the increased half-life for NKG2D allows OMCP to effectively antagonize

NKG2D-mediated immunity in our murine infection models despite the lower affinity of the murine NKG2D receptor for OMCP compared to human NKG2D receptor. CPXV infections in primates or rodents that express the higher affinity NKG2D receptor would likely be protected from NKG2D-mediated immunity by OMCP to a greater degree.

To understand the molecular basis for the long half-life of OMCP for NKG2D, we previously determined the structure of OMCP alone, and here, we report the structure of OMCP bound to NKG2D. The structure of OMCP alone was grossly similar to that of host NKG2D ligands, containing an atypical MHCI-like platform domain. Host NKG2D ligands bind with the helices of their platform domains oriented diagonally within the symmetric binding groove of NKG2D. Thus it was expected that OMCP was a viral mimic of host NKG2D ligands and would interact with NKG2D analogously.

The structure of OMCP-NKG2D instead revealed a novel orientation for an NKG2D ligand in the NKG2D binding groove. Alterations within the $\alpha 2$ domain helix allow OMCP to arrange its helices perpendicularly within the binding groove. This reorientation places the H2a and H2b helices directly in contact with the core binding sites of NKG2D and also forms the largest and most continuous binding interface with NKG2D. Because the forces (hydrogen bonds, van der Waals, hydrophobic interactions) that mediate protein-protein interactions are individually weak, a large, continuous interface with high shape complementary allows for a cumulatively strong interaction between proteins. This change in the binding orientation of OMCP reveals how the MHCI-like platform used by host ligands can be adapted by a pathogen to enhance NKG2D binding.

Since host NKG2DLs and OMCP have a similar MHCI-like platform, it is reasonable to wonder why no host ligand has evolved an analogous high-affinity interaction with NKG2D. One likely reason is that the host immune response must be carefully calibrated to balance the need for protection against the threat of autoimmunity. Since the expression of NKG2DLs on the cell surface signals for effector functions, even a small amount of high affinity host ligand on the cell

surface could trigger an immune response, and the resulting tissue damage could be deleterious for the host. Indeed, NKG2D-expressing cells and/or aberrant expression of host NKG2DLs have been implicated in diabetes, celiac disease and rheumatoid arthritis [56-59]. Viruses are not constrained by autoimmune selective pressures. Therefore, CPXV and MPXV were free to evolve a viral NKG2DL with the highest possible affinity to maximize immune evasion potential.

Interestingly, OMCP triggers NKG2D signaling when attached to a target cell membrane, despite the novel orientation of OMCP relative to host NKG2DLs. The interaction of host NKG2DLs with the dimeric NKG2D bears broad structural similarity to the interaction between MHC molecules with their cognate T cell receptors (TCRs). In both cases, the NKG2DL/MHC lies diagonally across the surface created by the dimeric NKG2D/TCR. However, there are several examples of MHC-TCR complexes that, like OMCP-NKG2D, interact with unconventional orientations [60-64]. Several of these complexes involved autoimmune MHC-TCR complexes that were tilted or rotated outside of the normal range for MHC-TCR complexes [60, 64]. While these receptors could induce TCR signaling at high MHC concentrations, they failed to assemble characteristic immunological synapses [65]. A striking example of unconventional binding was found when an *in vitro* peptide library-MHC-TCR (H2-L^d-42F3) screen produced a p3A1-H2-L^d-42F3 complex with an interface rotated ~40° relative to other H2-L^d-42F3 complexes. This rotation places the TCR nearly parallel with the MHC peptide-binding groove and shifted the interface center almost entirely on one of the MHC α helices - an orientation strikingly similar to the interface of OMCP-NKG2D [64]. Interestingly, the p3A1-H2-L^d-42F3 complex failed to induce TCR signaling [64]. Thus, unlike OMCP/NKG2D, the orientation of MHC relative to TCR is an important factor for signaling.

OMCP-NKG2D and p3A1-H2-L^d-42F3 have opposite signaling outcomes, despite having very similar orientations. TCR signaling requires co-receptor binding to either the $\alpha 2/\beta 2$ or $\alpha 3$ domains of MHCII or MHCI, respectively. The failure of p3A1-H2-L^d-42F3 to signal, and of other

unconventional MHC-TCR complexes to form true immunological synapses, is potentially due to the inability of co-receptors to form correct quaternary structures for signaling [63, 64, 66]. Signaling by NKG2D is not known to require co-receptor stimulation and the majority of NKG2DLs lack the co-receptor binding $\alpha 2/\beta 2$ or $\alpha 3$ domains of true MHC molecules. This difference in co-receptor dependency likely explains why OMCP (when attached via GPI anchor) is still competent to stimulate NKG2D-signaling compared to MHC-TCR complexes with unconventional binding orientations. Further, it suggests that clustering of NKG2D on the cell surface is the major determinant of NKG2D-mediated activation.

Materials and Methods

Cell lines and mice.

The generation of Ba/F3 transductants expressing NKG2D, FcRL5, RAE1 δ , MULT1, and OMCP-Thy1.1 was previously described [37, 67]. C57BL/6 (B6) mice were obtained from the National Cancer Institute (Charles River, MA). NKG2D-deficient mice on a B6 background were a kind gift from Bojan Polic (University of Rijeka, Croatia) [68], and FcRL5-deficient mice on a B6 background were obtained from Riccardo Dalla Favera (Columbia University). Mice were maintained under specific pathogen-free conditions and used between 8 and 12 weeks of age. Female mice utilized in survival studies were observed daily after infection for 28 days and moribund mice were euthanized per institutional guidelines.

Ethics Statement.

All experiments were conducted in accordance with institutional guidelines for animal care and use based on the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health. The protocol was approved by the Animal Studies Committee at Washington University (#20110104). Human PBMC were collected under IRB approved protocol #201110275 at

Washington University in St Louis. Informed consent was obtained from all volunteers donating blood.

CPXV and infection of mice.

Brighton Red strain CPXV was obtained from the ATCC and propagated in BS-C-1 cells. Virus was purified from infected BS-C-1 cell lysates by centrifugation through a 36% sucrose gradient. The titer of the viral stock was determined using a standard plaque assay on BS-C-1 cells [69, 70]. Intraperitoneal (i.p.) infections were done at a dose of 2×10^6 pfu CPXV/mouse unless otherwise indicated. For intranasal infections, mice were first anesthetized i.p. with ketamine/xylazine short acting sedative. Mice were infected with the appropriate inoculum of virus in a 50 μ L volume intranasally. All i.n. infections were done at a dose of 2.5×10^4 pfu CPXV/mouse unless otherwise indicated.

Generation of recombinant cowpox viruses.

Recombinant virus lacking expression of OMCP (Δ V018) and its revertant were produced by transient dominant selection [71, 72]. A PCR product encoding the VACV p7.5 5' UTR and promoter was amplified and spliced to another PCR product encoding the *E. coli gpt* sequence. The final product was cloned into plasmid pUC19 to create pUC19 p7.5 *gpt*. OMCP was then amplified in two fragments in order to insert two tandem early termination codons. This product was cloned into pUC19 7.5 *gpt* to create pUC19 p7.5 *gpt* OMCPSTOP. The revertant plasmid pUC19 p7.5 *gpt* OMCP was generated by excising the OMCPSTOP sequence and replacing it with wild-type OMCP. OMCP D132R mutant CPXV was generated in a similar manner with primers to insert a point mutation at D132. The mutated D132R construct was also spliced to *E. coli gpt* and cloned for transient dominant selection.

Confluent CV-1 cells were then infected with CPXV (for CPXV ΔV018 or D132R generation) or CPXV ΔV018 (for revertant generation) at an MOI of 0.5. Two hours post-infection, the cells were transfected with pUC19 p7.5 *gpt* OMCPSTOP for CPXV ΔV018, pUC19 p7.5 *gpt* OMCP-D132R for D132R generation, or pUC19 p7.5 *gpt* OMCP for revertant generation using Lipofectamine2000. The infected/transfected cells were lysed after 48 hours, and *gpt*⁺ viruses were selected by infecting confluent B-SC-1 cells with the transfected virus stock in *gpt* selection medium (DMEM, 2% FCS, 25 μg/mL mycophenolic acid, 250 μg/mL xanthine, and 15 μg/mL hypoxanthine). *Gpt*⁺ viral plaques underwent two further rounds of selection in *gpt* selection media and two subsequent rounds of purification in non-selective media. Finally, plaques were tested for their ability to grow in both selective and non-selective media. Plaques that grew only in non-selective media (indicating loss of the *gpt* marker) were picked and underwent a total of five plaque purifications. Virus stocks were subsequently propagated in B-SC-1 cells and titered following standard protocols using B-SC-1 cells [70].

Identification of NKG2D-binding null mutant D132R. A high throughput *in vitro* selection approach based on combinatorial cell surface display was utilized to identify NKG2D-binding null mutants. The sequence of OMCP was globally mutagenized using error-prone PCR, and the mutated amplicons were spliced to a signal-less Thy1.1 cDNA via overlap extension PCR. This library of mutated OMCPs fused to unmutated Thy1.1 was cloned into the pMXs-IRES-EGFP retroviral transfer vector (kind gift of Toshio Kitamura, University of Tokyo) to generate a molecular library for transduction into Ba/F3 cells. The transductants were then sorted for green fluorescence and anti-Thy1.1 expression to yield a cellular library whose members all had surface expression of OMCP, filtering out mutations giving frameshifts, premature stop codons, and folding-incompetent OMCP. This OMCP library was sorted for NKG2D binding using NKG2D-tetramers. Sorted cells were cloned by limiting dilution and analyzed. The retroviral cassettes of cells lacking or having reduced NKG2D-binding activity were amplified and

sequenced. Utilizing this approach, we identified Asp132 as a critical residue for NKG2D binding.

Protein expression and purification. OMCP_{BR} and human NKG2D expression constructs were previously described [38]. The (D132R) OMCP_{BR} protein was prepared identically to WT OMCP_{BR}. (23D/95D) OMCP-NKG2D complex was reconstituted by oxidative co-refolding from purified inclusion bodies, as described previously [38]. Refolded protein was slowly diluted 10-fold with water and captured on a 5 ml HiTrap Q HP column (GE Healthcare) using a Profinia instrument (Bio-Rad). The captured protein was washed with 50 mM Tris, pH 8.5, 20 mM NaCl and bulk eluted with 50 mM Tris, pH 8.5, 250 mM NaCl. The eluted protein was then concentrated and further purified by gel filtration chromatography on a Superdex S75 column (16/60; Amersham Biosciences). Fractions containing mono-dispersed OMCP-NKG2D complex (~50 KDa) were pooled and buffer exchanged into 25mM Ammonium acetate pH 7.4.

Mammalian-derived proteins were expressed by transient transfection of HEK293F cells (Life Technologies) using PEI. The culture medium was collected 3 days and 6 days after transfection, and then purified by standard Ni-NTA chromatography in accordance with the manufacturer's protocol (Gold Biotechnology) and were subsequently buffer exchanged into phosphate-buffered saline (PBS). Pentamerized proteins were made by fusing the target protein with a C-terminal cartilage oligomeric matrix protein (COMP) domain [73].

Crystallization, data collection and processing. Native protein crystals were grown by hanging drop vapor diffusion at 20°C by streak seeding into a well solution containing 15% PEG 3350, 0.2M MgCl₂, 0.1M Bis-Tris pH 6.75. Crystals were cryoprotected with well solution containing 15% glycerol before flash freezing directly in a liquid nitrogen bath. Diffraction data were collected at the Advanced Light Source synchrotron (beamline 4.2.2). Native (23D/95D) OMCP-hNKG2D crystal diffraction data were collected at 100 K and at a wavelength of 1.00004

Å. Additional diffraction data statistics are summarized in Table 1. Data processing with HKL2000 [74] showed the crystals belonged to the primitive monoclinic space group $P2_1$ (space group #4). The asymmetric unit of the crystal contained two copies of the (23D/95D) OMCP-hNKG2D complex.

Model building and refinement. The structures of human NKG2D (1MPU)[49] and OMCP (4FFE)[38] were used as search models for molecular replacement through Phenix [75]. Iterative refinement and manual rebuilding were performed using Phenix and Coot [76], respectively. Both 2Fo-Fc and Fo-Fc maps were used for manual building and to place solvent molecules. The final model yielded an R_{work} of 16.6% and R_{free} of 21.4%, with 4% of all reflections set aside for free R factor cross-validation. Progress in refinement was also measured using the MOLPROBITY webserver [77]. The final Ramachandran statistics for the model were 98% favored and 0% outliers. Additional refinement statistics are summarized in Table 1. Images of structures were produced using the program PyMol [78].

Structure analysis. Analysis of the contact residues, buried surface area and shape complementarity of the OMCP-NKG2D interface were carried out using the programs Ligplot+ [79], PISA [80] and SC [81]. Structural programs as compiled by the SBGrid consortium [82]. Analysis of NKG2D conservation was performed using the ConSurf server [83-86]. GenBank numbers for species used in Consurf analysis are: Humans (30749494), Borean orangutan (21902299), Chimpanzee (57113989), Gibbon (332232684), Macaque (355785888), Green Monkey (635063485), Common marmoset (380848799), Mouse (148667521), Brown rat (149049263), Guinea Pig (348569092), Ground squirrel (532114387), Deer mouse (589967905), Naked mole rat (512868733), Prairie vole (532053033), European Shrew (505834608), Star-nosed mole (507978716), Chinese hamster (537136230), and Cat (410963826).

Atomic coordinates

The atomic coordinates (accession code 4PDC) have been deposited in the Protein Data Bank, Research Collaboratory for Structural Bioinformatics (Rutgers University, New Brunswick, NJ)

***In vitro* NK cell killing assays.** Splenocytes from C57BL/6 mice were preactivated with 200 U/ml IL-2 for 24 hours and used as cytotoxic effectors against stably transduced Ba/F3 cell lines in standard killing assays. Target cells were carboxyfluorescein succinimidyl ester (CFSE) labeled and co-incubated with activated splenocytes at 37°C, 5% CO₂ for 4 hours at effector:target ratios of 10:1, 20:1, and 40:1. Killing percentage was determined by incorporation of the dead cell exclusion dye 7-amino-actinomycin D (7AAD) in the CFSE⁺ target population as assessed by flow cytometry. Percent specific lysis was calculated using the formula $[(\text{experimental dead \%} - \text{background dead \%}) / (\text{maximum release dead \%} - \text{background dead \%})] \times 100$. Single cell suspensions of splenocytes used in killing assays were generated using standard protocols[87].

Antibodies, and Flow cytometry.

Recombinant OMCP, D132R OMCP, and West Nile Virus glycoprotein domain III (DIII) were produced and biotinylated as previously reported [37, 88]. APC labeled tetramers were made by incubating biotinylated OMCP protein with APC-streptavidin in a 4:1 molar ratio for 10 min at room temperature. APC eFlour 780 anti-CD3 (145-2C11) and eFlour 450 anti-CD19 (1D3) were purchased from eBioscience (San Diego, CA). FITC CD19 (1D3) and PE anti-NK1.1 (PK136) were from BD Biosciences (San Jose, CA). FITC anti-CD5 (53-7.3) was from Biolegend (San Diego, CA). Antibodies and staining of human PBMCs were performed as previously reported [39]. Single cell suspensions of splenocytes and isolation of peritoneal cells and Ficoll purification of human PBMCs were performed using standard protocols [87, 89]. To block

nonspecific binding of antibodies to FcRs, murine cells were incubated in 2.4G2 (anti-FcγRII/III) supernatants (hybridoma from ATCC) prior to staining with labeled Abs. Data was collected on a FACScan flow cytometer (BD Pharmingen) with DXP multicolor upgrade (Cytex Development, Fremont, CA) and FlowJo CE software (TreeStar, Ashland, OR) or FACSCalibur flow cytometer (BD Biosciences) with CellQuest collection software (BD Biosciences). Data analysis was performed using FlowJo software (TreeStar).

Plaque assay.

Tissues harvested from mice were weighed and collected in tubes containing 1mL of DMEM and stored at -80°C until processing. Frozen organs were homogenized using glass dounce homogenizers and collected in 1mL of DMEM. The tissue homogenate was serially diluted in D2.5 media (DMEM, 2.5% FBS, 1% L-glutamine, 1% penicillin-streptomycin, 1% non-essential amino acids) and used in standard plaque assays on BS-C-1 cells [69]. Each organ was titrated on 5x10⁵ BS-C-1 cells plated in wells of 6 well plates in duplicate. Infected cell cultures were maintained in D2.5 media at 37°C, 5% CO₂ for 48 hours. Cells were stained at 48 hours post infection using a crystal violet solution and plaques were counted. Viral titers were calculated from a minimum of 3 dilutions done in duplicate.

Statistical analysis.

Data analysis was done with Microsoft Excel and GraphPad Prism (GraphPad Software, La Jolla, CA). Unless otherwise noted, unpaired, two-tailed t tests were used to determine statistically significant differences. Error bars in the figures represent SDs from the mean value. The Log-rank (Mantel-Cox) test was used in the comparison of all Kaplan Meier survival curves.

Acknowledgements

- 1 E.L., X.W., C.N., and D.H.F. were supported in part by NIH/NIAID R01 AI019687, U19
- 2 AI109948 and NIAID contracts HHSN272200700058C & HHSN272201200026C. M.M.S, T.L.G,
- 3 D.L, and A.R.F were supported in part by NIH/NIAID R01AI073552. A.S.K. supported by NIH
- 4 PO1 AI116501. We thank Helen M. Lazear for help with illustrations

References

1. Hansen TH, Bouvier M. MHC class I antigen presentation: learning from viral evasion strategies. *Nature reviews Immunology*. 2009;9(7):503-13. Epub 2009/06/06. doi: 10.1038/nri2575. PubMed PMID: 19498380.
2. Griffin BD, Verweij MC, Wiertz EJ. Herpesviruses and immunity: the art of evasion. *Vet Microbiol*. 2010;143(1):89-100. Epub 2010/03/23. doi: 10.1016/j.vetmic.2010.02.017. PubMed PMID: 20303681.
3. Karre K, Ljunggren HG, Piontek G, Kiessling R. Selective rejection of H-2-deficient lymphoma variants suggests alternative immune defence strategy. *Nature*. 1986;319(6055):675-8. Epub 1986/02/20. doi: 10.1038/319675a0. PubMed PMID: 3951539.
4. Orange JS, Fasset MS, Koopman LA, Boyson JE, Strominger JL. Viral evasion of natural killer cells. *Nature immunology*. 2002;3(11):1006-12. Epub 2002/10/31. doi: 10.1038/ni1102-1006. PubMed PMID: 12407408.
5. Lisnic VJ, Krmpotic A, Jonjic S. Modulation of natural killer cell activity by viruses. *Current opinion in microbiology*. 2010;13(4):530-9. Epub 2010/06/19. doi: 10.1016/j.mib.2010.05.011. PubMed PMID: 20558100; PubMed Central PMCID: PMC2920364.
6. Finton KA, Strong RK. Structural insights into activation of antiviral NK cell responses. *Immunological reviews*. 2012;250(1):239-57. Epub 2012/10/11. doi: 10.1111/j.1600-065X.2012.01168.x. PubMed PMID: 23046134; PubMed Central PMCID: PMC3471384.
7. Li Y, Mariuzza RA. Structural Basis for Recognition of Cellular and Viral Ligands by NK Cell Receptors. *Front Immunol*. 2014;5:123. Epub 2014/04/12. doi: 10.3389/fimmu.2014.00123. PubMed PMID: 24723923; PubMed Central PMCID: PMC3972465.
8. Raulet DH. Roles of the NKG2D immunoreceptor and its ligands. *Nature reviews Immunology*. 2003;3(10):781-90. Epub 2003/10/03. doi: 10.1038/nri1199. PubMed PMID: 14523385.
9. Draghi M, Pashine A, Sanjanwala B, Gendzekhadze K, Cantoni C, Cosman D, et al. NKp46 and NKG2D recognition of infected dendritic cells is necessary for NK cell activation in the human response to influenza infection. *J Immunol*. 2007;178(5):2688-98. Epub 2007/02/22. PubMed PMID: 17312110.
10. Pappworth IY, Wang EC, Rowe M. The switch from latent to productive infection in epstein-barr virus-infected B cells is associated with sensitization to NK cell killing. *Journal of virology*. 2007;81(2):474-82. Epub 2006/11/03. doi: 10.1128/JVI.01777-06. PubMed PMID: 17079298; PubMed Central PMCID: PMC1797427.
11. Welte SA, Sinzger C, Lutz SZ, Singh-Jasuja H, Sampaio KL, Eknigk U, et al. Selective intracellular retention of virally induced NKG2D ligands by the human cytomegalovirus UL16 glycoprotein. *European journal of immunology*. 2003;33(1):194-203. Epub 2003/02/21. doi: 10.1002/immu.200390022. PubMed PMID: 12594848.
12. Ward J, Bonaparte M, Sacks J, Guterman J, Fogli M, Mavilio D, et al. HIV modulates the expression of ligands important in triggering natural killer cell cytotoxic responses on infected primary T-cell blasts. *Blood*. 2007;110(4):1207-14. Epub 2007/05/22. doi: 10.1182/blood-2006-06-028175. PubMed PMID: 17513617; PubMed Central PMCID: PMC1939902.
13. Obeidy P, Sharland AF. NKG2D and its ligands. *Int J Biochem Cell Biol*. 2009;41(12):2364-7. Epub 2009/07/28. doi: 10.1016/j.biocel.2009.07.005. PubMed PMID: 19631280.
14. Eagle RA, Trowsdale J. Promiscuity and the single receptor: NKG2D. *Nature reviews Immunology*. 2007;7(9):737-44. Epub 2007/08/04. doi: 10.1038/nri2144. PubMed PMID: 17673918.
15. Lodoen M, Ogasawara K, Hamerman JA, Arase H, Houchins JP, Mocarski ES, et al. NKG2D-mediated natural killer cell protection against cytomegalovirus is impaired by viral gp40 modulation of retinoic acid early inducible 1 gene molecules. *The Journal of experimental medicine*. 2003;197(10):1245-53. Epub 2003/05/21. doi: 10.1084/jem.20021973. PubMed PMID: 12756263; PubMed Central PMCID: PMC2193789.
16. Lodoen MB, Abenes G, Umamoto S, Houchins JP, Liu F, Lanier LL. The cytomegalovirus m155 gene product subverts natural killer cell antiviral protection by disruption of H60-NKG2D interactions. *The Journal of experimental medicine*. 2004;200(8):1075-81. Epub 2004/10/13. doi: 10.1084/jem.20040583. PubMed PMID: 15477345; PubMed Central PMCID: PMC2211837.

17. Krmpotic A, Hasan M, Loewendorf A, Saulig T, Halenius A, Lenac T, et al. NK cell activation through the NKG2D ligand MULT-1 is selectively prevented by the glycoprotein encoded by mouse cytomegalovirus gene m145. *The Journal of experimental medicine*. 2005;201(2):211-20. Epub 2005/01/12. doi: 10.1084/jem.20041617. PubMed PMID: 15642742; PubMed Central PMCID: PMC2212792.
18. Lenac T, Budt M, Arapovic J, Hasan M, Zimmermann A, Simic H, et al. The herpesviral Fc receptor fcr-1 down-regulates the NKG2D ligands MULT-1 and H60. *The Journal of experimental medicine*. 2006;203(8):1843-50. Epub 2006/07/13. doi: 10.1084/jem.20060514. PubMed PMID: 16831899; PubMed Central PMCID: PMC2118374.
19. Cosman D, Mullberg J, Sutherland CL, Chin W, Armitage R, Fanslow W, et al. ULBPs, novel MHC class I-related molecules, bind to CMV glycoprotein UL16 and stimulate NK cytotoxicity through the NKG2D receptor. *Immunity*. 2001;14(2):123-33. Epub 2001/03/10. PubMed PMID: 11239445.
20. Chalupny NJ, Rein-Weston A, Dosch S, Cosman D. Down-regulation of the NKG2D ligand MICA by the human cytomegalovirus glycoprotein UL142. *Biochem Biophys Res Commun*. 2006;346(1):175-81. Epub 2006/06/06. doi: 10.1016/j.bbrc.2006.05.092. PubMed PMID: 16750166.
21. Thomas M, Boname JM, Field S, Nejentsev S, Salio M, Cerundolo V, et al. Down-regulation of NKG2D and NKp80 ligands by Kaposi's sarcoma-associated herpesvirus K5 protects against NK cell cytotoxicity. *Proceedings of the National Academy of Sciences of the United States of America*. 2008;105(5):1656-61. Epub 2008/01/31. doi: 10.1073/pnas.0707883105. PubMed PMID: 18230726; PubMed Central PMCID: PMC2234200.
22. Cerboni C, Neri F, Casartelli N, Zingoni A, Cosman D, Rossi P, et al. Human immunodeficiency virus 1 Nef protein downmodulates the ligands of the activating receptor NKG2D and inhibits natural killer cell-mediated cytotoxicity. *The Journal of general virology*. 2007;88(Pt 1):242-50. Epub 2006/12/16. doi: 10.1099/vir.0.82125-0. PubMed PMID: 17170457.
23. Wen C, He X, Ma H, Hou N, Wei C, Song T, et al. Hepatitis C virus infection downregulates the ligands of the activating receptor NKG2D. *Cellular & molecular immunology*. 2008;5(6):475-8. Epub 2009/01/03. doi: 10.1038/cmi.2008.60. PubMed PMID: 19118515.
24. Stern-Ginossar N, Elefant N, Zimmermann A, Wolf DG, Saleh N, Biton M, et al. Host immune system gene targeting by a viral miRNA. *Science*. 2007;317(5836):376-81. Epub 2007/07/21. doi: 10.1126/science.1140956. PubMed PMID: 17641203.
25. Nachmani D, Stern-Ginossar N, Sarid R, Mandelboim O. Diverse herpesvirus microRNAs target the stress-induced immune ligand MICB to escape recognition by natural killer cells. *Cell host & microbe*. 2009;5(4):376-85. Epub 2009/04/22. doi: 10.1016/j.chom.2009.03.003. PubMed PMID: 19380116.
26. Bauman Y, Nachmani D, Vicenshtein A, Tsukerman P, Drayman N, Stern-Ginossar N, et al. An identical miRNA of the human JC and BK polyoma viruses targets the stress-induced ligand ULBP3 to escape immune elimination. *Cell host & microbe*. 2011;9(2):93-102. Epub 2011/02/16. doi: 10.1016/j.chom.2011.01.008. PubMed PMID: 21320692.
27. Gainey MD, Rivenbark JG, Cho H, Yang L, Yokoyama WM. Viral MHC class I inhibition evades CD8+ T-cell effector responses in vivo but not CD8+ T-cell priming. *Proceedings of the National Academy of Sciences of the United States of America*. 2012;109(47):E3260-7. Epub 2012/11/01. doi: 10.1073/pnas.1217111109. PubMed PMID: 23112205; PubMed Central PMCID: PMC3511129.
28. Byun M, Verweij MC, Pickup DJ, Wiertz EJ, Hansen TH, Yokoyama WM. Two mechanistically distinct immune evasion proteins of cowpox virus combine to avoid antiviral CD8 T cells. *Cell host & microbe*. 2009;6(5):422-32. Epub 2009/11/18. doi: 10.1016/j.chom.2009.09.012. PubMed PMID: 19917497; PubMed Central PMCID: PMC2791900.
29. Byun M, Wang X, Pak M, Hansen TH, Yokoyama WM. Cowpox virus exploits the endoplasmic reticulum retention pathway to inhibit MHC class I transport to the cell surface. *Cell host & microbe*. 2007;2(5):306-15. Epub 2007/11/17. doi: 10.1016/j.chom.2007.09.002. PubMed PMID: 18005752.
30. McCoy WH, Wang X, Yokoyama WM, Hansen TH, Fremont DH. Cowpox virus employs a two-pronged strategy to outflank MHCI antigen presentation. *Molecular immunology*. 2013. Epub 2013/01/15. doi: 10.1016/j.molimm.2012.11.011. PubMed PMID: 23312338.

31. McCoy WHt, Wang X, Yokoyama WM, Hansen TH, Fremont DH. Structural mechanism of ER retrieval of MHC class I by cowpox. *PLoS biology*. 2012;10(11):e1001432. Epub 2012/12/05. doi: 10.1371/journal.pbio.1001432. PubMed PMID: 23209377; PubMed Central PMCID: PMC3507924.
32. Alzhanova D, Edwards DM, Hammarlund E, Scholz IG, Horst D, Wagner MJ, et al. Cowpox virus inhibits the transporter associated with antigen processing to evade T cell recognition. *Cell host & microbe*. 2009;6(5):433-45. Epub 2009/11/18. doi: 10.1016/j.chom.2009.09.013. PubMed PMID: 19917498; PubMed Central PMCID: PMC2791678.
33. Dasgupta A, Hammarlund E, Slifka MK, Fruh K. Cowpox virus evades CTL recognition and inhibits the intracellular transport of MHC class I molecules. *J Immunol*. 2007;178(3):1654-61. Epub 2007/01/24. PubMed PMID: 17237415.
34. Luteijn RD, Hoelen H, Kruse E, van Leeuwen WF, Grootens J, Horst D, et al. Cowpox Virus Protein CPXV012 Eludes CTLs by Blocking ATP Binding to TAP. *J Immunol*. 2014;193(4):1578-89. Epub 2014/07/16. doi: 10.4049/jimmunol.1400964. PubMed PMID: 25024387.
35. Fang M, Lanier LL, Sigal LJ. A role for NKG2D in NK cell-mediated resistance to poxvirus disease. *PLoS pathogens*. 2008;4(2):e30. Epub 2008/02/13. doi: 10.1371/journal.ppat.0040030. PubMed PMID: 18266471; PubMed Central PMCID: PMC2233669.
36. Song H, Josleyn N, Janosko K, Skinner J, Reeves RK, Cohen M, et al. Monkeypox virus infection of rhesus macaques induces massive expansion of natural killer cells but suppresses natural killer cell functions. *PloS one*. 2013;8(10):e77804. Epub 2013/10/23. doi: 10.1371/journal.pone.0077804. PubMed PMID: 24147080; PubMed Central PMCID: PMC3798392.
37. Campbell JA, Trossman DS, Yokoyama WM, Carayannopoulos LN. Zoonotic orthopoxviruses encode a high-affinity antagonist of NKG2D. *J Exp Med*. 2007;204(6):1311-7. doi: 10.1084/jem.20062026. PubMed PMID: 17548517.
38. Lazear E, Peterson LW, Nelson CA, Fremont DH. Crystal structure of the cowpox virus-encoded NKG2D ligand OMCP. *Journal of virology*. 2013;87(2):840-50. Epub 2012/11/02. doi: 10.1128/JVI.01948-12. PubMed PMID: 23115291; PubMed Central PMCID: PMC3554055.
39. Campbell JA, Davis RS, Lilly LM, Fremont DH, French AR, Carayannopoulos LN. Cutting edge: FcR-like 5 on innate B cells is targeted by a poxvirus MHC class I-like immuno-evasin. *J Immunol*. 2010;185(1):28-32. Epub 2010/06/04. doi: 10.4049/jimmunol.1000240. PubMed PMID: 20519648.
40. Carayannopoulos LN, Naidenko OV, Kinder J, Ho EL, Fremont DH, Yokoyama WM. Ligands for murine NKG2D display heterogeneous binding behavior. *European journal of immunology*. 2002;32(3):597-605. Epub 2002/02/22. doi: 10.1002/1521-4141(200203)32:3<597::AID-IMMU597>3.0.CO;2-E. PubMed PMID: 11857333.
41. Mistry AR, O'Callaghan CA. Regulation of ligands for the activating receptor NKG2D. *Immunology*. 2007;121(4):439-47. Epub 2007/07/07. doi: 10.1111/j.1365-2567.2007.02652.x. PubMed PMID: 17614877; PubMed Central PMCID: PMC2265965.
42. Strong RK, McFarland BJ. NKG2D and Related Immunoreceptors. *Adv Protein Chem*. 2004;68:281-312. Epub 2004/10/27. doi: 10.1016/S0065-3233(04)68008-9. PubMed PMID: 15500864.
43. Deng L, Mariuzza RA. Structural basis for recognition of MHC and MHC-like ligands by natural killer cell receptors. *Semin Immunol*. 2006;18(3):159-66. Epub 2006/06/02. doi: 10.1016/j.smim.2006.03.004. PubMed PMID: 16737824; PubMed Central PMCID: PMC2519613.
44. Li P, McDermott G, Strong RK. Crystal structures of RAE-1beta and its complex with the activating immunoreceptor NKG2D. *Immunity*. 2002;16(1):77-86. Epub 2002/02/05. PubMed PMID: 11825567.
45. Li P, Morris DL, Willcox BE, Steinle A, Spies T, Strong RK. Complex structure of the activating immunoreceptor NKG2D and its MHC class I-like ligand MICA. *Nature immunology*. 2001;2(5):443-51. Epub 2001/04/27. doi: 10.1038/87757. PubMed PMID: 11323699.
46. Li P, Willie ST, Bauer S, Morris DL, Spies T, Strong RK. Crystal structure of the MHC class I homolog MIC-A, a gammadelta T cell ligand. *Immunity*. 1999;10(5):577-84. Epub 1999/06/15. PubMed PMID: 10367903.

47. Radaev S, Rostro B, Brooks AG, Colonna M, Sun PD. Conformational plasticity revealed by the cocrystal structure of NKG2D and its class I MHC-like ligand ULBP3. *Immunity*. 2001;15(6):1039-49. Epub 2002/01/05. PubMed PMID: 11754823.
48. Adams EJ, Luoma AM. The adaptable major histocompatibility complex (MHC) fold: structure and function of nonclassical and MHC class I-like molecules. *Annu Rev Immunol*. 2013;31:529-61. Epub 2013/01/10. doi: 10.1146/annurev-immunol-032712-095912. PubMed PMID: 23298204.
49. McFarland BJ, Kortemme T, Yu SF, Baker D, Strong RK. Symmetry recognizing asymmetry: analysis of the interactions between the C-type lectin-like immunoreceptor NKG2D and MHC class I-like ligands. *Structure*. 2003;11(4):411-22. Epub 2003/04/08. PubMed PMID: 12679019.
50. Chapman JL, Nichols DK, Martinez MJ, Raymond JW. Animal models of orthopoxvirus infection. *Veterinary pathology*. 2010;47(5):852-70. doi: 10.1177/0300985810378649. PubMed PMID: 20682806.
51. Lefkowitz EJ, Upton C, Changayil SS, Buck C, Traktman P, Buller RM. Poxvirus Bioinformatics Resource Center: a comprehensive Poxviridae informational and analytical resource. *Nucleic acids research*. 2005;33(Database issue):D311-6. Epub 2004/12/21. doi: 10.1093/nar/gki110. PubMed PMID: 15608205; PubMed Central PMCID: PMC540064.
52. Strong RK. Asymmetric ligand recognition by the activating natural killer cell receptor NKG2D, a symmetric homodimer. *Molecular immunology*. 2002;38(14):1029-37. Epub 2002/04/17. PubMed PMID: 11955595.
53. Radaev S, Sun PD. Structure and function of natural killer cell surface receptors. *Annu Rev Biophys Biomol Struct*. 2003;32:93-114. Epub 2002/12/10. doi: 10.1146/annurev.biophys.32.110601.142347. PubMed PMID: 12471063.
54. Copeland RA, Pompliano DL, Meek TD. Drug-target residence time and its implications for lead optimization. *Nat Rev Drug Discov*. 2006;5(9):730-9. Epub 2006/08/05. doi: 10.1038/nrd2082. PubMed PMID: 16888652.
55. O'Callaghan CA, Cerwenka A, Willcox BE, Lanier LL, Bjorkman PJ. Molecular competition for NKG2D: H60 and RAE1 compete unequally for NKG2D with dominance of H60. *Immunity*. 2001;15(2):201-11. Epub 2001/08/25. PubMed PMID: 11520456.
56. Groh V, Bruhl A, El-Gabalawy H, Nelson JL, Spies T. Stimulation of T cell autoreactivity by anomalous expression of NKG2D and its MIC ligands in rheumatoid arthritis. *Proceedings of the National Academy of Sciences of the United States of America*. 2003;100(16):9452-7. Epub 2003/07/25. doi: 10.1073/pnas.1632807100. PubMed PMID: 12878725; PubMed Central PMCID: PMC170939.
57. Hue S, Mention JJ, Monteiro RC, Zhang S, Cellier C, Schmitz J, et al. A direct role for NKG2D/MICA interaction in villous atrophy during celiac disease. *Immunity*. 2004;21(3):367-77. Epub 2004/09/11. doi: 10.1016/j.immuni.2004.06.018. PubMed PMID: 15357948.
58. Meresse B, Chen Z, Ciszewski C, Tretiakova M, Bhagat G, Krausz TN, et al. Coordinated induction by IL15 of a TCR-independent NKG2D signaling pathway converts CTL into lymphokine-activated killer cells in celiac disease. *Immunity*. 2004;21(3):357-66. Epub 2004/09/11. doi: 10.1016/j.immuni.2004.06.020. PubMed PMID: 15357947.
59. Ogasawara K, Hamerman JA, Hsin H, Chikuma S, Bour-Jordan H, Chen T, et al. Impairment of NK cell function by NKG2D modulation in NOD mice. *Immunity*. 2003;18(1):41-51. Epub 2003/01/18. PubMed PMID: 12530974.
60. Hahn M, Nicholson MJ, Pyrdol J, Wucherpfennig KW. Unconventional topology of self peptide-major histocompatibility complex binding by a human autoimmune T cell receptor. *Nature immunology*. 2005;6(5):490-6. Epub 2005/04/12. doi: 10.1038/ni1187. PubMed PMID: 15821740; PubMed Central PMCID: PMC3415330.
61. Sethi DK, Schubert DA, Anders AK, Heroux A, Bonsor DA, Thomas CP, et al. A highly tilted binding mode by a self-reactive T cell receptor results in altered engagement of peptide and MHC. *The Journal of experimental medicine*. 2011;208(1):91-102. Epub 2011/01/05. doi: 10.1084/jem.20100725. PubMed PMID: 21199956; PubMed Central PMCID: PMC3023130.
62. Wucherpfennig KW, Call MJ, Deng L, Mariuzza R. Structural alterations in peptide-MHC recognition by self-reactive T cell receptors. *Curr Opin Immunol*. 2009;21(6):590-5. Epub

- 2009/08/25. doi: 10.1016/j.coi.2009.07.008. PubMed PMID: 19699075; PubMed Central PMCID: PMC2787854.
63. Yin Y, Li Y, Mariuzza RA. Structural basis for self-recognition by autoimmune T-cell receptors. *Immunological reviews*. 2012;250(1):32-48. Epub 2012/10/11. doi: 10.1111/imr.12002. PubMed PMID: 23046121.
64. Adams JJ, Narayanan S, Liu B, Birnbaum ME, Kruse AC, Bowerman NA, et al. T cell receptor signaling is limited by docking geometry to peptide-major histocompatibility complex. *Immunity*. 2011;35(5):681-93. Epub 2011/11/22. doi: 10.1016/j.immuni.2011.09.013. PubMed PMID: 22101157; PubMed Central PMCID: PMC3253265.
65. Schubert DA, Gordo S, Sabatino JJ, Jr., Vardhana S, Gagnon E, Sethi DK, et al. Self-reactive human CD4 T cell clones form unusual immunological synapses. *The Journal of experimental medicine*. 2012;209(2):335-52. Epub 2012/02/09. doi: 10.1084/jem.20111485. PubMed PMID: 22312112; PubMed Central PMCID: PMC3280872.
66. Li Y, Yin Y, Mariuzza RA. Structural and biophysical insights into the role of CD4 and CD8 in T cell activation. *Front Immunol*. 2013;4:206. Epub 2013/07/26. doi: 10.3389/fimmu.2013.00206. PubMed PMID: 23885256; PubMed Central PMCID: PMC3717711.
67. Campbell JA, Davis RS, Lilly LM, Fremont DH, French AR, Carayannopoulos LN. Cutting Edge: FcR-Like 5 on Innate B Cells Is Targeted by a Poxvirus MHC Class I-Like Immuno-evasin. *The Journal of Immunology*. 2010;185(1):28-32. doi: 10.4049/jimmunol.1000240.
68. Zafirova B, Mandarić S, Antulov R, Krmpotić A, Jonsson H, Yokoyama WM, et al. Altered NK cell development and enhanced NK cell-mediated resistance to mouse cytomegalovirus in NKG2D-deficient mice. *Immunity*. 2009;31(2):270-82. doi: 10.1016/j.immuni.2009.06.017. PubMed PMID: 19631564.
69. Earl PL, Cooper N, Wyatt LS, Moss B, Carroll MW. Preparation of cell cultures and vaccinia virus stocks. *Current protocols in protein science / editorial board, John E Coligan [et al]*. 2001;Chapter 5:Unit5.12. doi: 10.1002/0471140864.ps0512s13. PubMed PMID: 18429178.
70. Earl PL, Cooper N, Wyatt LS, Moss B, Carroll MW. Preparation of cell cultures and vaccinia virus stocks. *Current protocols in protein science / editorial board, John E Coligan [et al]*. 2001;Chapter 5:Unit5 12. Epub 2008/04/23. doi: 10.1002/0471140864.ps0512s13. PubMed PMID: 18429178.
71. Falkner FG, Moss B. Transient dominant selection of recombinant vaccinia viruses. *Journal of virology*. 1990;64(6):3108-11. Epub 1990/06/01. PubMed PMID: 2159565; PubMed Central PMCID: PMC249504.
72. Kato SE, Greco FA, Damaso CR, Condit RC, Moussatche N. An alternative genetic method to test essential vaccinia virus early genes. *Journal of virological methods*. 2004;115(1):31-40. Epub 2003/12/06. PubMed PMID: 14656458.
73. Tomschy A, Fauser C, Landwehr R, Engel J. Homophilic adhesion of E-cadherin occurs by a co-operative two-step interaction of N-terminal domains. *EMBO J*. 1996;15(14):3507-14. Epub 1996/07/15. PubMed PMID: 8670853; PubMed Central PMCID: PMCPMC451947.
74. Otwinowski Z, Minor W. Processing of X-ray diffraction data collected in oscillation mode. *Macromolecular Crystallography, Pt A*. 1997;276:307-26. PubMed PMID: ISI:A1997BH42P00020.
75. Adams PD, Grosse-Kunstleve RW, Hung LW, Ioerger TR, McCoy AJ, Moriarty NW, et al. PHENIX: building new software for automated crystallographic structure determination. *Acta crystallographica Section D, Biological crystallography*. 2002;58(Pt 11):1948-54. Epub 2002/10/24. PubMed PMID: 12393927.
76. Emsley P, Cowtan K. Coot: model-building tools for molecular graphics. *Acta crystallographica Section D, Biological crystallography*. 2004;60(Pt 12 Pt 1):2126-32. Epub 2004/12/02. doi: 10.1107/S0907444904019158. PubMed PMID: 15572765.
77. Chen VB, Arendall WB, 3rd, Headd JJ, Keedy DA, Immormino RM, Kapral GJ, et al. MolProbity: all-atom structure validation for macromolecular crystallography. *Acta crystallographica Section D, Biological crystallography*. 2010;66(Pt 1):12-21. Epub 2010/01/09. doi: 10.1107/S0907444909042073. PubMed PMID: 20057044; PubMed Central PMCID: PMC2803126.
78. Schrodinger, LLC. The PyMOL Molecular Graphics System, Version 1.3r1. 2010.

79. Laskowski RA, Swindells MB. LigPlot+: multiple ligand-protein interaction diagrams for drug discovery. *Journal of chemical information and modeling*. 2011;51(10):2778-86. Epub 2011/09/17. doi: 10.1021/ci200227u. PubMed PMID: 21919503.
80. Krissinel E, Henrick K. Inference of macromolecular assemblies from crystalline state. *Journal of molecular biology*. 2007;372(3):774-97. Epub 2007/08/08. doi: 10.1016/j.jmb.2007.05.022. PubMed PMID: 17681537.
81. Lawrence MC, Colman PM. Shape complementarity at protein/protein interfaces. *Journal of molecular biology*. 1993;234(4):946-50. Epub 1993/12/20. doi: 10.1006/jmbi.1993.1648. PubMed PMID: 8263940.
82. Morin A, Eisenbraun B, Key J, Sanschagrin PC, Timony MA, Ottaviano M, et al. Collaboration gets the most out of software. *eLife*. 2013;2:e01456. Epub 2013/09/17. doi: 10.7554/eLife.01456. PubMed PMID: 24040512; PubMed Central PMCID: PMC3771563.
83. Ashkenazy H, Erez E, Martz E, Pupko T, Ben-Tal N. ConSurf 2010: calculating evolutionary conservation in sequence and structure of proteins and nucleic acids. *Nucleic acids research*. 2010;38(Web Server issue):W529-33. Epub 2010/05/19. doi: 10.1093/nar/gkq399. PubMed PMID: 20478830; PubMed Central PMCID: PMC2896094.
84. Landau M, Mayrose I, Rosenberg Y, Glaser F, Martz E, Pupko T, et al. ConSurf 2005: the projection of evolutionary conservation scores of residues on protein structures. *Nucleic acids research*. 2005;33(Web Server issue):W299-302. Epub 2005/06/28. doi: 10.1093/nar/gki370. PubMed PMID: 15980475; PubMed Central PMCID: PMC1160131.
85. Glaser F, Pupko T, Paz I, Bell RE, Bechor-Shental D, Martz E, et al. ConSurf: identification of functional regions in proteins by surface-mapping of phylogenetic information. *Bioinformatics*. 2003;19(1):163-4. Epub 2002/12/25. PubMed PMID: 12499312.
86. Celniker G, Nimrod G, Ashkenazy H, Glaser F, Martz E, Mayrose I, et al. ConSurf: Using Evolutionary Data to Raise Testable Hypotheses about Protein Function. *Isr J Chem*. 2013;53(3-4):199-206. doi: 10.1002/ijch.201200096. PubMed PMID: ISI:000317859800010.
87. Dokun AO, Kim S, Smith HR, Kang HS, Chu DT, Yokoyama WM. Specific and nonspecific NK cell activation during virus infection. *Nat Immunol*. 2001;2(10):951-6. doi: 10.1038/ni714. PubMed PMID: 11550009.
88. Nybakken GE, Oliphant T, Johnson S, Burke S, Diamond MS, Fremont DH. Structural basis of West Nile virus neutralization by a therapeutic antibody. *Nature*. 2005;437(7059):764-9. Epub 2005/09/30. doi: nature03956 [pii] 10.1038/nature03956. PubMed PMID: 16193056.
89. Zhang X, Goncalves R, Mosser DM. The isolation and characterization of murine macrophages. *Curr Protoc Immunol*. 2008;Chapter 14:Unit 14.1. doi: 10.1002/0471142735.im1401s83. PubMed PMID: 19016445.
90. Stewart DE, Sarkar A, Wampler JE. Occurrence and role of cis peptide bonds in protein structures. *Journal of molecular biology*. 1990;214(1):253-60. Epub 1990/07/05. doi: 10.1016/0022-2836(90)90159-J. PubMed PMID: 2370664.
91. Craveur P, Joseph AP, Poulain P, de Brevern AG, Rebehmed J. Cis-trans isomerization of omega dihedrals in proteins. *Amino Acids*. 2013;45(2):279-89. Epub 2013/06/04. doi: 10.1007/s00726-013-1511-3. PubMed PMID: 23728840.

1 **Figure Legends**

2

3 **Figure 1. Cowpox virus lacking OMCP is significantly attenuated *in vivo*.** WT
4 CPXV **(A)** is more virulent than $\Delta V018$ CPXV **(B)**, as evidenced by increased lethality at
5 all tested doses. The revertant CPXV control **(C)** phenocopies WT CPXV as is
6 expected. 8-12 week old female WT B6 mice were infected with 2×10^6 (solid line), 1×10^6
7 (dashed line), or 0.6×10^6 (dotted line) of the indicated CPXV strain, and survival was
8 assessed daily for 28 days. Data are aggregated from two-seven independent
9 experiments with 5-10 mice per group.

10

11 **Figure 2. The structure of OMCP in complex with NKG2D.** **(A)** OMCP bound to
12 NKG2D. OMCP is colored magenta and the protomers of NKG2D are colored cyan ("A")
13 and yellow ("B"). NKG2D^A makes contacts primarily with the H2a helix and NKG2D^B
14 with H2b. Mutations introduced to facilitate alternate crystal packing are shown in red.
15 The S193-S194 bond is shown as a ball on each NKG2D protomer. The asparagines of
16 putative hNKG2D glycosylation sites are shown in orange. The asparagine of the
17 confirmed N-glycan site of OMCP is shown green (data not shown) **(B)** View of the
18 interface between OMCP-NKG2D. The $\alpha 2$ domain of OMCP is shown in the front with
19 the $\alpha 1$ domain behind. OMCP and NKG2D are shown with cartoon representations for
20 the main chain, with the side chains of contact residues shown as sticks. Hydrogen
21 bonds and salt bridges are indicated with green dotted lines.

22

23 **Figure 3. The interface of OMCP and NKG2D.** **(A)** The local environment of the
24 OMCP-NKG2D binding interface surrounding the D132R residue. The D132R mutation
25 ablates OMCP-NKG2D binding. **(B)** APC labeled OMCP tetramers (solid line) were
26 used to stain splenic NK cells (NK1.1+, CD3-) and peritoneal B1-a B cells (CD19+,

1 CD5+), Staining with APC labeled WNV DIII tetramer control shown in gray histograms.
2 Representative results from three independent experiments.

3

4 **Figure 4. NKG2D mediates all of the virulence of OMCP during intranasal CPXV**
5 **infection.** WT CPXV (solid line), Δ V018 CPXV (dashed line) and D132R OMCP CPXV
6 (dotted line) infected via the intranasal route differentially influence survival in **(A)** WT
7 B6, **(B)** NKG2D-deficient, and **(C)** FcRL5-deficient mice. Kaplan-Meier analysis shows
8 survival of female mice following intranasal infection with 2.5×10^4 pfu of the designated
9 CPXV strain. The data are aggregated from two to four independent experiments. The
10 Log-rank (Mantel-Cox) test was used in the comparison of all Kaplan Meier survival
11 curves. NS not significant, **p<0.01 ***p<0.001

12

13 **Figure 5. Glycosylated OMCP binds to NKG2D only.** **(A)** Sorted Ba/Fc cells
14 transduced (EGFP positive) with NKG2D or FcRL5 were stained with biotinylated
15 monomeric OMCPs or with antibodies to NKG2D or FcRL5. A representative experiment
16 is shown. **(B)** Pentamers of OMCP, (D132R) OMCP, or ULBP3 were incubated with
17 ficoll-purified human PBMCs. PBMCs were stained with antibodies to CD56, CD19,
18 CD3, and CD16. NK (CD56+, CD3-), T (CD3+), and B (CD19+, CD3-) cells were then
19 assessed by flow cytometry for protein binding. Representative results from four
20 independent experiments.

21

22 **Figure 6. Differences in the $\beta 5'$ - $\beta 5$ loop (L2) of human and murine NKG2D. (A-B)**
23 Superimposition of mNKG2D (grey) (PDB ID: 1HQ8) with the structure of OMCP-
24 hNKG2D (yellow and cyan). Core binding residues Y152 (Y168) and Y199 (Y215) are
25 positionally conserved, while core binding residue M184 (I200) is not. **(C)** Surface
26 representation of OMCP (magenta) interacting with the $\beta 5'$ - $\beta 5$ loop. **(D)** Conservation of

M184 and Q185. Only the NKG2D of mice, rats, guinea pigs, and flying foxes (not shown) differ. Conservation score is as computed by the ConSurf server.

Figure 7. A novel NKG2D binding adaptation. Surface representation of NKG2D and surface and cartoon representations of OMCP, MICA and ULBP3. Buried surface areas for NKG2D^A and NKG2D^B are indicated in cyan and yellow, respectively. Buried surface area by NKG2D is indicated for OMCP (magenta), MICA (green), and ULBP3 (orange). The core binding residues of NKG2D and NKG2D-binding elements of NKG2DLs are indicated. Alignment by secondary structure of NKG2DLs (PDB ID: OMCP (4FFE), MICA (1HYR), MICB (1JE6), ULBP3 (1KCG) and RAE-1 β (1JSK)). Contact residues are indicated for OMCP (magenta), MICA (green), ULBP3 (orange) and RAE-1 β (bold and italics). Secondary structure elements are noted above the sequence (arrow for beta sheets, cylinders for alpha helices). Predicted glycan sites are highlighted in black.

Figure 8. Activation of NK cells by cell-associated OMCP. Model depicting NKG2D interaction with (A) host, (B) cancer-induced, (C) viral, or (D) chimeric ligands. Binding interactions that lead to NKG2D-mediated signaling are indicated by DAP10 tyrosine phosphorylation (red filled circles). (E) IL-2-activated splenocytes were used as cytotoxic effectors against stably transduced Ba/F3 cell lines. Splenocytes were activated with 200 U/ml of IL-2 for 24 hours. Labeled target cells were co-incubated with activated splenocytes for 4 hours at effector:target ratios of 10:1, 20:1, and 40:1. Killing was measured by incorporation of 7AAD by CFSE-labeled target cells using flow cytometry. Representative data from five independent experiments is shown

Figure S1. Δ V018 virus and D132R virus do not have a growth defect *in vitro*.

Multi-step growth curves for WT CPXV, Δ V018, D132R, and revertant viruses show that

1 all four cowpox virus strains have similar replication kinetics in a multi-step growth curve.
2 B-SC-1 cells were infected at an MOI of 0.01 with the indicated viruses. Infected cells
3 were harvested at the indicated time points and freeze-thawed three times. Lysates
4 were titrated by plaque assay on B-SC-1 monolayers. Results are representative of three
5 experiments done in duplicate.

6

7 **Figure S2. Electron density supporting a *cis* peptide conformation.** Stereo view of
8 the $\beta 5$ - $\beta 6$ loop of hNKG2D. Residues 193-Ala-Ser-Ser-Phe-Lys-197 is displayed for the
9 OMCP-hNKG2D structure (yellow) and the structure of hNKG2D alone (grey). The 2Fo-
10 Fc map for OMCP-hNKG2D is displayed at 2σ . *Cis* conformations are underrepresented
11 in crystal structures and are often under reported due to resolution limits and
12 assumptions made during structure refinement [90, 91].

13

14 **Figure S3. Glycosylation of mammalian-expressed OMCP.** OMCP has two putative
15 N-linked glycosylation sites, N12 and N88 **(A)** WT, glycosylated OMCP migrates slowly
16 as a diffuse band. OMCP mutated at either N12 or N12/N88 migrate quickly as sharp
17 bands. Mutation of N88 alone does not effect OMCP migration. **(B)** N-linked glycans
18 were completely removed from 1 ug of OMCP by incubation with 10U Endo-F in 50mM
19 sodium phosphate (pH7.5) at 37C for 16h.

20

21

Table 1: Data collection and refinement statistics^a

	OMCP _{BR} -hNKG2D
Data collection	
Space group	P2 ₁
Cell dimensions	
a, b, c (Å)	43.3, 101.1, 91.4
$\alpha\beta\gamma$ (°)	90.0, 91.6, 90.0
Resolution (Å)	50-2.0 (2.07-2.00)
R_{sym}	11.8 (48.5)
I / σ	14.5 (3.8)

Completeness (%)	93.5 (91.5)
Redundancy	6.2 (5.3)
Refinement	
Resolution (Å)	44-2.0
Total reflections	309693
Unique reflection	50139
R_{work}	16.6% (21.0%)
R_{free}	21.4% (29.5%)
Wilson B-factor	21.62
Protein residues	791
Water molecules	524
R.M.S. deviations	
Bond lengths (Å)	0.003
Bond angles (°)	0.79

^a As defined by PHENIX [75]


Table 2: Interface contacts between NKG2D and OMCP

NKG2D-A	OMCP	Bond type
Lys150	Asp132	Salt bridge
Lys150	Trp127	H bond
Lys150	Trp127	Φ (3)
Ser151	Lys126	H bond
Ser151	Trp127	Φ (1)
Tyr152	Phe122	H bond
Tyr152	Phe122	Φ (9)
Tyr152	Lys126	Φ (5)
Met184	Thr118	H bond
Met184	Thr119	Φ (1)
Met184	Phe122	Φ (5)
Gln185	Arg66	Φ (1)
Leu191	Phe122	Φ (1)
Tyr199	Phe122	Φ (4)
Glu201	Arg66	Salt bridge
Thr205	Arg66	H bond
NKG2D-B	OMCP	Bond type


Leu148	Trp127	Φ (1)
Ser151	Glu131	H bond
Tyr152	Asp132	H bond
Tyr152	Glu131	Φ (3)
Tyr152	Met135	Φ (5)
Ile182	Ile49	Φ (2)
Glu183	Arg142	Salt bridge
Met184	Met135	Φ (1)
Met184	Arg138	Φ (2)
Met184	Arg142	H bond
Lys186	Arg142	Φ (1)
Leu191	Met135	Φ (1)
Glu201	Arg138	Salt bridge

1
2
3
4
5
6
7
8


Hydrogen bonds (H bonds), salt bridges and carbon-to-carbon hydrophobic interactions (Φ) are shown for each contact residue. The number of hydrophobic interactions between contact residues is designated in parenthesis.




A




B





A





B

















Supplemental Figure 1





A



1: OMCP
2: Mock
3: (N12Q) OMCP
4: (N88Q) OMCP
5: (N12Q/N88Q) OMCP

B

