

1 Breeding behaviour and productivity of Black-necked crane (*Grus nigricollis*) in Ladakh,

2 Indian Trans-Himalaya

3 Pankaj Chandan¹, Tanveer Ahmed², Afifullah Khan^{2*}

4 ¹WWF-India, 172-B, Lodhi Estate, New Delhi, India-110003

5 ²Department of Wildlife Sciences, Aligarh Muslim University, Aligarh, India- 202002

6 *Corresponding author:

7 Email: afifkhan@gmail.com (AF)

8

9 **Abstract**

10 A long-term study was conducted to understand some aspects of breeding biology of
11 Black-necked crane (*Grus nigricollis*) in Changthang, Ladakh. Data on aspects such as
12 the breeding season, courtship, mating, egg laying and incubation period, nest site
13 fidelity, egg morphometry, breeding productivity and recruitment rate were collected
14 between 2003 and 2012. Black-necked crane started arriving from last week of March
15 to first half of April and showed fidelity at ten nesting sites. Courtship and mating
16 peaked early morning (0700 hours), around noon (1100 hours) and in late evening
17 (0600 hours) while the nest building at evening (1600 hours). The egg laying period
18 initiated in May and extended up to July. The average incubation period was 33.88 ± 0.3
19 days. Hatching success, nest survival rate and fledgling increment rate of the bird were
20 73.3%, 0.55 ± 0.03 and 0.41 ± 0.02 respectively with an overall breeding productivity of
21 0.73. The present population of Black-necked crane in Changthang, Ladakh seems to
22 stable with an average recruitment rate of 15.7 ± 1.4 .

23 **Introduction**

24 The breeding of a species is largely influenced by the availability of food, water and
25 temperature in boreal and temperate zones, while intensity of illumination and micro-
26 habitat conditions apart from other demographic factors play an important role in tropics.
27 The date at which each individual actually lays eggs is related to the food supply at the
28 time of laying [1]. Laying cannot begin until food has become sufficiently abundant for
29 the females needed for general body maintenance and for feeding their chicks. Most

30 bird species breed around the time when food supplies are readily available and climatic
31 conditions are favourable. In deserts, or in relatively well watered areas, the sexual
32 cycles of vertebrates may be inhibited by drought and stimulated to sudden activity by
33 rainfall [2].

34 For water birds like cranes, factors reported to influence nest success include, water
35 depth at the nest [3-5], vegetation type around the nest [6-9], nest concealment [6], and
36 land use practices [5]. A few studies have suggested influence of weather (e.g. annual
37 precipitation) on the reproductive success [7]. The land use practices were reported to
38 affect nest success [10] while predation is considered as one of the most important
39 factors responsible for nest loss [11].

40 In Trans-Himalayan landscape, slowly melting glaciers and frozen wetlands mark the
41 beginning of the biological activity for various water bird species including Black-necked
42 cranes *Grus nigricollis*. However, available information on the breeding biology of black-
43 necked crane in the Indian Changthang is scanty and largely based on convenience
44 sampling, which is due to extreme and harsh climate of the area and limited funds and
45 logistics. Except few long term studies [12-13], studies conducted so far on black-
46 necked crane during the breeding season are mainly confined to explanatory notes on
47 status at few sites and effects of dogs on the nesting birds [14-19]. Considering the
48 paucity of scientific information on the breeding biology of black-necked crane, a long-
49 term study to understand and collect information on aspects such as breeding season,
50 courtship and display, nest site fidelity, egg laying, incubation and breeding productivity
51 was conducted. There was also a need to ascertain recruitment rate of black-necked

52 crane and factors influencing it, so that effective conservation and management strategy
53 for long-term survival of the species can be suggested.

54 Materials and Methods

55 Study Area

56 Changthang Cold Desert Wildlife Sanctuary; henceforth Changthang (32°25'-34°35' N &
57 77°30'-79°29' E) declared in 1987 is located in the eastern part of Ladakh, Jammu and
58 Kashmir state of India (Fig. 1). At an elevation between 3600 and 7000 m amsl.,
59 Changthang encompasses an area of about 22,588 km². The Zanskar mountain Range
60 forms its south-western border and the Karakoram Range the north-west. To the east, it
61 is contiguous with Tibetan portion of Changthang, the westernmost extension of Tibetan
62 plateau. The terrain is extremely rugged with a high proportion of open sandy plains,
63 cliffs and exposed rocks. After Taklang La, the study area opens into an extensive
64 plateau. The soil (pH ranges between 7 and 11), is sandy clay and to a lesser extent,
65 loam or loamy clay [20-21], with poor organic matter and nitrogen content [22]. Extreme
66 cold and arid conditions prevail in the area with monthly average maximum temperature
67 during winters often remain less than 100 C while in summer it goes up as high as 270
68 C. The monthly average minimum temperature during winters fluctuates between sub-
69 zero and -14 degree Celsius. Most of the precipitation is in the form of snowfall, and the
70 average annual rainfall is about <100 mm.

71 Changthang is virtually treeless, except for small isolated patches of *Myricaria elegans*
72 and *Yricaria germanica* in addition to poplar (*Populus* spp.) and willow (*Salix* spp.)

73 plantations at few places near human habitations. The major vegetation communities
74 include *Caragana-Eurotia*, *Artemisia-Tanacetum*, *Stipa-Oxytropis-Alyssum* and *Carex*
75 *melanantha-Leymus sccalinus*. At the higher altitudes (5000 m and above) sparse fell-
76 field communities with moss or cushion-like growth forms, e.g. *Thylacospermum*
77 *caespitosum*, *Arenaria bryophylla*, *Androsace saementosa* and lichens occupy the area.
78 Changthang encompasses numerous brackish and freshwater wetlands; as small as
79 one hectare to as large as 120 km². Stream banks and marsh meadows around the
80 wetlands exhibit characteristic sedge-dominated vegetation represented by species like
81 *Carex*, *Kobresia*, *Scirpus*, *Triglochin*, *Pucciniella*, *Ranunculus* and *Polygonum* (Rawat &
82 Adhikari, 2005). The shallow parts of the wetlands support a dense growth of aquatic
83 plants such as *Hippuris vulgaris*, *Potamogeton pectinatus*, *Potamogeton perfoliatus*,
84 *Zannichellia palustris*, and *Ranunculus natans*. These wetlands have been identified as
85 the breeding grounds of several migratory and resident avian species including Black-
86 necked Crane [18, 19, 23, 24].

87 **Fig 1. Location of Changthang cold desert Sanctuary in Ladakh, Jammu &
88 Kashmir**

89 **Methodology**

90 To record breeding status of black-necked cranes' population in Changthang, surveys
91 were initiated in 2000, when two pairs of breeding cranes were seen, each at Tsomoriri
92 and Tsokar basin. In subsequent years (2001 and 2002), extensive surveys were
93 conducted covering large portion of Changthang plateau. All previously described sites
94 were visited and occurrence of cranes was recorded at 21 different locations within

95 Changthang. Once the distribution of cranes was understood, systematic surveys were
96 initiated between 2003 and 2012 to collect information on various aspects of breeding
97 biology.

98 Information on arrival and departure of Black-necked crane was collected from four
99 different locations viz. Staklung, Chushul, Tsokar and Yaya Tso, since collection of data
100 from all sites was not possible on weekly basis as the sites were widely distributed over
101 an area of more than 20,000 Km². These four sites, however, represent altitudinal and
102 habitat variations between different breeding territories of Black-necked crane in
103 Changthang. The initial and last sighting of Black-necked crane at any breeding ground
104 was considered as arrival and departure dates respectively. Egg laying dates along with
105 clutch size was also recorded each year. In addition to this information from locals and
106 herders in the nearby villages around the nesting sites were also collected to ascertain
107 the dates. Measurements of confirmed infertile eggs (34 eggs) were taken using a
108 calliper.

109 Focal-animal sampling method (Altmann 1974) was used to determine the time budget
110 of Black-necked crane during breeding season. Occurrence of breeding related
111 behaviour such as courtship, mating and nesting were recorded at every five minute
112 interval through a movable hide from morning (0600 hours) to late evening (1800
113 hours).

114 Data Analysis

115 The egg volume was estimated using the following formula as described by (Hoyt
116 1979). Nesting success, hatching success, nestling survival rate, fledgling increment

117 rate and overall breeding productivity were calculated to determine breeding success at
118 various stages of breeding Black-necked cranes using the following formulae:

119 Egg volume = 0.51 (Length X Breadth²)

120 Nest success =
$$\frac{\text{Total No. of active nests}}{\text{Total nests}} * 100$$

121 Hatching success =
$$\frac{\text{Total No. of eggs hatched}}{\text{Total eggs}} * 100$$

122 Nestling survival rate =
$$\frac{\text{Nestlings survived and reached the fledgling stage}}{\text{Number of eggs hatched}}$$

123 Fledgling increment rate =
$$\frac{\text{Number of fledglings}}{\text{Number of eggs laid}}$$

124 Breeding productivity =
$$\frac{\text{Number of chicks survived and fledged}}{\text{Breeding pair}}$$

125 Recruitment rate =
$$\frac{\text{Number of chicks survived and fledged}}{\text{Total population in the study area}} * 100$$

126 Data on courtship, mating and nest building were analysed by summarizing frequency
127 of occurrences on hourly basis and percentages were depicted to understand the
128 activity rhythm. Differences in mean incubation period across different years and sites
129 were analyzed using Kruskal-Wallis One Way ANOVA. One sample t-test was used to
130 check the differences in mean hatching success and nestling survival rates. All
131 statistical analysis was conducted in SPSS 16.0.

132 **Results**

133 **Breeding behaviour of Black-necked crane**

134 The arrival period of Black-necked crane ranged from last week of March to first week of
135 May and peaked during first half of April in Changthang, (Table S1). The courtship and
136 mating activity started within two to three days of arrival of cranes in their respective
137 breeding territories. Courtship and mating peaked early morning (0700 hours), around
138 noon (1100 hours) and late evening (0600 hours) (Fig. 2). Nest building activity
139 remained at its low in the morning hours and increased between 1000 and 1200 hours
140 and peaked at evening (1600 hours) (Fig. 2). The species showed site fidelity at 10
141 nesting sites while at three sites, nesting occurred at different places. At the rest five
142 sites, nesting did not occur regularly (Table S2).

143 **Figure 2: Diurnal rhythm of courtship- mating and nest building activity in
144 Changthang, Ladakh**

145 A total of 146 nests of Black-necked crane were observed at 23 locations; 90.42% of
146 nests were active while the rest (9.58%) nests were inactive. Of the total active nests,
147 76.03% were successful in raising at least one chick, while 23.97% of nests failed in
148 raising any chick. The maximum nesting success (ca 86%) was recorded in 2005 and
149 2008 and a minimum (64.29%) was in 2012. The nesting success varied over different
150 years of the study period (Fig. 3).

151 Most of the crane pairs laid their eggs during June (77.27%), nearly one fourth during
152 May (19.09%) and few (3.64%) during July. The clutch size in most of the monitored
153 nests (16 nesting sites) was two except one nesting site where single egg was recorded
154 in 2005. The eggs of Black-necked crane had an average length breadth, weight and

155 volume of 107.32 ± 1.02 mm, 63.17 ± 0.33 mm, 210.85 ± 3.17 g and 218.95 ± 3.67 cm³
156 respectively (Table S3). The length, breadth, volume and weight of the eggs did not
157 differ significantly over the years. Incubation started immediately after laying of the first
158 egg with an average incubation period of 33.88 ± 0.3 days (range 30-38 days). The
159 maximum (36.00 ± 1.0 days) incubation period was recorded during 2012 whereas
160 minimum (32.33 ± 0.6 days) during 2010 (Fig. 3). The incubation period also did not
161 differ significantly across years and various nesting sites.

162 **Figure 4: Average incubation period during different years in Changthang,**

163 **Ladakh**

164 **Breeding productivity of Black-necked crane**

165 A total 266 eggs were laid during the study period, of which, 195 eggs (73.3%) were
166 successfully hatched. The hatching success was maximum (92.9%) during 2008 and
167 minimum (59.3%) during 2009 (Table 1). The hatching success of Black-necked crane
168 differed significantly between different years of study (KW = 18.83, df = 9, p<0.01). Of
169 the total eggs successfully hatched and produced nestlings, 108 chicks (56%) reached
170 the fledgling stage. Nestling survival rate ranged between 0.44 and 0.68 with a mean
171 value of 0.56 ± 0.03 (Table 1). The overall mean fledgling increment rate was $0.41 \pm$
172 0.03 (range 0.27-0.55). Overall breeding productivity of Black-necked crane was 0.74
173 while it was maximum during 2010 (0.92) and lowest (0.50) in 2011 (Table 1). The
174 mean recruitment rate was $15.70 \pm 1.4\%$ and it ranged between 7.91% and 22.03%
175 during 2012 and 2006 respectively (Table 1).

176

177

178 Table 1: Breeding performance of Black-necked crane in Changthang, Ladakh

Year	Total cranes	Breeding pairs	Hatching Success (%)	Nestling Success rate	Fledgling Success rate	Breeding Productivity (%)	Recruitment rate (%)
2003	60	16	73.3	0.45	0.33	0.63	16.67
2004	64	15	70.4	0.68	0.48	0.87	20.31
2005	58	15	76.7	0.43	0.33	0.67	17.24
2006	59	15	74.1	0.65	0.48	0.87	22.03
2007	58	16	66.7	0.61	0.41	0.69	18.97
2008	81	15	92.9	0.50	0.46	0.87	16.05
2009	65	15	59.3	0.56	0.33	0.60	13.85
2010	73	12	70.8	0.65	0.46	0.92	15.07
2011	79	14	61.5	0.44	0.27	0.50	8.86
2012	139	14	90.0	0.61	0.55	0.79	7.91
Overall			73.6±3.4	0.56±0.03	0.41±0.03	0.74	15.70±1.4

179

180 **Discussion**

181 For many species of cranes, one of the most crucial unresolved questions related to
182 breeding biology is one that concerns with breeding productivity over a longer period of
183 time. An often asked research question while studying the breeding ecology of a
184 species is to know the reproductive success of a nesting bird. Many consider clutch
185 size, nesting success, and nesting mortality as critical site specific information required
186 for long term conservation of the species. To bridge this information gap the present
187 study has focused on some of these issues.

188 The Black-necked cranes start arriving in Ladakh during last week of March to the first
189 week of May with majority arriving in the first half of April. Their departure starts during
190 last week of October and continues until first week of November. However, Pfister [17]
191 reported the arrival of Black-necked cranes on their breeding ground from late April to
192 early May and departure from mid-October to November. The differences in the arrival
193 and departure may be attributed to short term observations of Pfister [17], restricted
194 only to one breeding season. Consistent with the present study, Dehao [25], Dwyer *et*
195 *al.* [26] and Bishop [27] reported the arrival of Black-necked cranes on their breeding
196 grounds in China from late March to mid-May and departure from mid-October to
197 November. This similarity in the arrival and departure of cranes in India and China is
198 due to the fact that Ladakh is part of the same eco-region. It is pertinent to note the late
199 arrival of Black-necked cranes at Yaya Tso. Since Yaya Tso is the highest breeding
200 place within Changthang, where the wetland remained frozen till about late April and
201 hence crane arrive during last week of April and first week of May when melting of snow
202 starts and weather is not so harsh.

203 The arrival was followed by courtship and mating which was frequently performed
204 during the month of April and May. In most of the cases the courtship and mating was
205 recorded during early morning, at about mid-day and in the evening. This is similar to
206 the observation made during various other studies conducted on the breeding behaviour
207 of Black-necked crane in China [26, 28-29].

208 The clutch size of most crane species is consistently two except grey-crowned and
209 black-crowned cranes [30]. These two species of crane usually lay more than two eggs.
210 The clutch of grey-crowned crane may even sometime consist of five eggs. However,

211 such a variation has neither been observed nor been reported from other breeding
212 ranges of Black-necked crane. Larger clutch size of grey-crowned and black-crowned
213 canes may be due to environmental conditions. Since these species inhabit tropical
214 areas therefore they require less energy in maintaining body temperature as compared
215 to Black-necked crane inhabiting alpine area. Furthermore, the former are resident while
216 the latter is migratory. Most migratory species have small clutch size as they have to
217 breed and rear chicks in relatively shorter span of time, before migration to their
218 wintering areas.

219 Mean egg weight of Black-necked crane (210 g) observed during the present study was
220 comparable as reported by Dehao *et al.* [25]. However, the mean egg weight during
221 present study was of infertile eggs. We did not collect data from the active nests to
222 avoid disturbance to the breeding pair and possible harm, if any, to the eggs. The
223 incubation period of 30-38 days during present study was little more as reported by
224 Dehao *et al.* [25] and Pfister [17]. The incubation period reported by the former was
225 based on observation of three nests while the latter observed only one nest. During
226 present study the sample size was large and therefore the variation was also large.

227 Hatching success of 27 avian species in the subarctic region ranged between 28 and
228 100 % and the major causes of hatching failure were attributed to weather conditions,
229 predation and poaching [31]. The avian species with small clutch size had higher
230 percentage of hatching success as compared to those had large clutch size [32]. Since
231 most species of cranes have small clutch size it is therefore expected to have higher
232 hatching success as compared to those avian species having large clutch size. The
233 assertion seems to be true in case of crane species. The hatching success of at least

234 10 crane species other than Black-necked crane ranged between 56 and 93%; the
235 lowest was reported for Grey crowned crane *Balearica regulorum* [33] and the highest
236 was in Red-crowned crane *Grus japonensis* [34]. The hatching success observed
237 during the current study is in conformity with that of the hatching species observed in
238 other crane species. Predation, flooding, abandonment and egg infertility have been
239 reported as major causes of hatching failure [26]. During present study average
240 hatching success was 73.6%. The reasons of failure include depredation that accounted
241 only about six per cent while two nests containing four eggs (about 1.3%) were washed
242 away due to flood. Rest 20 % losses were for unknown reasons as it was not practically
243 possible to continuously monitor all the nests, spread over such a large area as
244 Changthang. Low hatching success in certain years could be attributed to weather
245 condition.

246 High mortality of nestlings is probably a characteristic of majority of avian species and
247 Black-necked crane is not an exception. Other species of cranes also show similar
248 pattern of nestling survival. For instance, the fledgling percentage of conspecific Grey-
249 crowned crane was 60 [33], Red-crowned crane 56 [35], Demoiselle crane *Grus virgo*,
250 63 [36] and it was 65 % for Florida sandhill crane *Antigone canadensis pratensis* [37].
251 The main cause of fledgling mortality in Changthang was depredation as nearly 25%
252 fledglings were depredated by land and avian predators. The cause of mortality of rest
253 20% remained unknown.

254 A perusal of data on breeding productivity among different species of cranes revealed
255 that it ranged between 0.13 and 1.05. The productivity of wattled crane *Bugeranus*
256 *carunculatus* in Zambia was reported 0.13 [38]. It was 0.87 for common crane *Grus grus*

257 [39], and 0.70 for sandhill crane *Antigone canadensis* [37]. The average breeding
258 productivity (0.73) of Black-necked crane is more or less similar to other crane species.
259 Of importance is the recruitment rate of a species in order to understand the status of
260 population. During the present study, lowest recruitment rate observed during 2012 was
261 due to sudden increase in wintering population of Black-necked crane in Changthang.
262 The reasons of sudden surge in crane numbers were not known. Similar pattern was
263 also observed in 2008 when the total population of wintering crane had increased from
264 previous year's 58 to 81. During 2011, the low recruitment is attributed to the egg
265 predation and chick mortality. However, in rest of the years of study (excluding 2008
266 and 2012) the recruitment rate remained above 13%. Lovvorn and Kirkpatrick [40]
267 opined that a stable breeding population of Sandhill cranes requires a recruitment rate
268 of 10 to 12%. If the same criterion is applied, the present population of Black-necked
269 crane in Changthang can be considered as stable.

270 **Acknowledgements**

271 We are grateful to WWF Netherlands for long term funding support for this project.
272 Thanks are due to Mr. Jigmet Takpa, CCF & Regional Wildlife Warden, Ladakh for
273 granting permission to work in the area. A very special thanks to Mr. Ravi Singh,
274 George Archibald, Ms. Esther Blom, Dr. Taej Mundukur, Mr. A. K. Singh, Dr. Sejal
275 Worah, Dr. V. B. Mathur, Dr. Asad R. Rahmani, Maj. Gen. G. D. Bakshi, Mr. A. K.
276 Srivastava, Dr. Parikshit Gautam, Dr. C. M. Seth, Ms. Archana Chatterjee, Dr. Dipankar
277 Ghose, Late Parkash Gole, Col. R. T. Chacko, Prof. Jamal A. Khan, Dr. Gopi Sundar,
278 Mr. Otto Pfister, Mr. Kiran Rajashekriah, Mr. Saleem-ul- Haq, Mr. Tsering Angchuk and
279 Mr. Intesar Sohail for their help. We are also thankful to Dr. Li Fengshan, Mr. Blaise

280 Humbert-Droz, Mrs. Usha Ganguli-Lachungpa, Mr. Shakeel Ahmed, Kamal Mehdi,
281 Pushpendra Singh Jamwal, Rohit Ratan, Nisha Khatoon, Tsewang Rigzin, Anupam
282 Anand, Pijush Kr. Dutta, Partha S. Ghose, Priyadarshinee Shrestha, Lak Tsheden
283 Theengh, Mr. Ram Saroop and Mr. Dawa Tsering for their help and assistance.

284 **References**

- 285 1. Perrins CM. The timing of bird's breeding season. *Ibis*. 1970; 112: 242-255.
- 286 2. Keast JA and Marshall AJ. The influence of drought and rainfall on reproduction in
287 Australian desert birds. *Proc. Zool. Soc. London*. 1954; 124: 493-499.
- 288 3. Littlefield CD. Sandhill Crane nesting habitat, egg predators, and predator history on
289 Malheur National Wildlife Refuge, Oregon. *Northwestern Nat.* 1995; 76: 137-143.
- 290 4. Maxson SJ, Riggs MR. Nest habitat selection and nest success of greater Sandhill
291 Cranes in northwestern Minnesota. Unpublished report, Minnesota Department of
292 Natural Resources, St. Paul, MN. 1996. Available from:
293 https://www.researchgate.net/publication/266034453_NEST_HABITAT_SELECTION_AND_NEST_SUCCESS_OF_GREATER_SANDHILL_CRANES_IN_NORTHWESTERN_MINNESOTA
- 296 5. Littlefield CD, Cornely JE, Ehlers BD. Effects of an early spring burn on greater
297 Sandhill Crane nesting success at Malheur National Wildlife Refuge, Oregon.
298 *Proceedings North American Crane Workshop* 2001; 8: 45-47.

- 299 6. Littlefield CD, Ryder RA. Breeding biology of the Greater Sandhill Crane on Malheur
300 National Wildlife Refuge, Oregon. *Transactions of the North American Wildlife and*
301 *Natural Resources Conference*. 1968; 33: 444-454.
- 302 7. Drewien RC, Brown, WM, Kendall WL. Recruitment in Rocky Mountain Greater
303 Sandhill Cranes and comparisons with other crane populations. *J. Wildl. Manage.*
304 1995; 59: 339–356.
- 305 8. Stern M A, Pampush GJ, Delcarlo R E. Nesting ecology and productivity of greater
306 Sandhill Cranes at Sycan Marsh, Oregon. In: Lewis J C, editor, *Proceedings of the*
307 *1985 Crane Workshop* Platte River Whooping Crane Maintenance Trust, Grand
308 Island, Nebraska; 1987. pp. 249-256.
- 309 9. Urbanek RP, Bookhout TA. Nesting of Greater Sandhill Cranes on Seney National
310 Wildlife Refuge. In: Wood DA, editor. *Proceedings of 1988 North American Crane*
311 *Workshop*, Florida Game and Freshwater Fish Commission, Nongame Wildlife
312 Program Technical Report No. 12; 1992. Pp. 161-172.
- 313 10. Littlefield CD, Paullin DG. Effects of land management on nesting success of Sandhill
314 Cranes in Oregon. *Wildl. Soc. Bull.* 1990; 18: 63-65.
- 315 11. Sargeant AB, Arnold PM. Predator management for ducks on waterfowl production
316 areas in the northern plains. In: Clark DO, editor, *Proceedings Eleventh Vertebrate Pest*
317 *Conference*, University of California, Davis, California; 1984. pp.161-167.

- 318 12. Khan A, Chandan P, Takpa J, Hussain SA, Rattan R, Jamwal PS, Ahmed T. Diurnal
319 time budget of breeding Black-necked Crane (*Grus nigricollis*) in Changthang,
320 Ladakh, India. *Zoological Research* 2014; 35(S1): 158–166.
- 321 13. Chandan P, Khan A, Takpa J, Hussain SA, Mehdi K, Jamwal PS, Rattan R, Khatoon
322 N, Rigzin T, Anand A, Dutta PK, Ahmed T, Ghose PS, Shrestha P, Theengh LT.
323 Status and distribution of Black-necked Crane (*Grus nigricollis*) in India. *Zoological
324 Research* 2014; 35 (S1): 39–50.
- 325 14. Ali S. Status and Conservation of the Black-necked Crane (*Grus nigricollis*) and
326 other wildlife in Ladakh. Report submitted to World Wildlife Fund; 1976.
- 327 15. Gole P. Black-necked Cranes in Ladakh. In: Lewis JC, Masatomi H, editors. *Crane
328 research around the world: proceedings of the International Crane Symposium at
329 Sapporo, Japan in 1980 and papers from the World Working Group on Cranes,*
330 International Council for Bird Preservation. Baraboo, USA: International Crane
331 Foundation; 1981. pp. 197–203.
- 332 16. Hussain SA. Status of Black-necked Crane in Ladakh – 1983, Problems and
333 Prospects. *J. Bombay Nat. Hist. Soc.* 1985, 82: 449-58.
- 334 17. Pfister, O. The Breeding Ecology and Conservation of the Black-necked Crane (*Grus
335 nigricollis*) in Ladakh India. M.Sc. Thesis, University of Hull, Hull. 1998.
- 336 18. Chandan P, Gautam P, Chatterjee A. Nesting sites and breeding success of Black-
337 necked Crane (*Grus nigricollis*) in Ladakh, India. In: Boere GC, Galbraith CA, Stroud

- 338 DA editors. *Waterbirds around the world*, The Stationery office, Edinburgh, UK; 2005.
- 339 pp. 311-314.
- 340 19. Ahmed T, Chandan P, Khan A. Observations on some nesting birds of the Tso-Kar
341 Basin, Ladakh. *Indian BIRDS* 2019; 15 (1): 13–16.
- 342 20. Bhat GM. The soils of Kashmir. Agriculture Bulletin, Agriculture Department, J & K
343 Government; 1965.
- 344 21. Kachroo P, Sapru BL, Dhar U. Flora of Ladakh: An ecological and Taxonomic
345 Appraisal. Bishen Singh and Mahendrapal Singh, Dehradun, India; 1977.
- 346 22. Murti SK. Flora of Cold Deserts of Western Himalaya. Botanical Survey of India,
347 Calcutta; 2001.
- 348 23. Hussain SA, Singh RK, Badola R. An ecological survey of the Trans-Himalayan
349 wetlands of the proposed Changthang Biosphere Reserve, India, for conservation
350 planning. *Biosphere Conservation* 2008; 9(1): 53-63.
- 351 24. Ahmed T, Khan A, Chandan P. Status of Avifauna of Tsokar basin, Ladakh, Indian
352 Trans-Himalaya. In: Serajuddin M, editor. Biodiversity, Environment and Functional
353 Biology. Excellent Publishing House, New Delhi; 2014. pp. 116-124.
- 354 25. Dehao L. The distribution and status of the Black-necked Crane on the Tibetan
355 Plateau. In: Archibald GW and Pasquier RF. Editors. Proceedings of the 1983
356 International Crane Workshop, Bharatpur, India. Baraboo, Wisconsin, USA:
357 International Crane Foundation; 1987. pp. 45–49
- 358 26. Dwyer NC, Tanner GW. Nesting success in Florida Sandhill Crane. *Wilson Bull.*
359 1992; 104(1): 22-31.

- 360 27. Bishop MA. Report on Investigations of Status, Distribution and Winter Habitat Use
361 by Black-necked Cranes in South-central Tibet, 1990-1994, ICF Baraboo; 1996.
- 362 28. Dehao, L, Zhijun Z, Zhikang W, Zhumei L, Youhui W. On the structure and behavior
363 of the breeding population of Black-necked Cranes in Songpan Meadow in Sichuan
364 Province. In: Harris J, editor. Proceedings 1987 International Crane Workshop.
365 Qiqihar, Heilongjiang Province, People's Republic of China. Baraboo: International
366 Crane Foundation; 1991. pp. 83–87.
- 367 29. Yang R, Wu HQ, Yang XJ, Jiang WG, Zou L, L Xiang. Diurnal Time Budget of the
368 Black-necked Crane During the Breeding Season. *Waterbirds* 2007; 30(1): 80-85.
- 369 30. Allan DG. A review of the biology and conservation status of cranes. In: Beifuss RD,
370 Tarboton WR, Nathan NG editor. Proceedings African Crane and Wetland Training
371 Workshop. 1998; pp. 13-51.
- 372 31. Joseph R, Jehl Jr. Patterns of Hatching Success in Subarctic Birds. *Ecology*. 1971;
373 52 (1): 169-173.
- 374 32. Reid JM, Monaghan P, Ruxton GD. The consequences of clutch size for incubation
375 conditions and hatching success in starlings. *Funct. Ecol.* 2000; 14(5): 560-565.
- 376 33. Pomeroy DE. Aspects of the ecology of Crowned Cranes *Balearica regulorum* in
377 Uganda. *Scopus* 1980; 4: 29-35.
- 378 34. Ma, YC. The annual cycle of red-crowned crane. In: Lewis JC, Masatomi H, editors.
379 Crane research around the world: Proceedings of the International Crane Symposium

- 380 at Sapporo, Japan in 1980 and paper form the Working Group on crane, International
381 Council for Birds Preservation, Baraboo, Wisconsin; 1981. pp: 94-95
- 382 35. Masatomi H. The red crowned crane. In: Lewis JC, Masatomi H, editors. Crane
383 research around the world: Proceedings of the International Crane Symposium at
384 Sapporo, Japan in 1980 and paper form the Working Group on crane, International
385 Council for Birds Preservation, Sapporo. Japan; 1981. pp.81-85.
- 386 36. Winter SW. The Demoiselle Crane in the Agricultural Landscape of the Ukrainian Steppe
387 Zone. In: Harris J, editor. Proceedings of the 1987 International Crane Workshop.
388 International Crane Foundation, Baraboo, Wisconsin, USA; 1991. pp. 285-295.
- 389 37. Nesbitt SA. First reproductive success and individuals productivity in Sandhill crane.
390 *J. Wildl. Manage.* 2001; 56(3): 573-577.
- 391 38. Konard PM. Status and Ecology of Wattled Cranes in Africa. In: Lewis JC, Masatomi
392 H editor. Crane Research Around the World: Proceedings of the International Crane
393 Symposium at Sapporo, Japan in 1980. International Crane Foundation, Baraboo,
394 Wisconsin, USA. 1981; pp. 220-237.
- 395 39. Neumann T. Breeding Status of the Common Crane in the Federal Republic of
396 Germany. In: Archibald GW, Pasquier RF. editors. Proceedings of the 1983
397 International Crane Workshop. International Crane Foundation, Baraboo, Wisconsin,
398 USA. 1987; Pp. 243-245.
- 399 40. Lovvorn JR, Kirkpatrick CM. Recruitment and socially-specific flocking tendencies of
400 eastern Sandhill Crane. *Wilson Bull.* 1982; 94: 313-321.

401 **Supporting Information**

402 Table S1: Arrival and departure dates of Black-necked crane at four study sites during
403 various years in Changthang, Ladakh

404 Table S2: Nest sites fidelity at different breeding sites at Changthang, Ladakh

405 Table S3: Eggs morphometry of Black-necked crane in Changthang, Ladakh

406 **Keywords:** Breeding, Crane, Fledgling, Hatching, Nestling, productivity, Changthang,
407 Ladakh




Figure 1: Location of Changthang Cold Desert Sanctuary in Jammu & Kashmir, India





Figure 2: Diurnal rhythm of courtship- mating and nest building activity in Changthang, Ladakh



bioRxiv preprint doi: <https://doi.org/10.1101/809095>; this version posted October 17, 2019. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY 4.0 International license.

Figure 3: Percentage of successful and failed nesting during study period in Changthang, Ladakh




Figure 4: Average incubation period during different years in Changthang, Ladakh