

A trafficome-wide RNAi screen reveals deployment of early and late secretory host proteins and the entire late endo-/lysosomal vesicle fusion machinery by intracellular *Salmonella*

Alexander Kehl^{1,4}, Vera Göser¹, Tatjana Reuter¹, Viktoria Liss¹, Maximilian Franke¹,
Christopher John¹, Christian P. Richter², Jörg Deiwick¹ and Michael Hensel¹.

¹Division of Microbiology, University of Osnabrück, Osnabrück, Germany; ²Division of Biophysics, University of Osnabrück, Osnabrück, Germany; ³CellNanOs – Center for Cellular Nanoanalytics, Fachbereich Biologie/Chemie, Universität Osnabrück, Osnabrück, Germany; ⁴current address: Institute for Hygiene, University of Münster, Münster, Germany

Running title: Host factors for SIF formation

Keywords: siRNA knockdown, live cell imaging, *Salmonella*-containing vacuole, *Salmonella*-induced filaments

Address for correspondence:

Alexander Kehl

Institute for Hygiene

University of Münster

Robert-Koch-Str. 4148149 Münster, Germany

Tel.: +49(0)251/83-55233

E-mail: alexander.kehl@ukmuenster.de

or

18.11.2019

Host factors for SIF formation

26

27 Michael Hensel

28 Abteilung Mikrobiologie

29 CellNanOs – Center for Cellular Nanoanalytics Osnabrück

30 Fachbereich Biologie/Chemie, Universität Osnabrück

31 Barbarastr. 11

32 49076 Osnabrück, Germany

33 Tel: ++ 49 (0)541 969 3940

34 Fax: ++ 49 (0)541 969 3942

35 E-mail: Michael.Hensel@uni-osnabrueck.de

18.11.2019

Host factors for SIF formation

Abstract

The intracellular lifestyle of *Salmonella enterica* is characterized by the formation of a replication-permissive membrane-bound niche, the *Salmonella*-containing vacuole (SCV). A further consequence of the massive remodeling of the host cell endosomal system, intracellular *Salmonella* establish a unique network of various *Salmonella*-induced tubules (SIT). The bacterial repertoire of effector proteins required for the establishment for one type of these SIT, the *Salmonella*-induced filaments (SIF), is rather well-defined. However, the corresponding host cell proteins are still poorly understood. To identify host factors required for the formation of SCV and SIF, we performed a sub-genomic RNAi screen. The analyses comprised high-resolution live cell imaging to score effects on SIF induction, dynamics and morphology. The hits of our functional RNAi screen comprise: i) The late endo-/lysosomal SNARE (soluble N-ethylmaleimide-sensitive factor attachment protein receptor) complex, consisting of STX7, STX8, VTI1B, and VAMP7 or VAMP8, this is, in conjunction with RAB7 and the homotypic fusion and protein sorting (HOPS) tethering complex, a complete vesicle fusion machinery. ii) Novel interactions with the early secretory GTPases RAB1A and RAB1B, possibly providing a link to coat protein complex I (COPI) vesicles and reinforcing recently identified ties to the endoplasmic reticulum. iii) New connections to the late secretory pathway and/or the recycling endosome via the GTPases RAB3A, RAB8A, and RAB8B and the SNAREs VAMP2, VAMP3, and VAMP4. iv) An unprecedented involvement of clathrin-coated structures. The resulting set of hits allowed to characterize completely new host factor interactions, and strengthen observations from several previous studies.

Author Summary

The facultative intracellular pathogen *Salmonella enterica* serovar Typhimurium induces the reorganization of the endosomal system of mammalian host cells. This activity is dependent on translocated effector proteins of the pathogen. The host cells factors required for endosomal

18.11.2019

Host factors for SIF formation

62 remodeling are only partially known. To identify such factors for formation and dynamics of
63 endosomal compartments in *Salmonella*-infected cell, we performed a live cell imaging-based
64 RNAi screen a to investigate the role of 496 mammalian proteins involved in cellular logistics.
65 We identified that endosomal remodeling by intracellular *Salmonella* dependent on host factors
66 in following functional classes: i) the late endo-/lysosomal SNARE (soluble *N*-ethylmaleimide-
67 sensitive factor attachment protein receptor) complex, ii) the early secretory pathway,
68 represented by regulators GTPases RAB1A and RAB1B, iii) the late secretory pathway and/or
69 recycling endosomes represented by GTPases RAB3A, RAB8A, RAB8B, and the SNAREs
70 VAMP2, VAMP3, and VAMP4, and iv) clathrin-coated structures. The identification of these
71 new host factors provides further evidence for the complex manipulation of host cell transport
72 functions by intracellular *Salmonella* and should enable detailed follow-up studies on the
73 mechanisms involved.

18.11.2019

Host factors for SIF formation

Introduction

The food-borne, facultative intracellular pathogen *Salmonella enterica* serovar Typhimurium (STM) is the etiological agent of gastroenteritis in humans or systemic infections in mice [1].

An early step in disease is the active invasion of epithelial cells. This process is dependent on the translocation of effector proteins by STM into the host cell through a type 3 secretion system (T3SS) encoded on *Salmonella* pathogenicity island 1 (SPI1) [2, 3].

After invasion STM, similar to many other intracellular pathogens, establish a replicative niche in host cells, termed *Salmonella*-containing vacuole (SCV). This process is dependent on the function of a distinct T3SS, encoded by SPI2 [4, 5] and translocating another set of effectors [6]. Though initially associating with markers of the early endosome (EE) such as EEA1 and the small GTPase RAB5 [7, 8], the SCV finally acquires several markers of the late endosome (LE). These include lysosome-associated membrane proteins (LAMPs) [9, 10], the vacuolar ATPase [11], and RAB7 [12, 13]. Concurrently, other canonical organelle markers such as the mannose-6-phosphate receptor are excluded [14].

A unique feature of STM among intravacuolar bacteria is the formation of a diverse array of long tubular structures, *Salmonella*-induced tubules (SIT) [15]. These include the LAMP-decorated *Salmonella*-induced filaments (SIF), the first SIT discovered [16, 17]. Moreover, SIF have been structurally characterized, revealing the presence of a double membrane tubular network [18, 19]. The host-derived membranes forming SCV, SIF, and other tubular compartments are collectively termed *Salmonella*-modified membranes (SMM).

The repertoire of bacterial effector proteins necessary for formation of SMM is quite well-characterized, with the SPI2-T3SS effector protein SifA being the most important factor [20, 21]. However, much less is known about corresponding host factors required for biogenesis of SMM. One crucial factor in SIF biogenesis is the SifA- and kinesin-interacting protein SKIP (a.k.a. PLEKHM2). In conjunction with the effectors SifA and PipB2 [22, 23] and the small

18.11.2019

Host factors for SIF formation

100 GTPase ARL8B [24, 25], SKIP mediates kinesin-1 interaction and thus a link to the microtubule
101 cytoskeleton and organelle motility [26].

102 Several attempts were made to analyze the interactions of STM with host factors in a systematic
103 manner. These comprise RNA interference (RNAi) screens aiming at different parts of the STM
104 infection process. Two genome-scale screens targeted the invasion [27, 28], while three screens
105 focused on intracellular replication with two sub-genomic screens covering kinases and
106 corresponding phosphatases, respectively [29, 30], and a genome-wide screen [31].

107 Additionally, two recent proteomic studies also shed light on interactions of intracellular STM
108 with host cells. Vorwerk et al. [32] characterized the proteome of late SMM, while Santos et al.
109 focused on early and maturing SCV [33].

110 All of these studies identified host factors yet unprecedented in STM pathobiology, and showed
111 the general value of such systematic approaches. However, none of these approaches targeted
112 specifically SIF, thus a host-SIF interactome is far from complete. Therefore, we established a
113 targeted RNAi screen comprising 496 human genes mostly involved in cellular logistics to
114 identify host factors involved in the formation of SIF. Using stably LAMP1-GFP-transfected
115 HeLa cells, we performed automated microscopy on a spinning disk confocal microscope
116 (SDCM) system with time-lapse live cell imaging (LCI) of STM infection, and scored for
117 altered SIF formation as phenotypic readout. Investigating high-scoring hits of the RNAi
118 screen, we validated several so far unknown host-SIF interactions by LCI: (i) involvement of
119 the late endo-/lysosomal soluble *N*-ethylmaleimide-sensitive factor attachment protein receptor
120 (SNARE) complex and its interaction partners, (ii) interactions of SIF with early secretory
121 RAB1A/B, (iii) late secretory RAB3A, RAB8A/B, and VAMP2/3/4, and (iv) a connection to
122 clathrin-coated structures.

18.11.2019

Host factors for SIF formation

Results

Setup and evaluation of RNAi screen

We aimed to identify host cell factors that are required for the endosomal remodeling induced by intracellular STM in a SPI2-T3SS-dependent manner. For this, an RNAi screen was performed with siRNAs predominantly targeting mammalian genes involved in cellular logistics and trafficking (75.2% categorized in intracellular transport according to Gene Ontology terms). This subset of 496 genes was termed ‘Trafficoome’ and is listed in Table S1. Such a screen necessitates specific considerations and controls with the major ones described below, and further experimental issue are detailed in Suppl. Materials.

As a phenotypic readout for STM-induced endosomal remodeling, we scored the formation of SIF in infected cells. SIF show a highly dynamic behavior in their early phase after formation, with constant elongation and retraction [34, 35]. Thus, in contrast to previous RNAi screens done by analyzing fixed cells, we decided to perform this screen by LCI in order to obtain maximal phenotypic information. A previously established HeLa cell line stably transfected with LAMP1-GFP as the marker for SIF [18] was used as host cell.

As controls for STM-induced phenotypes, we used STM wild type (WT), capable in SIF induction, and an isogenic strain defective in SsaV, a central component of the SPI2-T3SS, and thus unable to induce SIF formation (Figure 1A). As a control for successful reverse transfection in general, we analyzed the lethal effect of an siRNA directed against polo-like kinase 1 (PLK1), a cell cycle control protein. The knockdown of this protein leads initially to a cell cycle arrest, and ultimately to cell death, as shown in Figure 1B. Besides, a phenotype-related control was established, i.e. knockdown of a host factor already known as essential for SIF formation. A host factor directly involved in SIF formation is SKIP [22]. This study already successfully used SKIP silencing, thus we used an siRNA with the same sequence as control. Real-time PCR indicated that the siRNA targeting SKIP yielded not a complete but sufficient

18.11.2019

Host factors for SIF formation

and significant knockdown with a reduction to ca. 22% (Figure 1C). Transfection with AllStars siRNA did not affect SIF formation and dynamics over the course of infection (Figure 1D, Movie 1), while SKIP knockdown abolished SIF formation (Figure 1D, Movie 2) and reduced intracellular replication of STM. Though this does not completely exclude off-target effects, the phenotypic/visual control showed at least the intended purpose of this siRNA being fulfilled. The partial knockdown explains the rare appearance of SIF. Taken together, the establishment of the proper controls allowed us the execution of a larger scale RNAi screen.

The RNAi trafficome screen

The complete workflow of the RNAi screen executed is summarized in Figure 2. First, siRNAs were automatically spotted onto 96-well plates. Additionally, each plate contained siAllStars, siPLK1, and siSKIP as negative and positive siRNA controls, and as phenotype-specific control, respectively. HeLa-LAMP1-GFP cells were seeded onto siRNAs for reverse transfection, incubated for 72 h, and subsequently infected with mCherry-labelled STM WT or *ssaV*. The formation of SIF was followed by LCI using an SDCM from 1-7 h post infection (p.i.) with hourly intervals.

We set out to execute the analysis by visual inspection following the example of Stein et al. [20], who performed a mutant library screen to identify bacterial factors involved in SIF formation. As the dynamic nature of SIF and phenotypic heterogeneity in the cellular context excluded fully automated analysis, we decided to perform the analysis by visual inspections and used a MATLAB-based tool named SifScreen to support data input and collection (Figure 2). This tool queried the presence of SIF in the examined field of view as the main feature in a binary manner (for detailed information see Suppl. Material).

Since siRNA silencing usually does not yield 100% loss of function, we did not expect a complete lack of SIF in each of the eight images per well. Furthermore, since a considerable number of cells were present per image (roughly 10-30 cells, depending on applied siRNA and position on plate) a single SIF-forming cell would have prompted a SIF-positive scoring, even

18.11.2019

Host factors for SIF formation

if generally a SIF-negative knockdown might have occurred. Thus, we decided to define an overall SIF-abolishing hit with a comparably high cutoff of 50%, i.e. if less than 50% of the images showed SIF. However, this did not take into consideration knockdowns possibly affecting cell viability in general or other circumstances compromising the analysis. Since these parameters were also queried by SifScreen, this allowed us to differentiate between ‘true hits’ and ‘possible hits,’ scoring the former and latter with values of 3 and 1, respectively. Additionally, to avoid a possible bias due to visual analysis, each screening plate was analyzed independently by two investigators. With the screen performed in biological triplicates, we subsequently compiled all scoring data for each host target, also pooling the results of the three individual siRNAs per target. This resulted in a list of final hits shown in Table S2, in which hits with a cumulative scoring of 1-4, 5-7, or ≥ 8 were classified as low-, mid- and high-ranking hits, respectively.

Approximately 81% (404 of 496) of the trafficome targets scored to varying degrees positive, underlining the general importance of trafficking processes for SIF formation. Table 1 shows selected high-ranking hits involved in trafficking and cytoskeleton biology. These hits clearly show the involvement of all protein classes necessary for the vesicle budding and fusion machinery, the core of cellular trafficking. These comprise: (i) small GTPases, especially Rab GTPases, as primary regulators [36-38]; (ii) vesicle coats and their adaptors as cargo and budding mediators [39-43]; (iii) cytoskeleton components as the basis for vesicle motility [44]; (iv) tethering factors as part of the fusion specification [45, 46]; (v) SNAREs as the primary fusion agents [45, 47, 48]. Besides, this list includes hits of diverse subcellular origin, encompassing the complete secretory and endo-/lysosomal system, i.e. endoplasmic reticulum (ER), Golgi apparatus, endo-/lysosomes. Supporting these allocations, the interaction network of the hits from Table 1 shows several distinct clusters (Figure 3). Two of them are connected to cytoskeleton biology (also interconnected if lower-ranking hits are included, data not shown). Another cluster is SNARE-centered, including RAB1A with RAB11A as a node between this

18.11.2019

Host factors for SIF formation

cluster and one of the cytoskeleton-related clusters. Lastly, one cluster is associated with COPI and clathrin-coated vesicles (CCVs). Collectively, the overall results of the trafficome screen confirm the general importance of host trafficking factors in SIF biogenesis, indicating a crucial role for a plethora of as yet unprecedented factors in STM pathobiology.

Validation of selected hits

To test the validity of our approach and the resulting hits, we focused on a subset of genes due to their presence in the noticeable interaction clusters depicted in Figure 3, or prior reports on involvement in STM pathobiology. HGS was chosen due to being the highest-ranking hit (Table S2) and the interaction of SPI1-T3SS effector SopB with endosomal sorting complex required for transport (ESCRT) complexes previously reported [49] with HGS being part of the ESCRT-0 complex. Furthermore, RAB1A and RAB11A were included as the highest-ranking Rab GTPases with RAB11A previously being shown to colocalize with SCV as well as SIF [32, 50]. Consistently, RAB7A served as another well-established SCV- and SIF-localizing control [12, 50-52]. STX5, STX7, VAMP7, and VAMP8 were chosen due to being the highest-ranking SNAREs (except VAMP8 lacking from the trafficome), the colocalization of STX7 with SIF [50], and the recent reported essential role of VAMP7 in SIF biogenesis [33]. The AAA ATPase VCP was included as another of the highest-ranking trafficking-related hits and another host factor already known to be important for proper SCV as well as SIF biogenesis via STM effector SptP [53]. Finally, the VPS11 core component of the class C core vacuole/endosome tethering (CORVET) / homotypic fusion and protein sorting (HOPS) group of multisubunit tethering complexes (MTCs) was chosen due to the HOPS complex functionally bridging RAB7 with late endo-/lysosomal SNAREs and the recent recognition of its essential role in STM replication and SCV and SIF biogenesis [54, 55].

The success of the siRNA knockdowns was confirmed by RT-PCR with a consistently significant decrease in mRNA in most cases down to 5-10% compared to control siAllStars (Figure S1, siSKIP served as the screen-inherent phenotype-specific control). Next, we exactly

18.11.2019

Host factors for SIF formation

quantified the reduction of SIF formation due to the silencing of the selected targets (Figure 4). The *ssaV* mutant strain and the knockdown with siSKIP served as screen-inherent SIF-abolishing controls. All knockdowns resulted in decreased SIF formation with the reduction being statistically significant except for siHGS and siSTX5. The siRAB7A had the highest impact, siVCP the second-highest and the others ranged similar. Thus, the knockdowns of VAMP7 and VCP meet previous data (even though SIF abolishment regarding VCP depletion here is less pronounced) [33, 53]. Altogether, the effect on SIF formation demonstrates that our approach confirms known host factors, but also allows the identification of novel factors to be crucial for SIF biology.

STM deploys membranes of early and late secretory, late endo-/lysosomal, and clathrin-coated origin in SIF biogenesis

The fact that host factors appear as hits in our screen, clearly indicates a physiologically relevant role in SIF biogenesis. However, whether this role is by direct interaction or an indirect one involving several intermittent steps, remains unclear. Thus, we decided to analyze the localization of selected hits with regard to SIF (Figure 5, Figure 6, Figure 7).

For analyses of RAB GTPases (Figure 5), we again used RAB7A as well as RAB9A as positive controls, both showing a clear colocalization with SIF. Of the several Rab GTPases included in the trafficome RAB1A showed the highest score (Table S2). RAB1 GTPases are responsible for anterograde ER-Golgi trafficking [56-59]. Importantly, RAB1A can be functionally substituted by RAB1B [60, 61] and an STM replication-targeted RNAi screen identified specifically RAB1B as a hit [31]. Hence, we analyzed the infection-related localization of both isoforms and detected a partial and a strong colocalization of RAB1A and RAB1B, respectively, with SIF (Figure 5).

Another high-ranking hit with relation to RAB proteins was RAB3GAP2 (Table S2), the non-catalytic subunit of the RAB3 inactivating GTPase-activating protein (GAP) complex [62]. RAB3 possesses four isoforms in mammals [63] and is involved in regulated exocytosis [64].

18.11.2019

Host factors for SIF formation

As neither the catalytic GAP subunit, RAB3GAP1, nor one of the four isoforms were present in the trafficome, we decided to analyze the localization of RAB3A and found a partial colocalization with SIF (Figure 5).

Besides, a mid-ranking hit was RAB8A (Table S2), a Golgi- and endosome-localized RAB likewise involved in exocytic processes [65]. Interestingly, its isoform RAB8B was observed to be excluded from maturing SCVs (≤ 3 h p.i.) [50]. Therefore, we analyzed the localization of both, RAB8A and RAB8B, and strikingly found a strong colocalization of not only RAB8A but also RAB8B with SIF (Figure 5).

As several RAB proteins participating in the late secretory system/exocytosis seem to play a role in SIF biogenesis, we additionally analyzed three SNAREs with exocytic roles not present in the trafficome: VAMP2, VAMP3, and VAMP4 [66-68], with VAMP2 also shown to be present on early SCV [69]. Apart from that, the presence of the two high-ranking SNARE hits STX7 and VAMP7 (Table S2) on SIF was previously shown [33, 50]. However, SNAREs, are part of complexes of usually four proteins participating in membrane fusion and consisting of a single protein v-SNARE (on the vesicle or incoming membrane) and a ternary t-SNARE subcomplex (on the target or accepting membrane). VAMP7 is the v-SNARE in the SNARE complex for heterotypic LE/lysosome fusions with the t-SNAREs STX7, STX8, and VTI1B [70, 71] being replaced by VAMP8 in homotypic LE fusions [72, 73]. In fact, the presence of VTI1B and STX8 on early SCV [69, 74] and their role in STM replication [54], as well as the involvement of VAMP8 in STM invasion were already shown [75]. However, their interaction with SIF remains unclear, except VAMP8 silencing causing SIF reduction identified here (Figure 4). Thus, we analyzed the localization of VTI1B, STX8, and VAMP8 using STX7 and VAMP7 as controls. As shown in Figure 6, we detected prominent association of STX7 and VAMP7 with SIF, as well as for VAMP2 and VAMP8. Colocalization of VTI1B, STX8, VAMP3, and VAMP4 with SIF was also observed. However, these SNARE subunits showed a more heterogeneous distribution, and only a fraction of SIF was positive for these candidates.

18.11.2019

Host factors for SIF formation

279 Notably, both recent proteomic studies [32, 33] and this screen identified AP2A1 as being
 280 present on SMM/SCV or scoring high-ranking (Table S2), respectively. AP2A1 represents one
 281 of the two core subunit isoforms of the canonical AP-2 adaptor complex usually acting in
 282 clathrin-mediated endocytosis (CME) at the plasma membrane [76, 77]. Accordingly, the main
 283 coat determinants implicated in the formation of CCVs were among the high-scoring hits,
 284 including both clathrin light chains, CLTA and CLTB, as well as the conventional heavy chain,
 285 CLTC (Table S2). Thus, we analyzed the localization of CLTA and observed a partial
 286 colocalization with SIF (Figure 7).

287 In conclusion, the colocalization of various host factors involved in cellular transport with SIF
 288 validated the results of the RNAi screen. These proteins are components of SIF tubules and, to
 289 variable extent, required for the formation of SIF. The data also support the highly diverse
 290 origin of host cell membranes involved in SIF formation.

18.11.2019

Host factors for SIF formation

Discussion

By applying a targeted RNAi screen, we identified several new host factors required for the formation of SIF, and partially characterized interactions of host proteins with SMM. Our data strengthen the involvement of the late endo-/lysosomal SNARE complex, and reveal new interactions of SIF with RAB1, RAB3, and RAB8 GTPases, exocytic SNAREs, as well as clathrin-coated structures. The implications of these findings as discussed below are depicted in Figure 8. Several host trafficome components identified here were previously shown as involved in infection biology in STM in general, and specifically in SCV and/or SIF biogenesis including: dynein – DYNC1H1 [78-80], filamin – FLNA [81], myosin II – MYH10 [82], VPS4A/B [49]. This also holds true for several mid-ranking hits: kinesin-1 – KIF5A/B [22-25, 83], PIKFYVE [84], RAB9A [85, 86], RAB14 [85], SCAMP3 [87].

Complementary data were recently provided by two proteomic studies. Our group analyzed the SMM proteome in the late phase of infection (8 h p.i.) [32] that contained several host proteins that are mid- or high-ranking hits in this screen (summarized in Table 2, first column). The colocalization of several of these proteins with SMM was shown by immunostaining or LCI. Santos et al. [33] determined the proteomes of early and maturing SCV (30 min p.i. and 3 h p.i., respectively) again identifying proteins appearing as hits in this screen (see Table 2, second and third column). Taken together, these data strongly validate the approach deployed here.

The approach reported here has a major advantage compared to studies based on organelle proteomics [32, 33]. Proteomics shown presence or absence of host factors on the organelle of interest, but a particular role in the biogenesis of this organelle cannot be implied directly. In our RNAi approach potentially each, or at least each high-ranking hit, points to a role in STM-induced endosomal remodeling. However, a functional role revealed by RNAi does not necessarily require colocalization of the host factor with the compartment, because a function may be mediated indirectly, involving several interacting partners. We analyzed the localization of selected factors (Figure 5, Figure 6, Figure 7) and found several differences in the host factor

18.11.2019

Host factors for SIF formation

sets identified by proteomics or by our approach. Nevertheless, there is a considerable overlap of host factors identified by both approaches as represented in Table 2.

The interaction of microbial pathogens with regulators of the early secretory system, such as *Legionella* and *Coxiella* with RAB1 [88-90], and *Brucella* with RAB2 [91, 92], is well established. However, an interaction of STM with the early secretory system, e.g. RAB2A, was only recently described by proteomic studies [32, 33]. We now expand this interaction by showing the physiological relevance of RAB1A in SIF formation (Figure 4), as well as the presence of both RAB1A and RAB1B on SIF (Figure 5). This is in striking contrast to the previous observation that RAB1A is detrimental to STM replication due to its role in antibacterial autophagy, which is counteracted by the SPI2-T3SS effectors SseFG [93, 94]. Future work has to clarify this apparent conundrum, at least regarding RAB1A. The direct association of RAB1A/B with SIF possibly connects several distinct trafficking events. First, RAB1B was shown to be involved in formation of the COPI vesicle coat, which participates in *intra*-Golgi and retrograde Golgi-to-ER transport. The formation of COPI is dependent on RAB1B due to its effector GBF1, which is an activating guanine exchange factor (GEF) of ARF1, the primary COPI-regulating small GTPase [95-97]. Second, the COPI components COPA and CPG1 were shown to partly colocalize with SIF [32]. In accordance, our screen identified the majority of COPI components as mid- or high-ranking hits (ARCN1, COPA/B1/B2/G1, Table 1 and Figure 3, though ARF1 did not score at all, and GBF1 only low). Thus, RAB1A and/or RAB1B might represent a physical link between COPI vesicles and SCV and/or SIF for the redirection of early secretory material as depicted in Figure 8iv.

The physical interaction of SIF with COPI vesicles might be, similar to the late endo-/lysosomal fusion machinery, additionally accompanied by tethering factors and SNAREs. The conserved oligomeric Golgi (COG) tethering complex was shown to be a RAB1 effector and directly bind COPI components [98, 99]. Interestingly, all components of the COG present in the trafficome scored mid- to high-ranking (COG1/2/3/5/7, Table S2). Additionally, COG binds STX5, a

18.11.2019

Host factors for SIF formation

SNARE that is part of several ER-Golgi and *intra*-Golgi transport-related SNARE complexes [100]. These complexes either consist of STX5, GOSR2/GS27/membrin, BET1, SEC22B [101], or STX5, GOSR1/GS28, BET1, YKT6 [102], or STX5, GOSR1, BET1L/GS15, YKT6 [103]. All of these SNAREs are present in the trafficome, but strikingly only the components of the first SNARE complex scored all mid- to high-ranking (Table 1 and Table S2). STM effectors partaking in this interaction might be SseF and SseG, as they were recently shown in a BioID screen [104] to interact with STX5 and SEC22B, besides PipB2 also interacting with SEC22B. However, the potential involvements indicated by these collective data remain to be elucidated.

Transport protein particle (TRAPP) complexes I, II, and III were identified as RAB1 GEFs [105-108], as well as COPII [TRAPPI; 109, 110, 111], and COPI tethers [TRAPP2; 107, 112]. TRAPPI is the core shared by all TRAPP complexes, with II and III processing unique additional subunits. TRAPPC8, the unique component of TRAPPIII, scored high-ranking. Other components were not present in this trafficome, except the TRAPPI core subunit TRAPPC2, which unexpectedly did not score at all (Table S2). So far TRAPPIII is only characterized to participate in autophagy [108, 113, 114]. Finally, the golgin USO1/p115 scored mid-ranking (Table S2). It is also a RAB1B effector and COPI and COPII tether [97, 115, 116], partly in conjunction with COG [117], besides being likewise able to bind STX5 [118]. For both TRAPP complexes and USO1, a specific role in SIF biogenesis remains to be elucidated. It has already been described that STM, depending on SseF/SseG, recruits to the SCV exocytic vesicles from the Golgi apparatus destined to the plasma membrane [119]. Which host factors are involved in this process was unclear, and our work now sheds light on this phenotype by showing the presence of exocytic RABs, RAB3A, RAB8A, and RAB8B (Figure 5) on SIF, as well as exocytic SNAREs, i.e. VAMP2, VAMP3, and VAMP4 (Figure 6).

Besides their involvement in exocytosis, VAMP4 and VAMP3 are also known to prominently participate in endosome-to-Golgi transport in conjunction with STX16, VTI1A, and STX6 or

18.11.2019

Host factors for SIF formation

STX10 for EEs or LEs, respectively [120, 121]. Although STX6 and STX10 were not included in the trafficome and STX16 ranked low, VTI1A was a mid-ranking hit (Table S2). The STM-mediated redirection of LAMP1-containing vesicles from the Golgi apparatus to the early SCV was shown to involve recruitment of STX6 and VAMP2 via SPI1-T3SS effector SipC [69]. Alternatively, this might happen via SPI2-T3SS effector PipB2 that was identified in the recent BioID screen as interactor of VAMP2 [104]. Furthermore, in homotypic EE fusion STX16 is replaced by STX13 [122], and STX13 was previously shown to be present on early SCV [74, 123]. While the exact role of RAB3A and the identity of SNAREs involved remain to be determined, this might indicate that the interception of secretory vesicles depends on a SNARE complex comprising a distinct combination of the abovementioned SNAREs as represented in Figure 8ii.

In addition to its exocytic role, RAB8A is involved in recycling processes as indicated by the localization on tubular recycling endosomes (RE) [124]. This localization depends on several factors such as RAB8 GEF RAB3IP/RABIN8 (which is also part of the trafficome, though it ranked low, Table S2), concurrently being an effector of RE master regulator RAB11 [125]. Another factor is the RE-localized MICALL1, which interacts with the dynamin-like ATPases EHD1 and EHD3 [124, 126, 127]. Interestingly, MICALL1 was identified in an RNAi screen with focus on STM intracellular replication [31], EHD1 and EHD2 were present in the proteome of maturing SCV [33], and EHD4 was present in late SMM [32]. Moreover, the association of the maturing SCV and late SMM with RAB11A/B was shown previously [32, 50], with RAB11A following RAB1A as the second highest-ranking RAB in our screen (Table S2). RAB8B was not present in our screen, but showed a clear colocalization with SIF. In a previous screen RAB8B was shown to be excluded from maturing SCV [50], however only early events up to 3 h p.i. were analyzed. Temporal differences in RAB8B recruitment could explain these observations. SPI2-T3SS effector SopD2 most likely plays a role in RAB8 recruitment, as it was previously shown to interact with RAB8 [104, 128]. Collectively, these

18.11.2019

Host factors for SIF formation

data strongly argue for a continued association of STM not only with exocytic compartments, but also with recycling compartments at later time points (summarized in Figure 8ii). Concurrently, a RAB11-to-RAB8 switch might occur on SMM, similar to the RAB5-to-RAB7 and RAB14-to-RAB9 switches, probably involving RAB3IP/RABIN8. However, unlike the enduring RAB5-to-RAB7 switch and similar to the RAB14-to-RAB9 switch, this seems to happen repeatedly in a transient manner due to the continued presence of RAB11 on SIF as with RAB14 [32].

The presence on SIF, and/or importance for SIF formation of RAB7, the HOPS complex, STX7, and VAMP7, as well as the direct fusion of late endo-/lysosomal-like VAMP7-positive vesicles with the SCV, was shown before [33, 50, 54]. This indicates the involvement of the complete canonical mammalian late endo-/lysosomal vesicle fusion machinery in SIF biogenesis. Whether this interaction cascade also employs the canonical STX7, VTI1B, and STX8 was not fully clarified. Here, we expand this cascade by showing the physiological relevance of STX7 for SIF formation (Figure 4), and the presence of VTI1B and STX8 on SIF (Figure 6, Figure 5) as depicted in Figure 8iii. This cascade is possibly expanded by the host protein PLEKHM1, as the recruitment of RAB7 and the HOPS complex by SifA via the host protein PLEKHM1 and its involvement in SCV biogenesis was recently revealed [55], most likely also being involved in SIF biogenesis. Taken together, SifA seems to recruit the complete late endo-/lysosomal fusion machinery. Thus, SifA performs a dual role besides the binding of SKIP and the SIF mobility connected with it. This is also corroborated by the identification of interactions of SifA with STX7 and VAMP7 by the recent BioID screen [104]. Alternatively or in addition, SopD2 might be likewise involved as it was also shown to interact with STX7 and VAMP7 besides VTI1B in the same study.

Data on involvement of clathrin-coated structures or adaptor protein complexes in infection processes of intracellular bacterial pathogens are scarce. The usurpation of CME during the internalization/invasion process is only recently being recognized to be employed by several

18.11.2019

Host factors for SIF formation

bacteria, best studied in *Listeria monocytogenes* infection [129, 130]. However, only one study on *Brucella abortus*, a pathogen also residing in a vacuole during intracellular lifestyle, demonstrated the association of clathrin with the *Brucella*-containing vacuole [131]. We now show such an association with CLTA also for STM (Figure 7). It is peculiar that proteomics, as well as our screen, indicate an involvement of the AP-2 complex, but one of the other adaptor complexes. This is noteworthy due to the CME-related AP-2 being primarily plasma membrane-localized, in contrast to the Golgi traffic-related AP-1 and AP-4, or the endo-/lysosomal traffic-related AP-3 and AP-5 [43]. Especially AP-3 deserves detailed analyses since its two core subunit isoforms scored in mid- and high-ranking range (AP3B1 and AP3D1, Table S2, see Figure 8i). An interaction of AP-3 with VAMP7 in mammalian cells [132-134], as well as an interaction with the complete late endo-/lysosomal SNARE complex in *Dictyostelium discoideum* was previously shown [135]. Several SPI2-T3SS effectors, i.e. PipB2, SopD2, and SseG, might participate in such a recruitment because a recent BioID screen revealed interaction with various AP-2 and AP-3 core subunits [104]. However, examination of other AP complexes also seems worthwhile, since the latter study indicates interactions with several of them and the trafficome screen did not comprehensively cover AP complex.

In summary, we successfully employed a sub-genomic RNAi screen to systematically identify new host factors, corresponding protein complexes, and pathways involved in SIF formation. By providing physiologically relevant data regarding SIF formation, this work further corroborates the promiscuous origin of SMM indicated by previous proteomics studies [32, 33]. Similar future screens can also reveal the biogenesis of several other SIT [15], and extend to the host cell types important for *Salmonella* pathogenesis.

Acknowledgements

We thank Monika Nietschke and Ursula Krehe for construction of plasmids and technical assistance. Special thanks go to Markus C. Kerr (Brisbane), André P. Mäurer, and Thomas F.

18.11.2019

Host factors for SIF formation

448 Meyer (Berlin) for advice and logistics in setting up the screen. We thank Martin Aepfelbacher
 449 (Hamburg), Thierry Galli (Paris), Wanjin Hong (Singapore), and Yulong Li (Beijing) for
 450 providing transfection vectors, and acknowledge DNASU und Addgene for provision of
 451 materials.

18.11.2019

Host factors for SIF formation

Materials and Methods

Bacterial strains and growth conditions

For infection STM NCTC 12023 WT and isogenic SPI2-T3SS-defective strain P2D6 harboring plasmid pFPV-mCherry/2 or isogenic GFP-expressing MvP1897 were used (for details see Table S6). Strains were routinely grown in Luria-Bertani (LB) broth (Difco, BD, Heidelberg, Germany) containing 50 µg/mL carbenicillin for plasmid selection at 37 °C with aeration.

Cell lines and cell culture

Experiments were performed using the parental HeLa cell line (ATCC No. CCL-2) or the lentivirus-transfected HeLa cell line stably expressing LAMP1-GFP [18]. Cells were routinely cultured in Dulbecco's modified Eagle's medium (DMEM) containing 4.5 g/L glucose, 4 mM stable glutamine, and sodium pyruvate (Biochrom, Berlin, Germany) supplemented with 10% inactivated fetal calf serum (iFCS; Gibco, Darmstadt, Germany) in an atmosphere of 5% CO₂ and 90% humidity at 37 °C.

siRNA library and individual siRNAs

The siRNA library used comprised siRNAs targeting 496 host proteins mostly involved in intracellular trafficking (but also other processes such as metabolism) with a threefold coverage. The siRNAs were part of a human whole-genome library obtained from Qiagen (Hilden, Germany) deposited at the Max Planck Institute for Infection Biology (Berlin, Germany). A volume of 4 µL of each siRNA (0.2 µM, end concentration of 5.2 nM) was spotted automatically onto 96-well Clear Bottom Black Cell Culture Microplates (Corning, Corning, NY, USA) and frozen at -20 °C before transfer. Additionally, each plate contained the same amount of the following siRNAs from Qiagen as knockdown controls: AllStars as negative and Hs_PLK1_7 (directed against the cell cycle protein polo-like kinase 1) as positive controls. A custom siRNA from Qiagen directed against SKIP served as a phenotype-specific control [22]

18.11.2019

Host factors for SIF formation

and was spotted on location. Information including target sequences for these siRNAs, as well as those ordered for validation experiments, are listed in Table S7.

Reverse transfection with siRNA

If not using 96-well screening plates as detailed above, the amount for an end concentration of 5 nM siRNA was spotted onto standard cell culture 6-well plates (for mRNA extraction, TPP, Trasadingen, Switzerland) or 8-well polymer bottom chamber slides (for quantification of SIF formation, μ -Slides, ibidi, Martinsried, Germany).

Next, a mixture of the transfection reagent HiPerFect (Qiagen, Hilden, Germany) and serum-free cell culture medium was applied and this was incubated for 5-10 min at room temperature (RT). Subsequently, 5,000, 125,000, or 20,000 cells per well of 96-well plates, 6-well plates, or 8-well chamber slides, respectively, were added in serum-containing medium and incubated for 72 h at 37 °C in a humidified atmosphere containing 5% CO₂.

Gene expression quantification

After reverse transfection with different siRNAs total RNA of cells was extracted using the RNeasy Mini Kit following the manufacturer's instructions (Qiagen, Hilden, Germany). Homogenization during extraction was performed using Qiagen QIAshredder columns. Then, 1 μ g of RNA digested with DNaseI (NEB, Frankfurt a. M., Germany) was used for reverse transcription of mRNA with the RevertAid First Strand cDNA Synthesis Kit (Thermo Scientific, Dreieich, Germany) following the manufacturer's instructions using the Oligo(dT)₁₈ primer. For RT-PCR 1 μ L of cDNA was used with the Thermo Scientific Maxima SYBR Green/Fluorescein qPCR Master Mix (2x). As reference gene the housekeeping gene *GAPDH* was selected [136]. For control of individual host factor knockdowns primers were used employing the PrimerBank database [137, 138]. Primers for those as well as *GAPDH* are listed in Table S8. Primer concentration was 150 nM each, and primer efficiency was determined for each primer pair. RT-PCR was performed in an iCycler instrument (Bio-Rad, Munich,

18.11.2019

Host factors for SIF formation

Germany) in triplicates in 96-well plates. Relative expression was determined using the $2^{-\Delta Ct}$ method [84, 139] with *GAPDH* expression set as 100%. Results were plotted using SigmaPlot 11 (Systat Software, Erkrath, Germany).

Construction of plasmids

Plasmids used in this study were either obtained from Addgene, kind gifts from various laboratories, or cloned by Gibson Assembly or restriction enzyme digests and are listed in Table S6. Oligonucleotides for the construction of plasmids encoding host proteins fused to mRuby2 or EGFP are listed in Table S8. First, N- or C-terminal mRuby2 vectors were cloned. For that, the vectors pEGFP-C1 and pEGFP-N1 were amplified and EGFP was exchanged for a fragment encoding mRuby2. Genes encoding host proteins were amplified from vectors obtained from DNASU (Table S6) and then inserted into mRuby2 vectors by Gibson Assembly. Plasmids encoding host proteins fused to EGFP were constructed using restriction enzyme digests. The vector pEGFP-C3 was digested with *KpnI* and *XbaI* or *KpnI* and *BamHI* and the larger fragment was recovered. The inserts were treated the same way and fragments were ligated.

Host cell transfection

For LCI for localization of host factors, HeLa or HeLa-LAMP1-GFP cells were seeded 1 d prior to transfection. About 20.000 or 150.000 cells were seeded in 8-well chamber slides (see above) or 3.5 mm glass bottom dishes (FluoroDish, WPI, Berlin, Germany), respectively. For transfection 0.5 or 2 μ g of plasmid DNA in 25 or 200 μ L serum-free medium were mixed with 1 or 4 μ L of FuGENE HD transfection reagent (DNA to reagent ratio of 1:2, Promega, Mannheim, Germany) and incubated for 10 min at RT. Medium on the cells was changed and transfection mixture applied. Cells were incubated for at least 18 h before infection with medium change during infection. For a complete list of transfection plasmids, see Table S6.

Infection experiments

18.11.2019

Host factors for SIF formation

Overnight cultures of STM were diluted 1:31 and grown for additional 3.5 h in LB broth in glass test tubes with agitation in a roller drum. HeLa cells were infected with STM WT or *ssaV* for screening approaches in 96-well plates with a multiplicity of infection (MOI) of 15, otherwise for colocalization analysis or SIF quantification in 8-well chamber slides or FluoroDishes with an MOI of 75 or 50, respectively. Infection only of 96-well plates was synchronized by centrifugation at 500 x g for 5 min, and in all cases proceeded for 25 min at 37 °C in a humidified atmosphere containing 5% CO₂. Cells were washed thrice with full medium or PBS for screening or non-screen LCI purposes, respectively, and incubated in full medium containing 100 µg/mL gentamicin for 1 h to eliminate extracellular bacteria. Then medium containing 10 µg/mL gentamicin was applied for the remainder of the experiment.

Live cell imaging

For LCI full medium was replaced by imaging medium consisting of Minimal Essential Medium (MEM) with Earle's salts, without NaHCO₃, without L-glutamine and without phenol red (Biochrom, Berlin, Germany) supplemented with 30 mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid) (Sigma-Aldrich, Taufkirchen, Germany), pH 7.4, containing 10 µg/mL gentamicin. Fluorescence imaging for screening purposes was performed using a Zeiss Cell Observer microscope with Yokogawa Spinning Disk Unit CSU-X1 (Carl Zeiss, Göttingen, Germany), Evolve 512 x 512 EMCCD camera (Photometrics, Tucson, AZ, USA), automated PZ-2000 stage (Applied Scientific Instrumentation, Eugene, OR, USA), and infrared-based focus system Definite Focus, operated by Zeiss ZEN 2012 software (blue edition). The microscope was equipped with live cell periphery consisting of a custom-made incubation chamber surrounding the microscope body and connected with "The Cube" heating unit (Life Imaging Services, Basel, Switzerland) maintaining 37 °C and the Incubation System S for CO₂ and humidity supply (PeCon, Erbach, Germany). Images were acquired using the Zeiss LD Plan-Neofluar 40x/0.6 Corr air objective (with bottom thickness correction ring). For acquisition of GFP and mCherry BP 525/50 (Zeiss) and LP 580 (Olympus, Hamburg,

18.11.2019

Host factors for SIF formation

Germany) filters, respectively, were applied. All images obtained were processed by the ZEN software. Non-screen LCI was performed using a Leica SP5 confocal laser-scanning microscope (CLSM) operated by Leica LAS AF software. The microscope was also equipped with live cell periphery consisting of ‘The Box’ incubation chamber (Life Imaging Services, Basel, Switzerland), a custom-made heating unit and a gas supply unit ‘The Brick’. Images were acquired using the HCX PL APO CS 100x/1.4 oil objective (Leica, Wetzlar, Germany), applying the polychroic mirror TD 488/543/633 for acquisition of GFP and mCherry. All images were processed by LAS AF software.

Quantification of SIF formation

After siRNA knockdown and infection, 100 infected HeLa-LAMP1-GFP cells per condition were examined live from 6-8 h p.i. for presence of SIF as exhibited by WT-infected cells, and percentage calculated. Results from biological triplicates were plotted using SigmaPlot 11.

Data analysis

For central entry and collection of scoring data, the MATLAB-based utility SifScreen was used. Categorization of targets/hits was executed using the Gene Ontology classification scheme [140, 141]. For visualization of protein interactions, the STRING v10 database with default settings was applied [142].

18.11.2019

Host factors for SIF formation

References

1. LaRock DL, Chaudhary A, Miller SI. Salmonellae interactions with host processes. Nat Rev Microbiol. 2015;13(4):191-205. doi: 10.1038/nrmicro3420. PubMed PMID: 25749450.
2. Galan JE, Curtiss R. Cloning and molecular characterization of genes whose products allow *Salmonella typhimurium* to penetrate tissue culture cells. Proc Natl Acad Sci USA. 1989;86:6383-7.
3. Ramos-Morales F. Impact of *Salmonella enterica* Type III Secretion System Effectors on the Eukaryotic Host Cell. ISRN Cell Biology. 2012;2012. doi: 10.5402/2012/787934.
4. Shea JE, Hensel M, Gleeson C, Holden DW. Identification of a virulence locus encoding a second type III secretion system in *Salmonella typhimurium*. Proc Natl Acad Sci U S A. 1996;93(6):2593-7.
5. Ochman H, Soncini FC, Solomon F, Groisman EA. Identification of a pathogenicity island required for *Salmonella* survival in host cells. Proc Natl Acad Sci U S A. 1996;93:7800-4.
6. Figueira R, Holden DW. Functions of the *Salmonella* pathogenicity island 2 (SPI-2) type III secretion system effectors. Microbiology. 2012;158(Pt 5):1147-61. Epub 2012/03/17. doi: mic.0.058115-0 [pii] 10.1099/mic.0.058115-0. PubMed PMID: 22422755.
7. Steele-Mortimer O, Meresse S, Gorvel JP, Toh BH, Finlay BB. Biogenesis of *Salmonella typhimurium*-containing vacuoles in epithelial cells involves interactions with the early endocytic pathway. Cell Microbiol. 1999;1(1):33-49.
8. Mukherjee K, Siddiqi SA, Hashim S, Raje M, Basu SK, Mukhopadhyay A. Live *Salmonella* recruits N-ethylmaleimide-sensitive fusion protein on phagosomal membrane and promotes fusion with early endosome. J Cell Biol. 2000;148(4):741-53.
9. Mills SD, Finlay BB. Comparison of *Salmonella typhi* and *Salmonella typhimurium* invasion, intracellular growth and localization in cultured human epithelial cells. MicrobPathog. 1994;17:409-23.
10. Oh YK, Alpuche-Aranda C, Berthiaume E, Jinks T, Miller SI, Swanson JA. Rapid and complete fusion of macrophage lysosomes with phagosomes containing *Salmonella typhimurium*. Infect Immun. 1996;64(9):3877-83. Epub 1996/09/01. PubMed PMID: 8751942; PubMed Central PMCID: PMC174306.
11. Garcia-del Portillo F, Zwick MB, Leung KY, Finlay BB. Intracellular replication of *Salmonella* within epithelial cells is associated with filamentous structures containing lysosomal membrane glycoproteins. Infect Agents Dis. 1993;2(4):227-31.
12. Meresse S, Steele-Mortimer O, Finlay BB, Gorvel JP. The rab7 GTPase controls the maturation of *Salmonella typhimurium*-containing vacuoles in HeLa cells. EMBO J. 1999;18(16):4394-403.
13. Brumell JH, Tang P, Mills SD, Finlay BB. Characterization of *Salmonella*-induced filaments (Sifs) reveals a delayed interaction between *Salmonella*-containing vacuoles and late endocytic compartments. Traffic. 2001;2(9):643-53. PubMed PMID: 11555418.
14. Garcia-del Portillo F, Finlay BB. Targeting of *Salmonella typhimurium* to vesicles containing lysosomal membrane glycoproteins bypasses compartments with mannose 6-phosphate receptors. J Cell Biol. 1995;129(1):81-97.
15. Schroeder N, Mota LJ, Meresse S. *Salmonella*-induced tubular networks. Trends Microbiol. 2011;19(6):268-77. Epub 2011/03/01. doi: 10.1016/j.tim.2011.01.006. PubMed PMID: 21353564.
16. Garcia-del Portillo F, Zwick MB, Leung KY, Finlay BB. *Salmonella* induces the formation of filamentous structures containing lysosomal membrane glycoproteins in epithelial cells. Proc Natl Acad Sci U S A. 1993;90(22):10544-8. PubMed PMID: 8248143.

18.11.2019

Host factors for SIF formation

- 618 17. Knuff K, Finlay BB. What the SIF is happening - The role of intracellular *Salmonella*-
619 induced filaments. Front Cell Infect Microbiol. 2017;7:335. doi: 10.3389/fcimb.2017.00335.
620 PubMed PMID: 28791257; PubMed Central PMCID: PMC5524675.
- 621 18. Krieger V, Liebl D, Zhang Y, Rajashekar R, Chlanda P, Giesker K, et al. Reorganization
622 of the endosomal system in *Salmonella*-infected cells: the ultrastructure of *Salmonella*-induced
623 tubular compartments. PLoS Pathog. 2014;10(9):e1004374. doi:
624 10.1371/journal.ppat.1004374. PubMed PMID: 25254663.
- 625 19. Liss V, Hensel M. Take the tube: remodelling of the endosomal system by intracellular
626 *Salmonella enterica*. Cell Microbiol. 2015;17(5):639-47. doi: 10.1111/cmi.12441. PubMed
627 PMID: 25802001.
- 628 20. Stein MA, Leung KY, Zwick M, Garcia-del Portillo F, Finlay BB. Identification of a
629 *Salmonella* virulence gene required for formation of filamentous structures containing
630 lysosomal membrane glycoproteins within epithelial cells. Mol Microbiol. 1996;20(1):151-64.
631 PubMed PMID: 8861213.
- 632 21. Zhao W, Moest T, Zhao Y, Guilhon AA, Buffat C, Gorvel JP, et al. The *Salmonella*
633 effector protein SifA plays a dual role in virulence. Sci Rep. 2015;5:12979. doi:
634 10.1038/srep12979. PubMed PMID: 26268777; PubMed Central PMCID: PMC4534788.
- 635 22. Boucrot E, Henry T, Borg JP, Gorvel JP, Meresse S. The intracellular fate of *Salmonella*
636 depends on the recruitment of kinesin. Science. 2005;308(5725):1174-8.
- 637 23. Henry T, Couillault C, Rockenfeller P, Boucrot E, Dumont A, Schroeder N, et al. The
638 *Salmonella* effector protein PipB2 is a linker for kinesin-1. Proc Natl Acad Sci U S A.
639 2006;103(36):13497-502.
- 640 24. Kaniuk NA, Canadien V, Bagshaw RD, Bakowski M, Braun V, Landekic M, et al.
641 *Salmonella* exploits Arl8B-directed kinesin activity to promote endosome tubulation and cell-
642 to-cell transfer. Cell Microbiol. 2011;13(11):1812-23. Epub 2011/08/10. doi: 10.1111/j.1462-
643 5822.2011.01663.x. PubMed PMID: 21824248.
- 644 25. Rosa-Ferreira C, Munro S. Arl8 and SKIP act together to link lysosomes to kinesin-1.
645 Dev Cell. 2011;21(6):1171-8. doi: 10.1016/j.devcel.2011.10.007. PubMed PMID: 22172677;
646 PubMed Central PMCID: PMC3240744.
- 647 26. Leone P, Meresse S. Kinesin regulation by *Salmonella*. Virulence. 2011;2(1):63-6. Epub
648 2011/01/11. doi: 14603 [pii]. PubMed PMID: 21217202.
- 649 27. Misselwitz B, Dilling S, Vonaesch P, Sacher R, Snijder B, Schlumberger M, et al. RNAi
650 screen of *Salmonella* invasion shows role of COPI in membrane targeting of cholesterol and
651 Cdc42. Mol Syst Biol. 2011;7:474. Epub 2011/03/17. doi: 10.1038/msb.2011.7. PubMed
652 PMID: 21407211.
- 653 28. Thornbrough JM, Gopinath A, Hundley T, Worley MJ. Human genome-wide RNAi
654 screen for host factors that facilitate *Salmonella* invasion reveals a role for potassium secretion
655 in promoting internalization. PLoS One. 2016;11(11):e0166916. doi:
656 10.1371/journal.pone.0166916. PubMed PMID: 27880807; PubMed Central PMCID:
657 PMC5120809.
- 658 29. Kuijl C, Savage ND, Marsman M, Tuin AW, Janssen L, Egan DA, et al. Intracellular
659 bacterial growth is controlled by a kinase network around PKB/AKT1. Nature.
660 2007;450(7170):725-30. Epub 2007/11/30. doi: nature06345 [pii]
661 10.1038/nature06345. PubMed PMID: 18046412.
- 662 30. Albers HM, Kuijl C, Bakker J, Hendrickx L, Wekker S, Farhou N, et al. Integrating
663 chemical and genetic silencing strategies to identify host kinase-phosphatase inhibitor networks
664 that control bacterial infection. ACS Chem Biol. 2014;9(2):414-22. doi: 10.1021/cb400421a.
665 PubMed PMID: 24274083; PubMed Central PMCID: PMC3934374.
- 666 31. Thornbrough JM, Hundley T, Valdivia R, Worley MJ. Human genome-wide RNAi
667 screen for host factors that modulate intracellular *Salmonella* growth. PLoS One.

18.11.2019

Host factors for SIF formation

- 2012;7(6):e38097. doi: 10.1371/journal.pone.0038097. PubMed PMID: 22701604; PubMed Central PMCID: PMC3372477.
32. Vorwerk S, Krieger V, Deiwick J, Hensel M, Hansmeier N. Proteomes of host cell membranes modified by intracellular activities of *Salmonella enterica*. Mol Cell Proteomics. 2015;14(1):81-92. doi: 10.1074/mcp.M114.041145. PubMed PMID: 25348832; PubMed Central PMCID: PMC4288265.
33. Santos JC, Duchateau M, Fredlund J, Weiner A, Mallet A, Schmitt C, et al. The COPII complex and lysosomal VAMP7 determine intracellular *Salmonella* localization and growth. Cell Microbiol. 2015;17(12):1699-720. doi: 10.1111/cmi.12475. PubMed PMID: 26084942.
34. Drecktrah D, Levine-Wilkinson S, Dam T, Winfree S, Knodler LA, Schroer TA, et al. Dynamic behavior of *Salmonella*-induced membrane tubules in epithelial cells. Traffic. 2008;9(12):2117-29. Epub 2008/09/13. doi: TRA830 [pii] 10.1111/j.1600-0854.2008.00830.x. PubMed PMID: 18785994.
35. Rajashekar R, Liebl D, Seitz A, Hensel M. Dynamic remodeling of the endosomal system during formation of *Salmonella*-induced filaments by intracellular *Salmonella enterica*. Traffic. 2008;9(12):2100-16. Epub 2008/09/27. doi: TRA821 [pii] 10.1111/j.1600-0854.2008.00821.x. PubMed PMID: 18817527.
36. Hutagalung AH, Novick PJ. Role of Rab GTPases in membrane traffic and cell physiology. Physiol Rev. 2011;91(1):119-49. doi: 10.1152/physrev.00059.2009. PubMed PMID: 21248164; PubMed Central PMCID: PMC3710122.
37. Bhuin T, Roy JK. Rab proteins: the key regulators of intracellular vesicle transport. Exp Cell Res. 2014;328(1):1-19. doi: 10.1016/j.yexcr.2014.07.027. PubMed PMID: 25088255.
38. Wandinger-Ness A, Zerial M. Rab proteins and the compartmentalization of the endosomal system. Cold Spring Harb Perspect Biol. 2014;6(11):a022616. doi: 10.1101/cshperspect.a022616. PubMed PMID: 25341920; PubMed Central PMCID: PMC34413231.
39. Brodsky FM. Diversity of clathrin function: new tricks for an old protein. Annu Rev Cell Dev Biol. 2012;28:309-36. doi: 10.1146/annurev-cellbio-101011-155716. PubMed PMID: 22831640.
40. Burd C, Cullen PJ. Retromer: a master conductor of endosome sorting. Cold Spring Harb Perspect Biol. 2014;6(2). doi: 10.1101/cshperspect.a016774. PubMed PMID: 24492709; PubMed Central PMCID: PMC3941235.
41. Popoff V, Adolf F, Brugger B, Wieland F. COPI budding within the Golgi stack. Cold Spring Harb Perspect Biol. 2011;3(11):a005231. doi: 10.1101/cshperspect.a005231. PubMed PMID: 21844168; PubMed Central PMCID: PMC3220356.
42. Lord C, Ferro-Novick S, Miller EA. The highly conserved COPII coat complex sorts cargo from the endoplasmic reticulum and targets it to the golgi. Cold Spring Harb Perspect Biol. 2013;5(2). doi: 10.1101/cshperspect.a013367. PubMed PMID: 23378591; PubMed Central PMCID: PMC3552504.
43. Park SY, Guo X. Adaptor protein complexes and intracellular transport. Biosci Rep. 2014;34(4). doi: 10.1042/BSR20140069. PubMed PMID: 24975939; PubMed Central PMCID: PMC4114066.
44. Anitei M, Hoflack B. Bridging membrane and cytoskeleton dynamics in the secretory and endocytic pathways. Nat Cell Biol. 2012;14(1):11-9. doi: 10.1038/ncb2409. PubMed PMID: 22193159.
45. Hong W, Lev S. Tethering the assembly of SNARE complexes. Trends Cell Biol. 2014;24(1):35-43. doi: 10.1016/j.tcb.2013.09.006. PubMed PMID: 24119662.
46. Chia PZ, Gleeson PA. Membrane tethering. F1000Prime Rep. 2014;6:74. doi: 10.12703/P6-74. PubMed PMID: 25343031; PubMed Central PMCID: PMC4166942.
47. Hong W. SNAREs and traffic. Biochim Biophys Acta. 2005;1744(3):493-517. PubMed PMID: 16038056.

18.11.2019

Host factors for SIF formation

48. Jahn R, Scheller RH. SNAREs--engines for membrane fusion. *Nat Rev Mol Cell Biol.* 2006;7(9):631-43. doi: 10.1038/nrm2002. PubMed PMID: 16912714.
49. Dukes JD, Lee H, Hagen R, Reaves BJ, Layton AN, Galyov EE, et al. The secreted *Salmonella dublin* phosphoinositide phosphatase, SopB, localizes to PtdIns(3)P-containing endosomes and perturbs normal endosome to lysosome trafficking. *Biochem J.* 2006;395(2):239-47. doi: 10.1042/BJ20051451. PubMed PMID: 16396630; PubMed Central PMCID: PMC1422764.
50. Smith AC, Heo WD, Braun V, Jiang X, Macrae C, Casanova JE, et al. A network of Rab GTPases controls phagosome maturation and is modulated by *Salmonella enterica* serovar Typhimurium. *J Cell Biol.* 2007;176(3):263-8. Epub 2007/01/31. doi: jcb.200611056 [pii] 10.1083/jcb.200611056. PubMed PMID: 17261845.
51. Harrison RE, Brumell JH, Khandani A, Bucci C, Scott CC, Jiang X, et al. *Salmonella* impairs RILP recruitment to Rab7 during maturation of invasion vacuoles. *Mol Biol Cell.* 2004;15:3146-54.
52. D'Costa VM, Braun V, Landekic M, Shi R, Proteau A, McDonald L, et al. *Salmonella* Disrupts Host Endocytic Trafficking by SopD2-Mediated Inhibition of Rab7. *Cell Rep.* 2015;12(9):1508-18. Epub 2015/08/25. doi: 10.1016/j.celrep.2015.07.063. PubMed PMID: 26299973.
53. Humphreys D, Hume PJ, Koronakis V. The *Salmonella* effector SptP dephosphorylates host AAA+ ATPase VCP to promote development of its intracellular replicative niche. *Cell Host Microbe.* 2009;5(3):225-33. doi: 10.1016/j.chom.2009.01.010. PubMed PMID: 19286132; PubMed Central PMCID: PMC2724103.
54. Sindhvani A, Arya SB, Kaur H, Jagga D, Tuli A, Sharma M. *Salmonella* exploits the host endolysosomal tethering factor HOPS complex to promote its intravacuolar replication. *PLoS Pathog.* 2017;13(10):e1006700. Epub 2017/10/31. doi: 10.1371/journal.ppat.1006700. PubMed PMID: 29084291; PubMed Central PMCID: PMC5679646.
55. McEwan DG, Richter B, Claudi B, Wigge C, Wild P, Farhan H, et al. PLEKHM1 regulates *Salmonella*-containing vacuole biogenesis and infection. *Cell Host Microbe.* 2015;17(1):58-71. doi: 10.1016/j.chom.2014.11.011. PubMed PMID: 25500191.
56. Segev N, Mulholland J, Botstein D. The yeast GTP-binding YPT1 protein and a mammalian counterpart are associated with the secretion machinery. *Cell.* 1988;52(6):915-24. PubMed PMID: 3127057.
57. Bacon RA, Salminen A, Ruohola H, Novick P, Ferro-Novick S. The GTP-binding protein Ypt1 is required for transport in vitro: the Golgi apparatus is defective in ypt1 mutants. *J Cell Biol.* 1989;109(3):1015-22. PubMed PMID: 2504726; PubMed Central PMCID: PMC2115776.
58. Plutner H, Cox AD, Pind S, Khosravi-Far R, Bourne JR, Schwaninger R, et al. Rab1b regulates vesicular transport between the endoplasmic reticulum and successive Golgi compartments. *J Cell Biol.* 1991;115(1):31-43. PubMed PMID: 1918138; PubMed Central PMCID: PMC2289927.
59. Tisdale EJ, Bourne JR, Khosravi-Far R, Der CJ, Balch WE. GTP-binding mutants of rab1 and rab2 are potent inhibitors of vesicular transport from the endoplasmic reticulum to the Golgi complex. *J Cell Biol.* 1992;119(4):749-61. PubMed PMID: 1429835; PubMed Central PMCID: PMC2289685.
60. Mukhopadhyay A, Nieves E, Che FY, Wang J, Jin L, Murray JW, et al. Proteomic analysis of endocytic vesicles: Rab1a regulates motility of early endocytic vesicles. *J Cell Sci.* 2011;124(Pt 5):765-75. doi: 10.1242/jcs.079020. PubMed PMID: 21303926; PubMed Central PMCID: PMC3039020.
61. Mukhopadhyay A, Quiroz JA, Wolkoff AW. Rab1a regulates sorting of early endocytic vesicles. *Am J Physiol Gastrointest Liver Physiol.* 2014;306(5):G412-24. doi:

18.11.2019

Host factors for SIF formation

- 10.1152/ajpgi.00118.2013. PubMed PMID: 24407591; PubMed Central PMCID: PMCPMC3949023.
62. Nagano F, Sasaki T, Fukui K, Asakura T, Imazumi K, Takai Y. Molecular cloning and characterization of the noncatalytic subunit of the Rab3 subfamily-specific GTPase-activating protein. *J Biol Chem.* 1998;273(38):24781-5. doi: 10.1074/jbc.273.38.24781. PubMed PMID: 9733780.
63. Touchot N, Chardin P, Tavitian A. Four additional members of the ras gene superfamily isolated by an oligonucleotide strategy: molecular cloning of YPT-related cDNAs from a rat brain library. *Proc Natl Acad Sci USA.* 1987;84(23):8210-4. doi: 10.1073/pnas.84.23.8210. PubMed PMID: 3317403; PubMed Central PMCID: PMCPMC299511.
64. Schlüter OM, Khvotchev M, Jahn R, Südhof TC. Localization versus function of Rab3 proteins. Evidence for a common regulatory role in controlling fusion. *J Biol Chem.* 2002;277(43):40919-29. doi: 10.1074/jbc.M203704200. PubMed PMID: 12167638.
65. Huber LA, Pimplikar S, Parton RG, Virta H, Zerial M, Simons K. Rab8, a small GTPase involved in vesicular traffic between the TGN and the basolateral plasma membrane. *J Cell Biol.* 1993;123(1):35-45. PubMed PMID: 8408203; PubMed Central PMCID: PMCPMC2119815.
66. Söllner T, Bennett MK, Whiteheart SW, Scheller RH, Rothman JE. A protein assembly-disassembly pathway in vitro that may correspond to sequential steps of synaptic vesicle docking, activation, and fusion. *Cell.* 1993;75(3):409-18. PubMed PMID: 8221884.
67. Borisovska M, Zhao Y, Tsytsyura Y, Glyvuk N, Takamori S, Matti U, et al. v-SNAREs control exocytosis of vesicles from priming to fusion. *EMBO J.* 2005;24(12):2114-26. doi: 10.1038/sj.emboj.7600696. PubMed PMID: 15920476; PubMed Central PMCID: PMCPMC1150890.
68. Cocucci E, Racchetti G, Rupnik M, Meldolesi J. The regulated exocytosis of energeosomes is mediated by a SNARE machinery that includes VAMP4. *J Cell Sci.* 2008;121(Pt 18):2983-91. doi: 10.1242/jcs.032029. PubMed PMID: 18713833.
69. Madan R, Rastogi R, Parashuraman S, Mukhopadhyay A. Salmonella acquires lysosome-associated membrane protein 1 (LAMP1) on phagosomes from Golgi via SipC protein-mediated recruitment of host Syntaxin6. *J Biol Chem.* 2012;287(8):5574-87. doi: 10.1074/jbc.M111.286120. PubMed PMID: 22190682; PubMed Central PMCID: PMC3285332.
70. Bogdanovic A, Bennett N, Kieffer S, Louwagie M, Morio T, Garin J, et al. Syntaxin 7, syntaxin 8, Vti1 and VAMP7 (vesicle-associated membrane protein 7) form an active SNARE complex for early macropinocytic compartment fusion in Dictyostelium discoideum. *Biochem J.* 2002;368(Pt 1):29-39. doi: 10.1042/BJ20020845. PubMed PMID: 12175335; PubMed Central PMCID: PMCPMC1222979.
71. Pryor PR, Mullock BM, Bright NA, Lindsay MR, Gray SR, Richardson SC, et al. Combinatorial SNARE complexes with VAMP7 or VAMP8 define different late endocytic fusion events. *EMBO Rep.* 2004;5(6):590-5. doi: 10.1038/sj.emboj.7400150. PubMed PMID: 15133481; PubMed Central PMCID: PMCPMC1299070.
72. Antonin W, Holroyd C, Fasshauer D, Pabst S, Von Mollard GF, Jahn R. A SNARE complex mediating fusion of late endosomes defines conserved properties of SNARE structure and function. *EMBO J.* 2000;19(23):6453-64.
73. Antonin W, Holroyd C, Tikkanen R, Honing S, Jahn R. The R-SNARE endobrevin/VAMP-8 mediates homotypic fusion of early endosomes and late endosomes. *Mol Biol Cell.* 2000;11(10):3289-98. PubMed PMID: 11029036; PubMed Central PMCID: PMCPMC14992.
74. Singh PK, Kapoor A, Lomash RM, Kumar K, Kamerkar SC, Pucadyil TJ, et al. *Salmonella* SipA mimics a cognate SNARE for host Syntaxin8 to promote fusion with early

18.11.2019

Host factors for SIF formation

- 819 endosomes. J Cell Biol. 2018;217(12):4199-214. doi: 10.1083/jcb.201802155. PubMed PMID:
820 30309979; PubMed Central PMCID: PMC6279372.
- 821 75. Dai S, Zhang Y, Weimbs T, Yaffe MB, Zhou D. Bacteria-generated PtdIns(3)P recruits
822 VAMP8 to facilitate phagocytosis. Traffic. 2007;8(10):1365-74. PubMed PMID: 17645435.
- 823 76. Merrifield CJ, Kaksonen M. Endocytic accessory factors and regulation of clathrin-
824 mediated endocytosis. Cold Spring Harb Perspect Biol. 2014;6(11):a016733. doi:
825 10.1101/cshperspect.a016733. PubMed PMID: 25280766; PubMed Central PMCID:
826 PMC64413230.
- 827 77. Traub LM, Bonifacino JS. Cargo recognition in clathrin-mediated endocytosis. Cold
828 Spring Harb Perspect Biol. 2013;5(11):a016790. doi: 10.1101/cshperspect.a016790. PubMed
829 PMID: 24186068; PubMed Central PMCID: PMC3809577.
- 830 78. Marsman M, Jordens I, Kuijl C, Janssen L, Neefjes J. Dynein-mediated vesicle transport
831 controls intracellular *Salmonella* replication. Mol Biol Cell. 2004;15(6):2954-64.
- 832 79. Abrahams GL, Müller P, Hensel M. Functional dissection of SseF, a type III effector
833 protein involved in positioning the *Salmonella*-containing vacuole. Traffic. 2006;7(8):950-65.
- 834 80. Guignot J, Caron E, Beuzon C, Bucci C, Kagan J, Roy C, et al. Microtubule motors
835 control membrane dynamics of *Salmonella*-containing vacuoles. J Cell Sci. 2004;117:1033-45.
- 836 81. Miao EA, Brittnacher M, Haraga A, Jeng RL, Welch MD, Miller SI. *Salmonella*
837 effectors translocated across the vacuolar membrane interact with the actin cytoskeleton. Mol
838 Microbiol. 2003;48(2):401-15. PubMed PMID: 12675800.
- 839 82. Wasylanka JA, Bakowski MA, Szeto J, Ohlson MB, Trimble WS, Miller SI, et al. Role
840 for myosin II in regulating positioning of *Salmonella*-containing vacuoles and intracellular
841 replication. Infect Immun. 2008;76(6):2722-35. Epub 2008/04/16. doi: IAI.00152-08 [pii]
842 10.1128/IAI.00152-08. PubMed PMID: 18411289; PubMed Central PMCID: PMC2423101.
- 843 83. Dumont A, Boucrot E, Drevensek S, Daire V, Gorvel JP, Pous C, et al. SKIP, the host
844 target of the *Salmonella* virulence factor SifA, promotes kinesin-1-dependent vacuolar
845 membrane exchanges. Traffic. 2010;11(7):899-911. Epub 2010/04/22. doi: TRA1069 [pii]
846 10.1111/j.1600-0854.2010.01069.x. PubMed PMID: 20406420.
- 847 84. Kerr MC, Wang JT, Castro NA, Hamilton NA, Town L, Brown DL, et al. Inhibition of
848 the PtdIns(5) kinase PIKfyve disrupts intracellular replication of *Salmonella*. EMBO J.
849 2010;29(8):1331-47. doi: 10.1038/emboj.2010.28. PubMed PMID: 20300065; PubMed Central
850 PMCID: PMC2868569.
- 851 85. Kuijl C, Pilli M, Alahari SK, Janssen H, Khoo PS, Ervin KE, et al. Rac and Rab GTPases
852 dual effector Nischarin regulates vesicle maturation to facilitate survival of intracellular
853 bacteria. EMBO J. 2013;32(5):713-27. doi: 10.1038/emboj.2013.10. PubMed PMID:
854 23386062; PubMed Central PMCID: PMC3590985.
- 855 86. Seixas E, Ramalho JS, Mota LJ, Barral DC, Seabra MC. Bacteria and protozoa
856 differentially modulate the expression of Rab proteins. PLoS One. 2012;7(7):e39858. doi:
857 10.1371/journal.pone.0039858. PubMed PMID: 22911692; PubMed Central PMCID:
858 PMC3401185.
- 859 87. Mota LJ, Ramsden AE, Liu M, Castle JD, Holden DW. SCAMP3 is a component of the
860 *Salmonella*-induced tubular network and reveals an interaction between bacterial effectors and
861 post-Golgi trafficking. Cell Microbiol. 2009;11(8):1236-53. Epub 2009/05/15. doi: CMI1329
862 [pii]
863 10.1111/j.1462-5822.2009.01329.x. PubMed PMID: 19438519; PubMed Central PMCID:
864 PMC2730479.
- 865 88. Campoy EM, Zoppino FC, Colombo MI. The early secretory pathway contributes to the
866 growth of the *Coxiella*-replicative niche. Infect Immun. 2011;79(1):402-13. doi:
867 10.1128/IAI.00688-10. PubMed PMID: 20937765; PubMed Central PMCID:
868 PMC3019900.

18.11.2019

Host factors for SIF formation

- 869 89. Kagan JC, Stein MP, Pypaert M, Roy CR. *Legionella* subvert the functions of rab1 and
870 sec22b to create a replicative organelle. J Exp Med. 2004;199(9):1201-11.
- 871 90. Müller MP, Peters H, Blumer J, Blankenfeldt W, Goody RS, Itzen A. The *Legionella*
872 effector protein DrrA AMPylates the membrane traffic regulator Rab1b. Science.
873 2010;329(5994):946-9. doi: 10.1126/science.1192276. PubMed PMID: 20651120.
- 874 91. Fugier E, Salcedo SP, de Chastellier C, Pophillat M, Muller A, Arce-Gorvel V, et al.
875 The glyceraldehyde-3-phosphate dehydrogenase and the small GTPase Rab 2 are crucial for
876 *Brucella* replication. PLoS Pathog. 2009;5(6):e1000487. doi: 10.1371/journal.ppat.1000487.
877 PubMed PMID: 19557163; PubMed Central PMCID: PMC2695806.
- 878 92. de Bolle X, Letesson JJ, Gorvel JP. Small GTPases and *Brucella* entry into the
879 endoplasmic reticulum. Biochem Soc Trans. 2012;40(6):1348-52. doi: 10.1042/BST20120156.
880 PubMed PMID: 23176479.
- 881 93. Huang J, Birmingham CL, Shahnazari S, Shiu J, Zheng YT, Smith AC, et al.
882 Antibacterial autophagy occurs at PI(3)P-enriched domains of the endoplasmic reticulum and
883 requires Rab1 GTPase. Autophagy. 2011;7(1):17-26. Epub 2010/10/29. doi: 13840 [pii].
884 PubMed PMID: 20980813; PubMed Central PMCID: PMC3039730.
- 885 94. Feng ZZ, Jiang AJ, Mao AW, Feng Y, Wang W, Li J, et al. The *Salmonella* effectors
886 SseF and SseG inhibit Rab1A-mediated autophagy to facilitate intracellular bacterial survival
887 and replication. J Biol Chem. 2018;293(25):9662-73. doi: 10.1074/jbc.M117.811737. PubMed
888 PMID: 29610274; PubMed Central PMCID: PMC6016468.
- 889 95. Alvarez C, Garcia-Mata R, Brandon E, Sztul E. COPI recruitment is modulated by a
890 Rab1b-dependent mechanism. Mol Biol Cell. 2003;14(5):2116-27. doi: 10.1091/mbc.E02-09-
891 0625. PubMed PMID: 12802079; PubMed Central PMCID: PMC165101.
- 892 96. Monetta P, Slavin I, Romero N, Alvarez C. Rab1b interacts with GBF1 and modulates
893 both ARF1 dynamics and COPI association. Mol Biol Cell. 2007;18(7):2400-10. doi:
894 10.1091/mbc.E06-11-1005. PubMed PMID: 17429068; PubMed Central PMCID:
895 PMC1924811.
- 896 97. Guo Y, Linstedt AD. Binding of the vesicle docking protein p115 to the GTPase Rab1b
897 regulates membrane recruitment of the COPI vesicle coat. Cell Logist. 2013;3:e27687. doi:
898 10.4161/cl.27687. PubMed PMID: 25332841; PubMed Central PMCID: PMC4187009.
- 899 98. Suvorova ES, Duden R, Lupashin VV. The Sec34/Sec35p complex, a Ypt1p effector
900 required for retrograde intra-Golgi trafficking, interacts with Golgi SNAREs and COPI vesicle
901 coat proteins. J Cell Biol. 2002;157(4):631-43. doi: 10.1083/jcb.200111081. PubMed PMID:
902 12011112; PubMed Central PMCID: PMC2173848.
- 903 99. Zolov SN, Lupashin VV. Cog3p depletion blocks vesicle-mediated Golgi retrograde
904 trafficking in HeLa cells. J Cell Biol. 2005;168(5):747-59. doi: 10.1083/jcb.200412003.
905 PubMed PMID: 15728195; PubMed Central PMCID: PMC171815.
- 906 100. Shestakova A, Suvorova E, Pavliv O, Khaidakova G, Lupashin V. Interaction of the
907 conserved oligomeric Golgi complex with t-SNARE Syntaxin5a/Sed5 enhances intra-Golgi
908 SNARE complex stability. J Cell Biol. 2007;179(6):1179-92. doi: 10.1083/jcb.200705145.
909 PubMed PMID: 18086915; PubMed Central PMCID: PMC2140037.
- 910 101. Xu D, Joglekar AP, Williams AL, Hay JC. Subunit structure of a mammalian ER/Golgi
911 SNARE complex. J Biol Chem. 2000;275(50):39631-9. doi: 10.1074/jbc.M007684200.
912 PubMed PMID: 11035026.
- 913 102. Zhang T, Hong W. Ykt6 forms a SNARE complex with syntaxin 5, GS28, and Bet1 and
914 participates in a late stage in endoplasmic reticulum-Golgi transport. J Biol Chem.
915 2001;276(29):27480-7. doi: 10.1074/jbc.M102786200. PubMed PMID: 11323436.
- 916 103. Tai G, Lu L, Wang TL, Tang BL, Goud B, Johannes L, et al. Participation of the syntaxin
917 5/Ykt6/GS28/GS15 SNARE complex in transport from the early/recycling endosome to the
918 *trans*-Golgi network. Mol Biol Cell. 2004;15(9):4011-22. doi: 10.1091/mbc.E03-12-0876.
919 PubMed PMID: 15215310; PubMed Central PMCID: PMC155336.

18.11.2019

Host factors for SIF formation

- 920 104. D'Costa VM, Coyaud E, Boddy KC, Laurent EMN, St-Germain J, Li T, et al. BioID
921 screen of *Salmonella* type 3 secreted effectors reveals host factors involved in vacuole
922 positioning and stability during infection. Nat Microbiol. 2019. doi: 10.1038/s41564-019-0580-
923 9. PubMed PMID: 31611645.
- 924 105. Sacher M, Barrowman J, Wang W, Horecka J, Zhang Y, Pypaert M, et al. TRAPP I
925 implicated in the specificity of tethering in ER-to-Golgi transport. Mol Cell. 2001;7(2):433-42.
926 PubMed PMID: 11239471.
- 927 106. Jones S, Newman C, Liu F, Segev N. The TRAPP complex is a nucleotide exchanger
928 for Ypt1 and Ypt31/32. Mol Biol Cell. 2000;11(12):4403-11. PubMed PMID: 11102533;
929 PubMed Central PMCID: PMC15082.
- 930 107. Yamasaki A, Menon S, Yu S, Barrowman J, Meerloo T, Oorschot V, et al. mTrs130 is
931 a component of a mammalian TRAPP II complex, a Rab1 GEF that binds to COPI-coated
932 vesicles. Mol Biol Cell. 2009;20(19):4205-15. doi: 10.1091/mbc.E09-05-0387. PubMed PMID:
933 19656848; PubMed Central PMCID: PMC2754934.
- 934 108. Lynch-Day MA, Bhandari D, Menon S, Huang J, Cai H, Bartholomew CR, et al. Trs85
935 directs a Ypt1 GEF, TRAPP III, to the phagophore to promote autophagy. Proc Natl Acad Sci
936 U S A. 2010;107(17):7811-6. Epub 2010/04/09. doi: 1000063107 [pii]
937 10.1073/pnas.1000063107. PubMed PMID: 20375281; PubMed Central PMCID:
938 PMC2867920.
- 939 109. Cai H, Yu S, Menon S, Cai Y, Lazarova D, Fu C, et al. TRAPP I tethers COPII vesicles
940 by binding the coat subunit Sec23. Nature. 2007;445(7130):941-4. doi: 10.1038/nature05527.
941 PubMed PMID: 17287728.
- 942 110. Yu S, Satoh A, Pypaert M, Mullen K, Hay JC, Ferro-Novick S. mBet3p is required for
943 homotypic COPII vesicle tethering in mammalian cells. J Cell Biol. 2006;174(3):359-68. doi:
944 10.1083/jcb.200603044. PubMed PMID: 16880271; PubMed Central PMCID:
945 PMC2064232.
- 946 111. Loh E, Peter F, Subramaniam VN, Hong W. Mammalian Bet3 functions as a cytosolic
947 factor participating in transport from the ER to the Golgi apparatus. J Cell Sci. 2005;118(Pt
948 6):1209-22. doi: 10.1242/jcs.01723. PubMed PMID: 15728249.
- 949 112. Cai H, Zhang Y, Pypaert M, Walker L, Ferro-Novick S. Mutants in *trs120* disrupt traffic
950 from the early endosome to the late Golgi. J Cell Biol. 2005;171(5):823-33. doi:
951 10.1083/jcb.200505145. PubMed PMID: 16314430; PubMed Central PMCID:
952 PMC2171297.
- 953 113. Lamb CA, Nuhlen S, Judith D, Frith D, Snijders AP, Behrends C, et al. TBC1D14
954 regulates autophagy via the TRAPP complex and ATG9 traffic. EMBO J. 2016;35(3):281-301.
955 doi: 10.15252/embj.201592695. PubMed PMID: 26711178; PubMed Central PMCID:
956 PMC4741301.
- 957 114. Shirahama-Noda K, Kira S, Yoshimori T, Noda T. TRAPP III is responsible for
958 vesicular transport from early endosomes to Golgi, facilitating Atg9 cycling in autophagy. J
959 Cell Sci. 2013;126(Pt 21):4963-73. doi: 10.1242/jcs.131318. PubMed PMID: 23986483.
- 960 115. Allan BB, Moyer BD, Balch WE. Rab1 recruitment of p115 into a *cis*-SNARE complex:
961 programming budding COPII vesicles for fusion. Science. 2000;289(5478):444-8. PubMed
962 PMID: 10903204.
- 963 116. Cao X, Ballew N, Barlowe C. Initial docking of ER-derived vesicles requires Uso1p and
964 Ypt1p but is independent of SNARE proteins. EMBO J. 1998;17(8):2156-65. doi:
965 10.1093/emboj/17.8.2156. PubMed PMID: 9545229; PubMed Central PMCID:
966 PMC1170560.
- 967 117. Sohda M, Misumi Y, Yoshimura S, Nakamura N, Fusano T, Ogata S, et al. The
968 interaction of two tethering factors, p115 and COG complex, is required for Golgi integrity.
969 Traffic. 2007;8(3):270-84. doi: 10.1111/j.1600-0854.2006.00530.x. PubMed PMID:
970 17274799.

18.11.2019

Host factors for SIF formation

- 971 118. Shorter J, Beard MB, Seemann J, Dirac-Svejstrup AB, Warren G. Sequential tethering
972 of Golgins and catalysis of SNAREpin assembly by the vesicle-tethering protein p115. J Cell
973 Biol. 2002;157(1):45-62. doi: 10.1083/jcb.200112127. PubMed PMID: 11927603; PubMed
974 Central PMCID: PMCPMC2173270.
- 975 119. Kuhle V, Abrahams GL, Hensel M. Intracellular *Salmonella enterica* redirect exocytic
976 transport processes in a *Salmonella* pathogenicity island 2-dependent manner. Traffic.
977 2006;7(6):716-30.
- 978 120. Mallard F, Tang BL, Galli T, Tenza D, Saint-Pol A, Yue X, et al. Early/recycling
979 endosomes-to-TGN transport involves two SNARE complexes and a Rab6 isoform. J Cell Biol.
980 2002;156(4):653-64. doi: 10.1083/jcb.200110081. PubMed PMID: 11839770; PubMed Central
981 PMCID: PMCPMC2174079.
- 982 121. Ganley IG, Espinosa E, Pfeffer SR. A syntaxin 10-SNARE complex distinguishes two
983 distinct transport routes from endosomes to the trans-Golgi in human cells. J Cell Biol.
984 2008;180(1):159-72. doi: 10.1083/jcb.200707136. PubMed PMID: 18195106; PubMed Central
985 PMCID: PMCPMC2213607.
- 986 122. Brandhorst D, Zwillig D, Rizzoli SO, Lippert U, Lang T, Jahn R. Homotypic fusion of
987 early endosomes: SNAREs do not determine fusion specificity. Proc Natl Acad Sci USA.
988 2006;103(8):2701-6. doi: 10.1073/pnas.0511138103. PubMed PMID: 16469845; PubMed
989 Central PMCID: PMCPMC1413832.
- 990 123. Smith AC, Cirulis JT, Casanova JE, Scidmore MA, Brumell JH. Interaction of the
991 *Salmonella*-containing vacuole with the endocytic recycling system. J Biol Chem.
992 2005;280(26):24634-41. PubMed PMID: 15886200.
- 993 124. Roland JT, Kenworthy AK, Peranen J, Caplan S, Goldenring JR. Myosin Vb interacts
994 with Rab8a on a tubular network containing EHD1 and EHD3. Mol Biol Cell. 2007;18(8):2828-
995 37. doi: 10.1091/mbc.E07-02-0169. PubMed PMID: 17507647; PubMed Central PMCID:
996 PMCPMC1949367.
- 997 125. Knodler A, Feng S, Zhang J, Zhang X, Das A, Peranen J, et al. Coordination of Rab8
998 and Rab11 in primary ciliogenesis. Proc Natl Acad Sci U S A. 2010;107(14):6346-51. doi:
999 10.1073/pnas.1002401107. PubMed PMID: 20308558; PubMed Central PMCID:
1000 PMCPMC2851980.
- 1001 126. Rahajeng J, Giridharan SS, Cai B, Naslavsky N, Caplan S. MICAL-L1 is a tubular
1002 endosomal membrane hub that connects Rab35 and Arf6 with Rab8a. Traffic. 2012;13(1):82-
1003 93. doi: 10.1111/j.1600-0854.2011.01294.x. PubMed PMID: 21951725; PubMed Central
1004 PMCID: PMCPMC3302426.
- 1005 127. Sharma M, Giridharan SS, Rahajeng J, Naslavsky N, Caplan S. MICAL-L1 links EHD1
1006 to tubular recycling endosomes and regulates receptor recycling. Mol Biol Cell.
1007 2009;20(24):5181-94. doi: 10.1091/mbc.E09-06-0535. PubMed PMID: 19864458; PubMed
1008 Central PMCID: PMCPMC2793294.
- 1009 128. Spano S, Gao X, Hannemann S, Lara-Tejero M, Galan JE. A Bacterial Pathogen Targets
1010 a Host Rab-Family GTPase Defense Pathway with a GAP. Cell Host Microbe. 2016;19(2):216-
1011 26. doi: 10.1016/j.chom.2016.01.004. PubMed PMID: 26867180; PubMed Central PMCID:
1012 PMCPMC4854434.
- 1013 129. Cossart P, Lebreton A. A trip in the "New Microbiology" with the bacterial pathogen
1014 *Listeria monocytogenes*. FEBS Lett. 2014;588(15):2437-45. doi:
1015 10.1016/j.febslet.2014.05.051. PubMed PMID: 24911203.
- 1016 130. Pizarro-Cerda J, Kuhbacher A, Cossart P. Entry of *Listeria monocytogenes* in
1017 mammalian epithelial cells: an updated view. Cold Spring Harb Perspect Med. 2012;2(11). doi:
1018 10.1101/cshperspect.a010009. PubMed PMID: 23125201; PubMed Central PMCID:
1019 PMCPMC3543101.
- 1020 131. Lee JJ, Kim DG, Kim DH, Simborio HL, Min W, Lee HJ, et al. Interplay between
1021 clathrin and Rab5 controls the early phagocytic trafficking and intracellular survival of *Brucella*

18.11.2019

Host factors for SIF formation

- 1022 *abortus* within HeLa cells. J Biol Chem. 2013;288(39):28049-57. doi:
1023 10.1074/jbc.M113.491555. PubMed PMID: 23940042; PubMed Central PMCID:
1024 PMCPMC3784717.
- 1025 132. Martinez-Arca S, Rudge R, Vacca M, Raposo G, Camonis J, Proux-Gillardeaux V, et
1026 al. A dual mechanism controlling the localization and function of exocytic v-SNAREs. Proc
1027 Natl Acad Sci U S A. 2003;100(15):9011-6. doi: 10.1073/pnas.1431910100. PubMed PMID:
1028 12853575; PubMed Central PMCID: PMCPMC166429.
- 1029 133. Kent HM, Evans PR, Schafer IB, Gray SR, Sanderson CM, Luzio JP, et al. Structural
1030 basis of the intracellular sorting of the SNARE VAMP7 by the AP3 adaptor complex. Dev Cell.
1031 2012;22(5):979-88. doi: 10.1016/j.devcel.2012.01.018. PubMed PMID: 22521722; PubMed
1032 Central PMCID: PMCPMC3549491.
- 1033 134. Salazar G, Craige B, Styers ML, Newell-Litwa KA, Doucette MM, Wainer BH, et al.
1034 BLOC-1 complex deficiency alters the targeting of adaptor protein complex-3 cargoes. Mol
1035 Biol Cell. 2006;17(9):4014-26. doi: 10.1091/mbc.E06-02-0103. PubMed PMID: 16760431;
1036 PubMed Central PMCID: PMCPMC1556383.
- 1037 135. Bennett N, Letourneur F, Ragno M, Louwagie M. Sorting of the v-SNARE VAMP7 in
1038 Dictyostelium discoideum: a role for more than one Adaptor Protein (AP) complex. Exp Cell
1039 Res. 2008;314(15):2822-33. doi: 10.1016/j.yexcr.2008.06.019. PubMed PMID: 18634783.
- 1040 136. Vreeburg RA, Bastiaan-Net S, Mes JJ. Normalization genes for quantitative RT-PCR in
1041 differentiated Caco-2 cells used for food exposure studies. Food Funct. 2011;2(2):124-9. doi:
1042 10.1039/c0fo00068j. PubMed PMID: 21779557.
- 1043 137. Wang X, Seed B. A PCR primer bank for quantitative gene expression analysis. Nucleic
1044 Acids Res. 2003;31(24):e154. PubMed PMID: 14654707; PubMed Central PMCID:
1045 PMCPMC291882.
- 1046 138. Wang X, Spandidos A, Wang H, Seed B. PrimerBank: a PCR primer database for
1047 quantitative gene expression analysis, 2012 update. Nucleic Acids Res. 2012;40(Database
1048 issue):D1144-9. doi: 10.1093/nar/gkr1013. PubMed PMID: 22086960; PubMed Central
1049 PMCID: PMCPMC3245149.
- 1050 139. Town L, McGlinn E, Fiorenza S, Metzis V, Butterfield NC, Richman JM, et al. The
1051 metalloendopeptidase gene *Pitrm1* is regulated by hedgehog signaling in the developing mouse
1052 limb and is expressed in muscle progenitors. Dev Dyn. 2009;238(12):3175-84. doi:
1053 10.1002/dvdy.22126. PubMed PMID: 19877269.
- 1054 140. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene
1055 ontology: tool for the unification of biology. The Gene Ontology Consortium. Nat Genet.
1056 2000;25(1):25-9. doi: 10.1038/75556. PubMed PMID: 10802651; PubMed Central PMCID:
1057 PMCPMC3037419.
- 1058 141. Huntley RP, Sawford T, Mutowo-Meullenet P, Shypitsyna A, Bonilla C, Martin MJ, et
1059 al. The GOA database: gene Ontology annotation updates for 2015. Nucleic Acids Res.
1060 2015;43(Database issue):D1057-63. doi: 10.1093/nar/gku1113. PubMed PMID: 25378336;
1061 PubMed Central PMCID: PMCPMC4383930.
- 1062 142. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, et al.
1063 STRING v10: protein-protein interaction networks, integrated over the tree of life. Nucleic
1064 Acids Res. 2015;43(Database issue):D447-52. doi: 10.1093/nar/gku1003. PubMed PMID:
1065 25352553; PubMed Central PMCID: PMCPMC4383874.
- 1066 143. Kehl A, Hensel M. Live cell imaging of intracellular *Salmonella enterica*. Methods Mol
1067 Biol. 2015;1225:199-225. doi: 10.1007/978-1-4939-1625-2_13. PubMed PMID: 25253257.
- 1068 144. Zhang Y, Hensel M. Evaluation of nanoparticles as endocytic tracers in cellular
1069 microbiology. Nanoscale. 2013;5(19):9296-309. doi: 10.1039/c3nr01550e. PubMed PMID:
1070 23942623.
- 1071 145. Knodler LA, Nair V, Steele-Mortimer O. Quantitative assessment of cytosolic
1072 *Salmonella* in epithelial cells. PLoS One. 2014;9(1):e84681. doi:

18.11.2019

Host factors for SIF formation

1073 10.1371/journal.pone.0084681. PubMed PMID: 24400108; PubMed Central PMCID:
1074 PMC3882239.
1075 146. Lam AJ, St-Pierre F, Gong Y, Marshall JD, Cranfill PJ, Baird MA, et al. Improving
1076 FRET dynamic range with bright green and red fluorescent proteins. *Nat Methods*.
1077 2012;9(10):1005-12. doi: 10.1038/nmeth.2171. PubMed PMID: 22961245; PubMed Central
1078 PMCID: PMCPMC3461113.
1079 147. Paumet F, Le Mao J, Martin S, Galli T, David B, Blank U, et al. Soluble NSF attachment
1080 protein receptors (SNAREs) in RBL-2H3 mast cells: functional role of syntaxin 4 in exocytosis
1081 and identification of a vesicle-associated membrane protein 8-containing secretory
1082 compartment. *J Immunol*. 2000;164(11):5850-7. doi: 10.4049/jimmunol.164.11.5850. PubMed
1083 PMID: 10820264.
1084 148. Martinez-Arca S, Alberts P, Zahraoui A, Louvard D, Galli T. Role of tetanus neurotoxin
1085 insensitive vesicle-associated membrane protein (TI-VAMP) in vesicular transport mediating
1086 neurite outgrowth. *J Cell Biol*. 2000;149(4):889-900. doi: 10.1083/jcb.149.4.889. PubMed
1087 PMID: 10811829; PubMed Central PMCID: PMCPMC2174569.
1088 149. Itakura E, Kishi-Itakura C, Mizushima N. The hairpin-type tail-anchored SNARE
1089 syntaxin 17 targets to autophagosomes for fusion with endosomes/lysosomes. *Cell*.
1090 2012;151(6):1256-69. doi: 10.1016/j.cell.2012.11.001. PubMed PMID: 23217709.

18.11.2019

Host factors for SIF formation

1092 Tables

1093 **Table 1.** High-ranking trafficome hits (scoring cutoff of ≥ 8) involved in trafficking and
1094 cytoskeleton biology.

Gene symbol	Full name ¹	Localization ²
Small GTPases and interacting proteins		
<i>Arf family</i>		
ARL1	ADP-ribosylation factor-like 1	Golgi
<i>Rab family</i>		
RAB1A	RAB1A, member RAS oncogene family	ER, Golgi, EE
RAB11A	RAB11A, member RAS oncogene family	RE, vesicle, PM
<i>Ras family</i>		
RHOB	ras homolog family member B	nucleus, LE, PM
RHOT1	ras homolog family member T1	mitochondrion OM
<i>Interacting proteins</i>		
G3BP2	GTPase activating protein (SH3 domain) binding protein 2	cytoplasm
RAB3GAP2	RAB3 GTPase activating protein subunit 2 (non-catalytic)	cytoplasm
Vesicle coats and adaptors		
<i>BBSome</i>		
BBS4	Bardet-Biedl syndrome 4	CS/MTOC
<i>Clathrin coats</i>		
AGFG1	ArfGAP with FG repeats 1	nucleus, vesicle
AP2A1	adaptor-related protein complex 2, α 1 subunit	CCV, PM
AP3D1	adaptor-related protein complex 3, β 1 subunit	CCV, Golgi
CLTA/B	clathrin, light chain A and B	CCV
CLTC	clathrin, heavy chain (Hc)	CCV
GGA3	golgi-associated, γ adaptin ear containing, ARF binding protein 3	Golgi, endosome
SYNRG	synergism, γ	Golgi
<i>COP-I</i>		
ARCN1	archain 1	Golgi, vesicle
COPA/B1/B2/G1	coatamer protein complex, subunit α , β , β' , and γ	Golgi, vesicle
TMED10	transmembrane emp24-like trafficking protein 10	ER, Golgi, vesicle
<i>COP-II</i>		
SEC24D	SEC24 family member D	ER, Golgi, vesicle
<i>Retromer</i>		
VPS35	vacuolar protein sorting 35 homolog	EE, LE
Tethering factors		
<i>Exocyst complex</i>		
EXOC5	exocyst complex component 5	cytoplasm
<i>TRAPPIII complex</i>		
TRAPPC8	trafficking protein particle complex 8	Golgi

18.11.2019

Host factors for SIF formation

SNAREs

Qa-SNAREs

STX5	syntaxin 5	Golgi
STX7	syntaxin 7	EE, LE

Qb,c-SNAREs

SNAP23	synaptosomal-associated protein, 23 kDa	PM
--------	---	----

R-SNAREs

SEC22B	SEC22 vesicle trafficking protein homolog B	ER, Golgi
VAMP7	vesicle-associated membrane protein 7	ER, Golgi, LE/ lysosome, vesicle, PM

Interacting proteins

NAPA	N-ethylmaleimide-sensitive factor attachment protein, α	membrane
STXBP2	syntaxin binding protein 2	PM

Cytoskeleton and motor proteins

Kinesins

KIF1A/B/C	kinesin family member 1A, B, and C	CS
-----------	------------------------------------	----

Dyneins

DYNC1H1	dynein, cytoplasmic 1, heavy chain 1	CS
---------	--------------------------------------	----

Microtubule-associated proteins

CEP57	centrosomal protein 57 kDa	CS/MTOC, nucleus
MAP1A	microtubule-associated protein 1A	CS

Myosins

MYH10	myosin, heavy chain 10, non-muscle	CS
-------	------------------------------------	----

Actin filament membrane linkers

ANK3	ankyrin 3, node of Ranvier (ankyrin G)	CS
FLNA	filamin A, α	CS

ESCRT complexes

Adaptors

HGS	hepatocyte growth factor-regulated tyrosine kinase substrate	EE, MVB
-----	--	---------

AAA ATPase

VPS4A/B	vacuolar protein sorting 4 homolog A and B	MVB
---------	--	-----

Miscellaneous

ERGIC1	endoplasmic reticulum-golgi intermediate compartment (ERGIC) 1	ER, Golgi
SNX15	sorting nexin 15	vesicle
SORT1	sortilin 1	nucleus, ER, Golgi, endo- /lysosome
VCP	Valosin-containing protein	nucleus, ER

1095 ¹ according to NCBI Gene

1096 ² subcellular localization according to UniProt; CCV = clathrin-coated vesicle, CS =
1097 cytoskeleton, EE = early endosome, ER = endoplasmic reticulum, LE = late endosome, MTOC
1098 = microtubule organizing center, MVB = multivesicular body, OM = outer membrane, PM =
1099 plasma membrane, RE = recycling endosome

1100

18.11.2019

Host factors for SIF formation

1101 **Table 2.** Host proteins (gene symbols) identified as hits in the trafficome screen that are also
1102 present in at least one distinct SMM proteome.

8 h p.i. SMM ¹	30 min p.i. SCV ²	3 h p.i. SCV ²
AP2A1 (AP-2)	-	AP2A1
-	-	BET1 (SNARE)
-	CLTC (clathrin)	-
COPA/G1 (COPI) ³	-	-
DYNC1H1 (dynein)	DYNC1H1	-
-	ERGIC1	ERGIC1
-	ERP29	-
-	-	EXOC5
FLNA (filamin)	FLNA	FLNA
G3BP2	-	G3BP2
-	IQGAP1	IQGAP1
-	KIF5B (kinesin-1)	KIF5B
-	MAP1B	-
MYH9/10 (myosin II)	MYH9	MYH9
NAPA/ α -SNAP	-	-
RAB2A ³	RAB2A	-
-	RAB4A	-
RAB7A ³	RAB7A	-
RAB11A ³	-	-
RAB14 ³	-	-
-	SEC22B (SNARE)	-
-	SEC24C (COPII)	-
TMED10 (COPI)	-	-
VCP	-	-

1103 ¹ [32]

1104 ² [33]

1105 ³ colocalization with SMM shown by fluorescence microscopy

18.11.2019

Host factors for SIF formation

Figure legends

Figure 1. RNAi screen setup and validation. A) Intracellular phenotypes of STM under screening conditions. HeLa-LAMP1-GFP cells were infected with mCherry-labelled STM WT or *ssaV* strains and imaged live 8 h p.i. by SDCM. Presence of STM in LAMP1-positive SCV (blue arrowhead), induction of SIF formation by STM WT (white arrowhead), and lack for SIF formation by STM *ssaV* strain. Scale bar, 10 μ m. B) Controls for siRNA-mediated knockdown. HeLa-LAMP1-GFP cells were reverse transfected with scrambled AllStars siRNA or PLK1 siRNA. Scale bar, 20 μ m. C) Validation of SKIP siRNA knockdown. HeLa-LAMP1-GFP cells were reverse transfected with AllStars or SKIP siRNA. Then, RT-PCR targeting *SKIP* was performed. Depicted is the mean with standard deviation of three biological replicates ($n = 3$) each performed in triplicates. Statistical analysis was performed using Student's *t*-test and indicated as: ***, $p < 0.001$. D) SKIP knockdown as control for inhibition of SIF formation. HeLa-LAMP1-GFP cells were first reverse transfected with AllStars or SKIP siRNA. Then, cells were infected with mCherry-labelled STM WT (MOI = 15) and imaged live 1-7 h p.i. by SDCM. Blue arrowheads indicate SIF-forming or non-SIF-forming single bacteria or microcolonies, white arrowheads indicate SIF. Scale bar, 10 μ m.

Figure 2. Basic workflow of the trafficome RNAi screen. 96-well plates with clear bottoms were automatically spotted with siRNAs. HeLa-LAMP1-GFP cells were seeded for reverse transfection. After 72 h of incubation, infection with STM was performed, followed by LCI using an SDCM system with hourly intervals of imaging. Phenotypic scoring was performed using the SifScreen utility. The MATLAB-based data input mask allows the entry of well- and position-specific information on general cell behavior and *Salmonella*/SMM phenotypes and the generation of a results report.

18.11.2019

Host factors for SIF formation

1132

1133 **Figure 3. Interaction network of selected trafficome hits.** The interaction of high-ranking
1134 hits (scoring cutoff of ≥ 8 , see also Table 1) was visualized using the STRING database
1135 (confidence view). Borders delineate clusters related to the cytoskeleton (i, iii), SNAREs (ii),
1136 or COPI and clathrin-coated vesicles (iv).

1137

1138 **Figure 4. Influence of host factor silencing on SIF formation.** HeLa-LAMP1-GFP cells were
1139 not transfected (mock), or reverse transfected with siAllStars or the indicated siRNA, infected
1140 with STM WT or SPI2-deficient *ssaV* expressing mCherry as indicated, and SIF counted.
1141 Depicted are means with standard deviation for three biological replicates ($n = 3$). Statistical
1142 analysis was performed against siAllstars + WT with Student's *t*-test and indicated as: n.s., not
1143 significant; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

1144

1145 **Figure 5. RAB proteins identified by the trafficome screen colocalize with SIF and SCV.**
1146 A) Direct interaction network of RAB1B as visualized by STRING. B) and C) HeLa cells either
1147 stably transfected with LAMP1-GFP (green), or transiently transfected with LAMP1-mCherry
1148 (red) were co-transfected with plasmids encoding various RAB GTPases (RAB7A, RAB1A,
1149 RAB1B, RAB3A, RAB8A, RAB8B, RAB9A, RAB11A) fused to GFP (green) or mRuby2 (red)
1150 and then infected with STM WT expressing mCherry or GFP. Living cells were imaged from
1151 6-9 h p.i. by CLSM and images are shown as maximum intensity projections (MIP). Insets
1152 magnify structures of interest and white arrowheads indicate colocalization with SIF. Scale
1153 bars, 10 μm (overviews), 1 μm (details).

1154

18.11.2019

Host factors for SIF formation

Figure 6. SNARE proteins identified by the trafficome screen colocalize with SIF and SCV. A) Direct interaction network of STX7 as visualized by STRING. B) and C) HeLa cells either stably transfected with LAMP1-GFP (green), or transiently transfected with LAMP1-mCherry (red) were co-transfected with plasmids encoding various SNAREs (STX7, VTIB, STX8, VAMP7, VAMP8, VAMP2, VAMP3, VAMP4) fused to GFP (green) or mRuby2 (red). Infection and imaging was performed as for Figure 5. Insets magnify structures of interest and white and blue arrowheads indicate colocalization with SIF and SCV, respectively. Scale bars, 10 μ m (overviews), 1 μ m (details).

Figure 7. CLTA identified by the trafficome screen colocalizes with SIF and SCV. A) Direct interaction network of CLTA as visualized by STRING. B) HeLa cells stably transfected with LAMP1-GFP (green) were co-transfected with a plasmid encoding CLTA fused to mRuby2 (red). Infection and imaging was performed as for Figure 5. Insets magnify structures of interest and white and blue arrowheads indicate colocalization with SIF and SCV, respectively. Scale bars, 10 μ m (overviews), 1 μ m (details).

Figure 8. Newly identified interactions of intracellular STM with host factors. Depicted are central eukaryotic endomembrane organelles possibly playing a role in the newly identified interplays of host factors with SIF. Magnifications show the interactions of clathrin (i), late secretory and/or recycling-related RAB3A, RAB8A/B, and VAMP2/3/4 (ii), late endo-/lysosomal VTI1B, STX8, and VAMP8 (iii), and early secretory RAB1A/B (iv) with other host factors added as discussed in the text. Solid lines represent interactions identified here or otherwise known, dashed lines represent putative interactions. COP, coat protein complex; EE, early endosome; ER, endoplasmic reticulum; LE, late endosome; Lys, lysosome; RE, recycling

18.11.2019

Host factors for SIF formation

1179 endosome; SCV, *Salmonella*-containing vacuole; SIF, *Salmonella*-induced filaments; STM, *S.*

1180 Typhimurium; SV, secretory vesicle.

1181

1182 **Suppl. Material**

1183 **Suppl. Tables**

1184 **Table S1.** Full list of the 496 host factors targeted in the siRNA screen including gene symbols,
1185 NCBI Gene IDs and accession numbers, full name, and aliases.

1186

1187 **Table S2.** Summary of the scoring of the executed siRNA screen with a full list of the
1188 trafficome, a list with hits only, and lists with low-, mid-, and high-ranking hits (scoring of 1-
1189 4, 5-7, or ≥ 8 , respectively).

1190

1191 **Table S3.** Requirements of an RNAi screen with LCI for the identification of host factors
1192 involved in SIF formation by intracellular *Salmonella*.

Type of prerequisite	Prerequisite	Necessity/Solution
biological	• label for structure of interest • <i>Salmonella</i> controls for structure of interest	• LAMP1 as marker of SIF • WT: SIF-positive, <i>ssaV</i>: SIF-negative
technical	• fast and gentle microscopic system • multiwell-compatible 37 °C incubation • multiwell-compatible sufficient magnification • autofocus system • autofocus-compatible multiwell plates	• spinning disk confocal microscope (Zeiss) • box incubation (custom-made) • LD 40x/0.6 objective (+ bottom thickness correction collar, Zeiss) • infrared-based system (Definite Focus, Zeiss) • 96 Well Clear Bottom Black Cell Culture Microplates (0.5 mm bottom, Corning)
siRNA-related	• harmlessness of transfection reagent • harmlessness of siRNA as reagent • general efficacy of siRNA • phenotype-interfering siRNA	• mock transfection control • scrambled AllStars siRNA control • lethal PLK1 siRNA control • SIF-abolishing SKIP siRNA • RT-PCR of SKIP knockdown

18.11.2019

Host factors for SIF formation

	• control of phenotype-interfering siRNA	
data analysis	• primary analysis	• manual visual inspection
	• collection of analyzed data	• SifScreen utility

1193

1194 **Table S4.** Characteristics of microscopy systems with respect to LCI and RNAi screening.

System	Advantage	Disadvantage
WFM	• fast	• lowest resolution (without deconvolution)
CLSM	• gentle to cells	• slow
	• highest resolution	• high local illumination, photo-damage
SDCM	• high resolution	• lower resolution as CLSM (due to camera)
	• fast	
	• gentle to cells	

1195

1196 **Table S5.** Suitability of various types of multi-well plates for screening with LD 40x objective
1197 and IRF.

Type	Bottom thickness in mm	Applicability	Representative suppliers
standard plastic	1	• no, bottom variation too high	Corning, Greiner, Thermo/Nunc, TPP
glass	0.17	• no, otherwise incompatible	Corning, Greiner, MatTek, Thermo/Nunc
cyclic olefin/synthetic	0.19	• no, otherwise incompatible	Corning, Greiner
special plastic	0.5	• yes	Corning

1198

1199 **Table S6.** Bacterial strains and plasmids used in this study.

Designation	Relevant genotype	Source/Reference
STM strains		
NCTC 12023	wild type	lab collection
P2D6	<i>ssaV::mTn5</i>	[4]
MvP1897	$\Delta phoN::P_{EM7}::sfGFP\ aph$	this study
Plasmids		
p3099	$P_{CMV}::eGFP::RAB8A$	This study
p3101	$P_{CMV}::eGFP::RAB9A$	This study
p3451	$P_{CMV}::hLAMP1::mCherry$	[18]
p4080	$P_{CMV}::mRuby2-C1$	This study
p4081	$P_{CMV}::mRuby2-N1$	This study

18.11.2019

Host factors for SIF formation

p4094	P _{CMV} ::mRuby2::STX8	This study
p4122	P _{CMV} ::mRuby2::RAB1B	This study
p4209	P _{CMV} ::mRuby2::RAB8B	This study
p4210	P _{CMV} ::mRuby2::RAB3A	This study
p4232	P _{CMV} ::mRuby2::CLTA	This study
p4250	P _{CMV} ::mRuby2::RAB1A	This study
pcDNA3-mRuby2	P _{CMV} ::mRuby2	Addgene (#40260), [146]
pEGFP VAMP3	P _{CMV} ::eGFP::VAMP3	Thierry Galli, Paris, Addgene (#42310), [147]
pEGFP VAMP7	P _{CMV} ::eGFP::VAMP7	Thierry Galli, Paris, Addgene (#42316), [148]
pEGFP VAMP8	P _{CMV} ::eGFP::VAMP8	Thierry Galli, Paris, Addgene (#42311), [147]
pEGFP-C1	P _{CMV} ::EGFP-C1	Clontech
pEGFP-C3	P _{CMV} ::EGFP-C3	Clontech
pEGFP-N1	P _{CMV} ::EGFP-N1	Clontech
pEGFP-Vamp4	P _{CMV} ::VAMP4::eGFP	Wanjin Hong, Singapur
pENTR223_RAB1A	RAB1A	DNASU GTPases (HsCD00509534)
pENTR223_RAB1B	RAB1B	DNASU GTPases (HsCD00509534)
pENTR223_RAB3A	RAB3A	DNASU GTPases (HsCD00507538)
pENTR223_RAB8B	RAB8B	DNASU GTPases (HsCD00288320)
pENTR223_STX8	STX8	DNASU 16967 (HsCD00507664)
pFPV-mCherry	P _{rpsM} ::mCherry	Addgene (#20956), [34]
pGL-Rab7 wt	P _{SV40} ::RAB7::eGFP	Martin Aepfelbacher, Hamburg
pLenti Vamp2	VAMP2::pHtomato	Yulong Li, Beijing
pHtomato		
pMRXIP GFP-Stx7	P _{CMV} ::GFP::STX7	Addgene (#45921), [149]
pMRXIP GFP-Vti1b	P _{CMV} ::GFP::VTI1B	Addgene (#45922), [149]

1200

1201 **Table S7.** Individual siRNA information used for validation.

Gene symbol	siRNA name	Catalog no.	Target sequence
---	AllStars	SI03650318	proprietary
HGS	Hs_HGS_6	SI02659650	GCACGTCTTTCCAGAATTCAA
PLK1	Hs_PLK1_7	SI02223844	CGCGGGCAAGATTGTGCCTAA
RAB1A	Hs_RAB1A_9	SI02662716	AACATAGAGTTAGACGGGAA
RAB7A	Hs_RAB7A_2	SI00066395	TCCCGTTAGATCAGCATTCTA
RAB11A	Hs_RAB11A_7	SI02663206	AAGAGCGATATCGAGCTATAA
SKIP/ PLEKHM2	custom	custom	AAAACGAAGAGCAGCTGTTCA
STX5	Hs_STX5A_4	SI00048636	CAGTGGAAATTGAAGAGCTAA
STX7	Hs_STX7_7	SI03064159	CAGAGGATGACCTCCGTCTTA
VAMP7	Hs_SYBL1_7	SI04212453	TAGGGCAATCGTGTGCTAAT
VAMP8	Hs_VAMP8_2	SI02652993	CCGACTAGGCGAATTCCTTA

18.11.2019

Host factors for SIF formation

VCP	Hs_VCP_7	SI03019730	AACAGCCATTCTCAAACAGAA
VPS11	Hs_VPS11_6	SI02778167	CAGCAATATATCCGAACCATT

1202

1203 **Table S8.** Oligonucleotides in this study.

Targeted gene	Designation	Sequence 5'-3'
Used for RT-PCR		
GAPDH	Hs-GAPDH-qPCR-For2	TGCACCACCAACTGCTTAGC
	Hs-GAPDH-qPCR-Rev2	GGCATGGACTGTGGTCATGAG
HGS	Hs-HGS-qPCR-For	CTCCTGTTGGAGACAGATTGGG
	Hs-HGS-qPCR-Rev	GTGTGGGTTCTTGTCTGTTGAC
RAB1A	Hs-RAB1A-qPCR-For	AGATTAAAAAGCGAATGGGTCCC
	Hs-RAB1A-qPCR-Rev	GCTTGACTGGAGTGCTCTGAAT
RAB7A	Hs-RAB7A-qPCR-For	TACAAAGCCACAATAGGAGCTG
	Hs-RAB7A-qPCR-Rev	GCAGTCTGCACCTCTGTAGAAG
RAB11A	Hs-RAB11A-qPCR-For	CAACAAGAAGCATCCAGGTTGA
	Hs-RAB11A-qPCR-Rev	GCACCTACAGCTCCACGATAAT
SKIP/ PLEKHM2	Hs-SKIP-qPCR-For	TAGAGTTCATTTCGTTTCGAGCTG
	Hs-SKIP-qPCR-Rev	AAGGCGGTCTTCAAAGTCCAG
STX5	Hs-STX5-qPCR-For	AAAGCGCAAGTCCCTCTTTGA
	Hs-STX5-qPCR-Rev	TGAGCAATTTGTTTGTTGAGGC
STX7	Hs-STX7-qPCR-For	GGCCCAGAGGATCTCTTCTAA
	Hs-STX7-qPCR-Rev	ACTGTTGCCTCAATTCAGGTG
VAMP7	Hs-VAMP7-qPCR-For	GAGGTTCCAGACTACTTACGGT
	Hs-VAMP7-qPCR-Rev	GACACTTGAGAACTCGCTATTCA
VAMP8	Hs-VAMP8-qPCR-For	TGTGCGGAACCTGCAAAGT
	Hs-VAMP8-qPCR-Rev	CTTCTGCGATGTCGTCTTGAA
VCP	Hs-VCP-qPCR-For	CAAACAGAAGAACCGTCCCAA
	Hs-VCP-qPCR-Rev	TCACCTCGGAACAACTGCAAT
VPS11	Hs-VPS11-qPCR-For	CAAGCCTACAACTACGGGTG
	Hs-VPS11-qPCR-Rev	GAGTGCAGAGTGGATTGCCA
Used for plasmid construction		
EGFP	Vf-pEGFP-C1	TCCGGACTCAGATCTCGAGCTCA
	Vr-pEGFP-C1	GGTGGCGACCGGTAGCGC
EGFP	Vf-pEGFP-N1	AGTAAAGCGGCCGCGACT
	Vr-pEGFP-N1	GGTGGCGACCGGTGGATC
RAB1A	1f-Rab1A	CTTCGAATTCTGCAGTCGACATGTCCAG CATGAATCCCGAATA
	1r-Rab1A	TCTAGATCCGGTGGATCCTCAGCAGCAA CCTCCACCTGAC
RAB1B	1f-Rab1B	CTTCGAATTCTGCAGTCGACATGAACCC CGAATATGACTACCTGTTT
	1r-Rab1B	TCTAGATCCGGTGGATCCTCAGCAACAG CCACCGCCAGC
RAB3A	1f-Rab3A	CTTCGAATTCTGCAGTCGACATGGCATC CGCCACAGACTC
	1r-Rab3A	TCTAGATCCGGTGGATCCTCAGCAGGCG CAGTCCTGGTG

18.11.2019

Host factors for SIF formation

RAB8B	1f-Rab8B	CTTCGAATTCTGCAGTCGACATGGCGAA GACGTACGAT
	1r-Rab8B	TCTAGATCCGGTGGATCCTCAGCAAAGT AGCGAGCAACG
mRuby2	1f-pcDNA3-mRu-C1	GCGCTACCGGTCGCCACCATGGTGTCTA AGGGCGAAGAG
	1r-pcDNA3-mRu-C1	TCGAGATCTGAGTCCGGACTTGTACAGC TCGTCCATC
mRuby2	1f-pcDNA3-mRu-N1	GATCCACCGGTCGCCACCATGGTGTCTA AGGGCGAAGAG
	1r-pcDNA3-mRu-N1	AGTCGCGGCCGCTTTACTTTACTTGTAC AGCTCGTCCATC
mRuby2	Vf-pmRuby2-C1	GGATCCACCGGATCTAGATAAC
	Vr-pmRuby2-C1	GTCGACTGCAGAATTCGAAG
mRuby2	Vf-pmRuby2-N1	CCGCGGGCCCGGGATCCA
	Vr-pmRuby2-N1	CTGCAGAATTCGAAGCTTGAGCTCGAGA
STX8	1f-STX8-C1	CTTCGAATTCTGCAGTCGACATGGCACC GGACCCCTGGT
	1r-STX8-C1	GTTATCTAGATCCGGTGGATCCGTTGGT CGGCCAGACTGCAA

1204

18.11.2019

Host factors for SIF formation

1205 **Suppl. Figure legends**

1206

1207 **Figure S1. Validation of host factor siRNA silencing.** HeLa-LAMP1-GFP cells were reverse
1208 transfected with siAllStars or the indicated siRNA. Total RNA was extracted, mRNA reverse
1209 transcribed, and the generated cDNA was used in RT-PCR. Depicted are means with standard
1210 deviation for three biological replicates ($n = 3$) each performed in triplicates. Statistical analysis
1211 was performed against siAllstars with Student's t -test and indicated as: ***, $p < 0.001$.

1212

1213 **Suppl. Movie captions**

1214 **Movie 1. Time-lapse acquisition of infected siAllStars-treated cells.** The movie corresponds
1215 to Figure 1D.

1216

1217 **Movie 2. Time-lapse acquisition of infected siSKIP-treated cells.** The movie corresponds to
1218 Figure 1D.




Figure 1




Figure 2




Figure 3





Figure 4







Figure 6

A



B





Figure 7





Figure 8