



Review

Forensic implications of genetic analyses from degraded DNA—A review

Reza Alaeddini^a, Simon J. Walsh^{b,*}, Ali Abbas^a^a School of Chemical and Biomolecular Engineering, University of Sydney, Sydney, Australia^b Forensic and Data Centres, Australian Federal Police, Canberra, Australia

ARTICLE INFO

Article history:

Received 28 August 2008

Received in revised form 30 August 2009

Accepted 11 September 2009

Keywords:

DNA degradation

Forensics

Cell death

Deoxyribonuclease

PCR

ABSTRACT

Forensic DNA identification techniques are principally based on determination of the size or sequence of desired PCR products. The fragmentation of DNA templates or the structural modifications that can occur during the decomposition process can impact the outcomes of the analytical procedures. This study reviews the pathways involved in cell death and DNA decomposition and the subsequent difficulties these present in DNA analysis of degraded samples.

© 2009 Elsevier Ireland Ltd. All rights reserved.

Contents

1. Introduction	148
2. Cell death and post-mortem decomposition	149
3. Cell death and DNA degradation processes	149
3.1. DNA enzymatic degradation	150
3.2. Nonenzymatic DNA degradation processes	150
3.2.1. Hydrolytic reactions	150
3.2.2. DNA crosslinkages	151
3.2.3. Oxidative reactions	151
3.2.4. Radiation	151
4. PCR amplification of degraded DNA	151
4.1. Failure of amplification	151
4.2. Preferential amplification	153
4.3. PCR miscoding lesions	153
4.4. Jumping PCR phenomenon	153
4.5. PCR inhibition versus DNA structural modifications	153
5. DNA repair strategies	153
6. DNA preservation	153
7. Quantification of DNA fragments	154
8. Practical applications of DNA degradation	154
9. Conclusion	154
Acknowledgements	154
References	154

1. Introduction

DNA is a negatively charged polynucleotide that usually exists in the form of double-helix. The two strands linked with

* Corresponding author. Fax: +61 2 6223 3270.

E-mail address: SimonJoseph.Walsh@afp.gov.au (S.J. Walsh).

hydrogenic bonds running in opposite directions. Each single strand DNA chain consists of a backbone made up of deoxyribose sugar molecules linked together at the 5' and 3' positions by phosphate ester groups. To accomplish the nucleotide structure a purine or pyrimidine base is attached as a C–N bond to the deoxyribose sugar molecules at the 1' position. In natural conditions DNA is a hydrated macromolecule, with 8–10 tightly bound water molecules per nucleotide residue [1]. As a highly reactive chemical residue, DNA is the target of several physical agents and chemical reactions. DNA analysis is widely used for human identification purposes after various kinds of tragic events such as crimes, mass disasters and terrorist attacks. In these circumstances, forensic investigation of biological items often requires the analysis of highly degraded material.

Polymerase chain reaction (PCR) is currently the method of choice for *in vitro* amplification of DNA molecules ahead of genetic analysis. During the decomposition process, the DNA templates can become highly fragmented or chemically modified, therefore reducing the yield of intact target fragments and consequently leading to failure or poor PCR amplification or mistyping of the target loci. Awareness of DNA degradation pathways and mechanisms would assist in choosing the most appropriate target samples and in designing improved techniques for sampling, storage, purification and probable repair. Moreover consideration of the original phenomena that drives in DNA degradation could assist in better interpretation of the results.

This study reviews the pathways associated with DNA degradation and their implications for forensic DNA profiling.

2. Cell death and post-mortem decomposition

The advent of death in higher order organisms and cessation of blood perfusion to the tissue leads to cellular oxygen and nutritional deprivation and consequently a drop in cellular adenosine tri-phosphate (ATP) production. The shift of ATP production reactions from oxidative phosphorylation to anaerobic reactions leads to accumulation of acidic by-products. Primarily it is the cells with higher energy demands or higher intracellular enzyme activities that are affected, however, these events eventually effect all cells in the body [2]. Since they require large amounts of oxygen to provide the energy to support the specialized cellular functions, the effects are most significant in well differentiated tissues such as heart, kidney, and brain. Different sections of the same organ can display different resistance against perfusion deficiency. In heart tissues for example necrosis occurs first in the subendocardial region following 20–40 min of ischemia, extends into the midmyocardium by 1–3 h, and often approaches transmural stage within 24 h [3,4].

Depending on several internal and external factors such as the nature of death signal, the tissue characteristics and the developmental stage of the tissues, cells would undergo one of two different death patterns; apoptosis or necrosis [5]. Apoptosis features an energy dependent programmed cell death, characterized by condensation of cytoplasm, loss of plasma membrane microvilli, segmentation of the nucleus, and extensive degradation of chromosomal DNA into oligomers of about 180 bp through activation of endogenous endonucleases [3]. Intracellular ATP levels remain unaltered until the very end of apoptosis process [6].

Alternatively as a cellular metabolic collapse with homeostatic breakdown, necrosis is a passive energy independent degenerative phenomenon. Necrosis is characterized by increased cell volume, swelling of cytoplasmic organelles, and, whilst the nuclei appear intact, there is chromatin condensation. This is followed by rupture of the membranes, organelle breakdown and spillage of lysosomal enzymes. Necrotic cells generate a random degradation pattern of DNA. Microscopic changes towards necrosis occur faster in higher

ambient temperatures [7]. Necrosis is typically induced by extremes in the external environmental conditions of the cell such as hypoxia, or through the action of membrane active toxicants and respiratory poisons such as cyanide [8]. Despite the existence of distinctive microscopic and biochemical aetiology, accurate discrimination between apoptosis and necrosis requires an integrated morphological and biochemical approach [9].

Increased membrane permeability and rupture of the cell membrane represent important steps in the necrotic process [10]. The onset and progression of ultrastructural changes in necrotic organs varies in different tissue. In liver, for example, the ultrastructural changes appear in 1 h whereas in skeletal muscles the changes take about 24 h post-mortem to appear [11]. Among several factors involved in cell death process the level of intracellular ATP determines whether a cell will die by apoptosis or necrosis [12,13].

At the molecular stage under both necrotic and apoptotic conditions the movement of Ca^{2+} plays a practically significant role in the pathogenesis of cell injury and cell death. Intracellular Ca^{2+} often increases within seconds or minutes following the injury [14]. Calcium channel blockers such as Verapamil reduce the amount of myocardial necrosis [15].

In physiological conditions the extracellular fluid Ca^{2+} is maintained at about 1.2 mM, whereas the cytoplasmic or nuclear and mitochondrial matrices Ca^{2+} concentrations are maintained at a very lower level about 100 nM. Such a difference between the compartments is maintained via the plasma membrane Ca^{2+} pumps and the $\text{Na}^+/\text{Ca}^{2+}$ exchangers [16]. During the first stages of cell death, Ca^{2+} released from the endoplasmic reticulum (ER) is taken up by mitochondria, however by the prolongation of the death phenomenon sustained release of Ca^{2+} from the ER stores combined with additional stress signals and the induction of mitochondrial membrane permeability may initiate Ca^{2+} -dependent forms of cell death [17,18]. Fig. 1 illustrates the pathways engaged in cell death and the progress towards nuclease activation in glucose and oxygen deprivation conditions.

3. Cell death and DNA degradation processes

Full blown cell death is associated with activation of different classes of various intracellular enzymes including lipases, nucleases, and different classes of proteases. Lysosomes are ubiquitous acidic vacuoles containing several hydrolytic enzymes. Most are active in acidic pH and capable of breaking down biomolecules [19]. The removal of histone proteins (the major proteins involved in chromatin structure) by lysosomal proteases would facilitate the process of DNA cleavage by endonucleases. Post-mortem DNA fragments are subject to degradation by endogenous nucleases released by host cells or exogenous nucleases released by microorganisms and environmental invertebrates. Moreover spontaneous degradation by hydrolysis and oxidation will further modify DNA structure at a much slower speed [20,21]. The apoptosis pathway is dependent mostly upon activation of a group of protease enzymes called as caspases [22,23]. Caspases, Granzyme A and serine proteases trigger the different intracellular pathways leading to nuclease activation [24].

The increase in free cytosolic calcium (Ca^{2+}) results in activation of membrane-bound phospholipases, which degrade membrane phospholipids and cause widespread disruption of membranes [25] and therefore release of enzymes.

A fundamental knowledge on the types of DNA damage encountered in forensic stains that would lead to amplification failure has not been yet provided. Quantitation of damaged DNA fragments is achievable by customised quantitative-PCR (Q-PCR) strategies [26].




Fig. 1. Biochemical pathways initiated by oxygen, serum and glucose deprivation leading to nuclease activation and cell death. AIF: apoptosis inducing factor.

3.1. DNA enzymatic degradation

Endogenous nucleases feature amongst the first agents to initiate the process of DNA fragmentation in the post-mortem period [27]. Based on the periodicities of DNA folding in the chromatin structure [28] provided the nucleosomal bound histone structure remains intact, DNA can undergo a series of successive steps of fragmentation. High molecular weight (HMW) fragments (300, 50 kb) can result from endonucleolytic cleavage of interphase chromosomes at the nuclease-sensitive sites that reside in chromatin fibres as a result of their folding into loop (mean size of 50 kb) and rosette (mean size of 300 kb) structures, respectively [28]. Ultimately, the ladder-like pattern of oligonucleosomal sized fragments noticeable on conventional electrophoresis yields are produced as a hallmark of apoptosis [29,30]. The digestion of chromatin proteins by lysosomal proteases would in turn facilitate the process of random digestion of DNA by endonucleases. This would produce DNA smears on gel electrophoresis most notably in post-mortem forensic cases. Several factors affect the rate of the fragmentation (and hence the detectability of the residual DNA) including the expression level of the enzymes, availability of cations and solution for the reactions, temperature and pH levels [31].

In post-mortem brain tissue, for example, the chromatin structure is properly preserved for at least 30 h after death [32]. Under favorable conditions such as low temperatures or where tissue dependent natural protective measures exist, nuclease enzymes may become inactive before they fully break down long strands of DNA into shorter, less informative fragments. The HMW fragments are produced in the presence of Mg^{2+} as the sole divalent cation [33], whereas oligonucleosomal fragments are produced when both Mg^{2+} and Ca^{2+} are present in the digestion buffer [34,35]. Ca^{2+}/Mg^{2+} DNase has been reported as a major feature in apoptosis [36]. Spleen, liver, kidney and thymus have expressed high levels of this enzyme whilst lung, brain, heart and testis showed little activity. In cell types lacking Ca^{2+} dependent activity, different DNA fragmentation takes place through the action of other endonuclease groups [37].

DNase activity in some tissues and conditions such as empyema might be so slight or absent that the DNA from the dead cells forms a solidified shape requiring external DNase injections to break it down [38]. Physiological concentrations of Zn^{2+} are also known to inhibit both DNA fragmentation and Ca^{2+}/Mg^{2+} nuclease activity [39].

Low molecular weight (LMW) DNA breaks often contain 3'-OH ends, which can be labeled by terminal deoxynucleotidyl transferase dUTP mediated nick end labeling (TUNEL) [40].

Table 1 summarises the endonuclease enzyme activities involved in cell death process.

The rupture of cell membranes during the later stages of cell death leads to the release of nutrient-rich fluids, which encourage the growth of environmental microorganisms and therefore further degradation of macromolecules [27]. More than 70% of soil microorganisms contain nuclease enzymes [48]. The pre-dominating soil microorganisms are able to decompose nucleic acids and most of their degradation products [49]. Factors such as the availability of the appropriate nutrients, pH, electron acceptors and donors, ionic concentration, and the absence of toxic metal ions will determine which microbial communities will colonize the post-mortem tissue [50].

Few environmental situations can properly exclude the microbial activity. Incorporation of molecules into bio-minerals might be one of those preservative conditions [51] as it happens in extracellular bone matrix [52]. The availability of liquid media increases the vulnerability to degradation. Such media is available during the very first stages of post-mortal period by the tissue matrix or later on through the accumulation of bacterial liquid products.

3.2. Nonenzymatic DNA degradation processes

The enzymatic reactions are followed by the much slower but persistent nonenzymatic or spontaneous DNA degradation processes [53]. DNA Phosphodiester bonds are highly resistant to nonenzymatic hydrolysis [54].

3.2.1. Hydrolytic reactions

The glycosidic base-sugar bond is the most susceptible bond to cleavage within the polynucleotide strand. Hence, it is the main target of hydrolytic attacks, resulting in base loss. In an alkaline hydrolysis reaction depurination is then followed by a primarily β elimination reaction which rapidly breaks the DNA strand at the 3'-phosphodiester bond of the apurinic sugar. In a medium with ionic concentrations similar to intracellular composition (37 °C and pH 7.21), the apurinic site has an estimated half life of between 288 and 335 h [55]. Primary amines (such as glycine, lysine, histidine and arginine), tris [56], Mg^{2+} ions, basic proteins [57] or aldehyde reagents [58] all can increase the rate of fragmentation. For

Table 1

Endonuclease activities involved in cell death. ND = not determined.

Cation dependence	DNA ends	Name	pH	Tissue distribution	Subcellular location	Original inhibitor	Activation	Ref.
Mg ²⁺ dependent	3' OH 5'P	DFF40 ^a	Neutral	Pancreas, spleen, prostate and ovary	Cytoplasm, nucleus	DFF45	Activated caspase-3	[44]
		GAAD ^b	Neutral	Natural killer cells	Cytotoxic T lymphocytes	Lysosome	IGAAD	[45]
		Endo G ^c	Neutral	Brain	Mitochondrion	ND	Granzyme A	[46]
Ca ²⁺ / Mg ²⁺ dependent	3' OH 5'P	Exo G ^d	6	ND	Mitochondrion	ND	ND	[47]
		Various ^e	Neutral ^e	Wide ^e	Nucleus	ND	ND	[31]
								[41]
Cation independent	3' P 5'OH	Various ^f	Acidic	Various	Lysosome, nucleus	ND ^f	ND ^f	[24]

^a DNA fragmentation factor 40 (caspase-activated nuclease), weakly activation by Ca²⁺ and Mg²⁺, formation of double strand DNA breaks, internucleosomal breakage.^b Granzyme A-activated DNase, NM₂₃-H₁, formation of single strand DNA nicks, high molecular weight breakage.^c Endonuclease G, internucleosomal breakage.^d Exonuclease G, 5'–3' exonuclease activity, combined endo and exonuclease activity.^e Internucleosomal breakage, inhibition reported by zinc, aurointricarboxylic acid and DR396 [41,42]. DNase X is mainly expressed in skeletal muscles and myocardium. Human DNase (gamma) is mainly distributed in macrophages, liver, spleen, and some other organs; pronounced expression of DNase₁L₂ was found in brain, lung, and placenta and it's an acidic exonuclease [43]. DNase₁L₂ was found predominantly in the cytoplasm [41].^f DNase II (alpha), DNase II (beta), L-DNase II. No preference in breakage pattern. L-DNase II is synthesized as LEI (leukocyte elastase inhibitor). The proteolytic cleavage of LEI by elastase and other serine proteases transforms it into the active form.

example, the polyamine putrescine (a bacterial metabolism product) at a concentration of 0.01 M causes a 25-fold increase in the rate of chain breakage at apurinic sites [58].

Being normally an acid catalysed reaction in pH < 6.0, the rate of depurination decreases with increasing pH and increases in solvents of low ionic strength. Above pH 6.0, the pH dependency of the reaction decreases and depurination proceeds mostly by a reaction that is independent to pH [59]. The reaction rate is also temperature dependant so it may proceed at a lower rate in conditions that approximate physiological temperatures. Theoretically for an 800 bp DNA fragment in 15 °C it takes 5000–10,000 years to achieve total degradation [58,60]. Pyrimidines are more stable, being released at 5% the rate of the purines [20]. In physiological ionic concentrations, at 15 °C, it would take approximately 100,000 years to destroy all human DNA by hydrolytic damage [27,20].

In addition, DNA bases with secondary amino groups such as adenine, cytosine, 5-methylcytosine and guanine undergo the hydrolytic loss of their amino groups (deamination) resulting in the generation of hypoxanthine, uracil, thymine, and xanthine, respectively. Cytosine residues are particularly prone to deamination and uracil formation; however the reaction is slow at 37 °C with each cytosine half life of about 30,000 years [20].

Whilst protected from hydrolytic damage, desiccated tissues are still prone to oxidation [61].

3.2.2. DNA crosslinkages

Abasic sites may proceed to DNA crosslinks between DNA and proteins [62] or between the ring-opened sugar of the abasic sugar and an amino group located on the opposite strand [63]. The crosslinking reaction between the two DNA strands is temperature dependent and is slower in low temperatures. It has been observed at a frequency of 1 crosslinkage for 6 apurinic sites after 85 days of storage at 4 °C [64]. Crosslinks can even be observed directly by electron microscopy in DNA preparations [61]. DNA strands would also react through their bases with reducing sugars in a nonenzymatic glycation reaction [65] followed by the generation of abasic sites [66].

3.2.3. Oxidative reactions

Oxygen-derived species such as superoxide radical (O₂^{•-}) and H₂O₂ are generated through ionizing radiation or the metabolic activities of aerobic microorganisms that colonize post-mortem tissue. Ionizing radiation can also produce reactive hydroxyl radicals in cells and tissues by interacting with cellular water.

Pyrimidines (in particular thymine) are more sensitive to oxidative damage than purines [67]. Oxidative damage mostly includes modifications of sugar residues, conversion of cytosine and thymine to hydantions, removal of bases and crosslinkages. During the decay process cytosine and thymine may be modified into Hydantoins by oxidation. This will block PCR amplification [68].

3.2.4. Radiation

Radiation can produce a variety of lesions including oxidative damages, single- and double-strand breaks, base modifications, destruction of sugars, intra and interstrands crosslinks and formation of dimers [69].

Fig. 2 illustrates the most common sites of oxidation, hydrolytic and nuclease enzymatic attacks on DNA fragments.

4. PCR amplification of degraded DNA

Forensic biology samples may contain highly fragmented DNA molecules replete with the various forms of damage summarised above. The current protocols in forensic analysis of human remains are based on size or sequence analysis [70,71] of PCR products. PCR based protocols although highly effective may encounter complications through the low copy numbers of the template or the modifications imposed on the template during the decay process. A diploid human cell contains ~6.6 pg of genomic DNA. A template DNA concentration < 100 pg genomic DNA (about 15–17 diploid copies of nuclear DNA markers such as autosomal STRs) is considered as low copy number [70].

Few practical studies have been undertaken on the underlying causes of artefacts encountered during amplification of degraded forensic specimens. In one study performed on selected aged forensic samples the frequency of artefacts including base substitutions and ambiguities was calculated as being at least 30-fold higher than the controls, with DNA oxidation products among the most abundant base modifications in aged samples [72]. The three main complications observed following PCR of degraded DNA samples are the failure of amplification, preferential amplification and miscoding lesions.

4.1. Failure of amplification

STR multiplexes usually work optimally with 1 ng DNA template with 28–30 amplification cycles [73]. One nanogram of DNA is approximately equal to 660 copies of genomic DNA. An increased incidence of PCR failures was observed when the starting




Fig. 2. Oxidation, hydrolytic and enzymatic attacks associated with DNA degradation. Redrawn with modifications from [20]. A: adenine, G: guanine, C: cytosine, T: thymine.

DNA amount was less than 60 copies [74]. Commercial multiplex kits used in forensic DNA typing amplify fragments in the size range of 100–450 bp [75]. Multiplexing the PCR reaction reduces the amount of sample material necessary for analysis and minimizes the experimental time and costs. Extra doses of *Thermus aquaticus* (Taq) polymerase and extra PCR cycles have been suggested to overcome the low profile amplification of low copy number templates [76].

In degraded samples there is a higher chance that shorter amplicons will be amplified compared to longer amplicons [77]. This is due to the higher likelihood that longer products will be degraded during the decay process. Therefore in multiplex genotyping profiles with a wide range of amplicon sizes the peak positions produce a curve in which the peak heights are inversely proportional to the amplicon lengths [78].

As amplification and profiling of shorter fragments is more successful in degraded templates, the newer versions of primer sets re-position the primers as close as possible to the repeat motif to reduce the overall amplicon lengths. These reduced sized targets are referred to as Mini-STRs [79]. The smaller products would also serve as better templates for analysis by alternative technologies such as time-of-flight mass spectrometry [80,81] and rapid microchannel electrophoretic separations [82]. This strategy reportedly reduced the required amount of DNA for amplification of genomic templates up to 100 pg/25 μ l [78].

In recent years forensic scientists have focused increased attention on single nucleotide polymorphisms (SNP) analysis. A relatively small array of 50 SNP loci gives likelihood ratios equivalent to approximately 12 STRs [83]. Short SNP fragments can be designed to analyze the degraded samples. Nevertheless there is a greater chance of base alterations in aged post-mortem samples [84].

Mitochondrial DNA (mtDNA) analysis is considered as a good alternative to genomic DNA identification measures, particularly in circumstances where there is an expectation that the template will be degraded or present in low copy numbers. MtDNA exists within cytoplasmic mitochondria as a separate small circular genome of 16,569 bp. Every mitochondrion may contain more than one copy of mtDNA [85]. Depending on cell type mammalian cells host between 80–680 mitochondria and 200–1700 mtDNA genome copies. This is equivalent to (on average) about 2.6 mtDNA genomes per mitochondrion [86]. The high copy number is one feature that makes mtDNA especially useful for investigations of human remains containing highly degraded DNA. MtDNA has

become the method of choice for working on skeletal remains such as bones [87–92] and teeth [93–95] as well as fingernails [96] and shed hairs [97] where nuclear DNA testing is typically less successful. Sequence analysis of hypervariable regions is the preferred mechanism for analysis of mtDNA variability. Compared to STR analysis this is a time-consuming process, and due to the haploid, non-Mendelian nature of mtDNA inheritance, the data are less powerful for identification purposes than a full 13-locus STR match [80].

For loci such as those targeted in forensic applications (shorter than 900 bp of 45–56% GC content), most DNA polymerases display similar amplification efficiencies. Hot-start modifications for appropriately pure templates did not appear to greatly influence amplification efficiency [98]. Although amplification efficiency increases as PCR target size reduces, the reduction of the PCR target size may also increase the chances of amplifying contaminating DNA, especially in highly degraded DNA specimens. Non-target STR products are amplified from highly degraded DNA samples [99]. Given the sensitivity required to work on ancient and forensic samples, Poinar proposed criteria of authenticity to ensure reproducibility within and between ancient and forensic laboratories [100]. Briefly the criteria consist of physical isolation of work area, negative control amplifications, testing the larger compared to shorter fragments amplification, quantifying the copy number of the DNA target, establishing the reproducibility of the results, cloning and sequencing of PCR products and independent replication of the experiments.

Whilst these technical complications associated with profiling degraded targets increase the potential for error, the increased use of DNA profiling has brought about a parallel improvement in interpretation methodology so that the probability of technical artifacts affecting the designation can be considered [101].

In low copy number samples repeated extractions and amplifications are recommended to authenticate the results [102]. Other analytical approaches recommended include reducing PCR volume [103], Increasing PCR cycle number [101], nested PCR [104], Whole genome amplification prior to the PCR [105], post-amplification removal of the interfering ions [106], and longer injection time at the outset of capillary electrophoresis [103]. The success rate of amplification can also be increased by using higher primer concentrations, however, the possibility of chimera or artifact formation may also increase concomitantly [107]. Whilst low copy number typing is appropriate for identification of missing persons and human remains and for developing investigative

leads, future developments are required to overcome its limitations in other applications [108].

Occurrence of artifacts discussed here could be reduced if quantitation of the sample indicated that template DNA contained fragments several hundred base pairs to initiate the PCR [27]. Such quantitation of template molecules can be achieved by using competitive PCR [109] and more recently, by using real-time quantitative PCR [110].

4.2. Preferential amplification

Subsequent to the degradation process the preferential amplification condition could happen in heterozygote individuals when one of the two alleles fail to amplify properly. Known as “allelic dropout” the effect is concealment of an allele, and therefore the potential incorrect genotyping of the individual as homozygote [61]. Preferential amplification may happen or escalate when there is differential denaturation or annealing temperature among the PCR products [111]. Similar complications might be seen when PCR inhibitors are present [112]. Utilization of Mini-STRs [113,114] reduces the probability of allele drop out [115].

4.3. PCR miscoding lesions

PCR miscoding lesions describe the misincorporation of erroneous bases during the extension phase of PCR which may result in misidentification of the samples. Two mechanisms are considered as the underlying cause of PCR miscoding lesions including the intrinsic errors of amplification enzyme error and the errors generated by post-mortem biochemical modifications of the original starting template molecules. Taq polymerase has an error rate of 2×10^{-4} [116] resulting in accumulation of 0.8 mutant positions on a 100 bp double strand sequence after 20 amplification cycles [117]. However the error rate of the Taq polymerase enzyme may vary more than 10 folds depending on the DNA sequence and amplification conditions [118].

The hydrolytic deamination of DNA strands mostly affects cytosine, 5-methylcytosine, adenine and guanine, resulting in formation of uracil, thymine, hypoxanthine and xanthine, respectively. This phenomenon leads to misincorporation of incorrect bases during PCR extension (A instead of G, and C instead of T) [53,61]. Strand specific studies have revealed that the majority of damage-induced miscoding lesions were (C to T) transitions resulting in (G to A) PCR misplacement [119]. However size-based STR analyses may still be satisfactory [120].

The polymerase enzyme based errors can be minimized by using high-fidelity polymerase enzymes with proofreading activity, such as pfu and Taq HiFi [121–123]. Nucleotide misincorporations during the early cycles of the PCR may propagate through the later cycles and cause incorrect sequences. Duplicate analysis would reduce the potential error rate associated with automated sequencings.

Pre-amplification procedures such as degenerated oligonucleotide PCR [124] and primer extension PCR [125,126] strategies have been introduced to enhance the retrieval of degraded DNA molecules. However, these strategies do not eliminate the risk of miscoding lesions.

4.4. Jumping PCR phenomenon

The ‘Jumping PCR’ phenomenon may take place in degraded DNA extracts if no amplifiable template DNA molecules spanning the complete fragment defined by the primers exist. The amplification may proceed through single primer amplifications of shorter available fragments. By annealing to another complementary segment those amplified fragments may serve as primers for further extensions of the products. The single primer extension

continues from both sides until a double stranded product with minimum desired length was produced after which a conventional amplification reaction would take place. Jumping PCR allows the amplification of regions longer than the longest amplifiable molecule in the extract [61]. Compatible results through repeated amplifications of the same template could verify the results in heterozygotic or modern DNA contamination cases [127,128].

4.5. PCR inhibition versus DNA structural modifications

Beside the potential errors in sampling, DNA purification, DNA quantitation, PCR and automated sequencings, a major concern in analysis of forensic specimens arises from the coexistence of PCR inhibitors in the extracted material. Through a variety of mechanisms, endo- or exogenous inhibitors would hamper the progression of PCR amplifications [129]. The inhibited and low copy number profiles may look similar at some points, however a detailed discussion on the composition or the effects of those materials on forensic genetic analysis is beyond the context of the current paper.

5. DNA repair strategies

The ability to repair DNA damages through the current technology is limited. Attempting to repair some of the nicks generated by DNA degradation in ancient DNA studies Pusch et al. introduced a pre-PCR repair method [130]. In that method *Escherichia coli* DNA polymerase I translates the nicks in the duplex DNA, and the remaining gaps are closed by the subsequent use of T₄ DNA ligase [130]. That sort of concerted action of the two enzymes repair can only be successful for molecules with unmodified 3' OH and/or 5' P termini which can be terminally elongated by DNA polymerase I and then sealed by T₄ DNA ligase [107].

In multiple displacement amplification (MDA) [131] whole genome amplification could be achieved from small quantities of genomic DNA by isothermal amplification with bacteriophage ϕ 29 DNA polymerase- and exonuclease-resistant random hexamer primers. However the modes of post-mortem damage products are not yet fully defined to allow the design of effective *in vitro* repair strategies [107].

In ancient DNA studies coprolites, were shown to contain large amounts of crosslinks between reducing sugars and other components hence producing Maillard products. Such crosslinks from coprolite extracts can be removed by N-phenacyl thiazolium bromide (PTB) [132]. In some cases DNA amplification from the coprolite was possible only after sugar-derived crosslinks had been resolved by PTB [133]. However the effectiveness of PTB has been challenged at least in some cases where different components were responsible for PCR inhibition and most notably removable by silica extractions [134]. The Maillard reaction products may be much more stable and especially more resistant to hydrolysis as a natural preservation mechanism [135].

Considering the overview proposed by Paabo et al. [53], Table 2 summarises the different types of DNA damage and their effects on PCR profiling.

6. DNA preservation

A dry environment would greatly reduce the rate of degradation. Presence of water as moisture encourages the growth of bacteria and also provides substrate for hydrolytic enzymes [10]. This may explain the higher stability of DNA in skeletal tissue and also employing desiccation techniques for DNA preservation. Constant low temperatures play a central role in the longevity of DNA molecules [20,136]. A 20 °C decrease in temperature results in 10–25-fold reduction in the rate of nucleotide bases decomposition [68].

Table 2

Overview of different types of DNA damage and their impact on forensic profiling.

Type of damage	Underlying cause	Effects on DNA template	Effects on profiling	Trouble shooting
Strand breaks	Nuclease activities, hydrolytic end products	Reduction of overall template copy numbers, size reduction	PCR failure, increased stutter peaks, inconsistent PCR outcomes	Consider resampling, multiple independent PCRs, size reduction, Mini-STRs, mitochondrial DNA, SNPs, increase PCR cycles up to 34, extra doses of Taq polymerase, DNA repair
Oxidative lesions	Bacterial metabolism, radiation	Base or sugar fragmentation	PCR failure	Same as above
Hydrolytic deamination	Hydrolytic loss of amino groups	Change of coding potential	PCR miscoding lesions	Same as above
DNA crosslinkage	Hydrolytic base loss, reduced sugar residues	DNA to DNA or DNA to other biomolecules linkage	PCR failure	Same as above, repair of crosslinkages

However based on the independent DNA retrieval studies DNA cannot survive for more than 1 million year in the most frozen conditions [137]. Enzyme-catalysed reactions are less sensitive to temperature changes compared to spontaneous reactions [138].

Environmental conditions have more influence on DNA degradation compared to time elapsed since deposition of the tissue. Neutral or slightly alkaline pH in the sample or in the soil favors DNA preservation [20]. Due to the rapid decomposition rate, tissues such as blood and kidney were found to be unsuitable for DNA fingerprinting after a period of 1 week [139]. Brain, lymph nodes and skeletal muscles preserved high molecular DNA up to 3 weeks, whereas kidney, thyroid and spleen retained it for 1 week and liver lost all high molecular DNA after 2 days. Brain tissue seems to be one of the best sources of DNA for post-mortem studies followed by muscle and blood and then other internal organs, whereas liver is consistently a poor source of DNA [140]. The morphological preservation of tissue although helpful in determination of DNA preservation is not always an accurate predictive measure [141]. This is typical of most naturally and artificially mummified remains in Egypt [142]. Binding of the DNA to mineral surfaces as it happens in bone samples may also lead to enhanced preservation [143]. Fermentation activities of anaerobic bacteria in the intestines or in the soil and presence of humic acids make further contribution to pH fall and therefore DNA degradation [10].

7. Quantification of DNA fragments

A number of DNA strand breakage assays are based on the general principle that under *in vitro* denaturation conditions of high pH, the rate at which single-stranded DNA is released from the duplex DNA is proportional to the number of strand breaks in the DNA molecule [144]. Those methods including DNA alkaline unwinding assay [145], DNA alkaline elution assay [146] and Fast Micromethod [147,148] however gained limited attention in forensic context due to the coexistence of microorganismal DNA in the analyte. Utilization of human specific DNA hybridization or PCR techniques would appropriately be considered in those cases. Single cell gel electrophoresis (also known as Comet assay) has been widely accepted as a simple, sensitive, and rapid tool for assessing DNA damage and repair [149]. Moreover various laboratories have attempted to use quantitative PCR (QPCR) to detect DNA damage in specific gene segments. If done properly the method has the potential to overcome the problematic shortage of template encountered in highly damaged DNA cases [26,150] and the detection of PCR inhibition phenomenon in amplification of forensic samples [151].

8. Practical applications of DNA degradation

The significant role of cations in DNA nuclease activities has been considered in the production of DNA preservation solid

media. FTA[®] cards which remove cations and maintain neutral pH and dryness have been successfully applied in storage and transportation of DNA containing material [152].

The level of DNA degradation could be assessed by PCR amplification and detection of two DNA target sequences of different lengths [151,153]. This would assist in selecting the best approach (e.g., STRs, Mini-STRs or SNPs analysis of genomic or mtDNA) for DNA profiling as well as interpretation and validation of the results.

Comet assay gave useful quantitative measures in early post-mortem period. Statistical correlation was found between DNA fragmentation and post-mortem interval from 0 to 56 h post-mortem in tissues such as leukocytes and muscle cells indicating that nuclear DNA is fragmented following death in an organ and time dependent manner. The first 24 h post-mortem represent the fastest degradation rate [154]. A close relationship was found between DNA degradation rate and post-mortem interval (PMI) for the first 72 h by comet assay of rat myocardial cells [155], indicating the usefulness of that application in estimation of time since death in early postmortal periods.

9. Conclusion

As a highly reactive chemical residue, DNA is the target of several physical agents and chemical reactions. Numerous modifications of DNA fragments or various substances accumulating in the tissue during decay process and co-extracted with DNA may affect DNA amplifications and therefore the outcome of DNA-based identification technologies.

The present study aimed at reviewing the causes and pathways associated with DNA degradation and their impact on DNA profiling. This informational resource can direct ideas on how to modify techniques to deal with DNA damage, and can alert the practitioner to vagaries that may be encountered in casework and possible ways to overcome them, or to take them into account in data interpretation.

Furthermore a few DNA repair technologies designed for restoration of DNA structure have been explained. However the ability to repair DNA damages through the current technology is limited and requires further investigations.

Acknowledgement

Authors wish to thank Professor Leigh Burgoyne for providing useful information about endonucleases.

References

- [1] T. Lindahl, The croonian lecture, 1996: endogenous damage to DNA, *Philos. Trans. R. Soc. Lond. B: Biol. Sci.* 351 (1347) (1996) 1529–1538.
- [2] A.A. Vass, *Beyond the grave understanding human decomposition*, 2001.

- [3] R.B. Jennings, C.E. Ganote, K.A. Reimer, Ischemic tissue injury, *Am. J. Pathol.* 81 (1) (1975) 179–198.
- [4] G. Ilse, K. Kovacs, N. Ryan, E. Horvath, D. Ilse, Autolytic changes in the rat adenohipophys. A histologic, immunocytologic and electron microscopic study, *Exp. Pathol. (Jena)* 17 (4) (1979) 185–195.
- [5] C.J. Zeiss, The apoptosis–necrosis continuum: insights from genetically altered mice, *Vet. Pathol.* 40 (5) (2003) 481–495.
- [6] S. Shimizu, Y. Eguchi, H. Kosaka, W. Kamiike, H. Matsuda, Y. Tsujimoto, Prevention of hypoxia-induced cell death by bcl-2 and bcl-xl, *Nature* 374 (6525) (1995) 811–813.
- [7] Y. Collan, M. Salmenpera, Electron microscopy of postmortem autolysis of rat muscle tissue, *Acta Neuropathol.* 35 (3) (1976) 219–233.
- [8] C. Dive, C.D. Gregory, D.J. Phipps, D.L. Evans, A.E. Milner, A.H. Wyllie, Analysis and discrimination of necrosis and apoptosis (programmed cell-death) by multiparameter flow-cytometry, *Biochim. Biophys. Acta* 1133 (3) (1992) 275–285.
- [9] D.V. Krysko, T. Vanden Berghe, K. D'Herde, P. Vandenabeele, Apoptosis and necrosis: detection, discrimination and phagocytosis, *Methods* 44 (3) (2008) 205–221.
- [10] H. Gill-King, Chemical and ultrastructural aspects of decomposition, in: M.H. Haglund, W.D. Sorg (Eds.), *Forensic Taphonomy: The Postmortem Fate of Human Remains*, Robert B. Stern, CRC press, Inc., 1997, pp. 93–108.
- [11] Y. Tomita, M. Nihira, Y. Ohno, S. Sato, Ultrastructural changes during in situ early postmortem autolysis in kidney, pancreas, liver, heart and skeletal muscle of rats, *Leg. Med. (Tokyo)* 6 (1) (2004) 25–31.
- [12] P. Nicotera, M. Leist, Energy supply and the shape of death in neurons and lymphoid cells, *Cell Death Differ.* 4 (6) (1997) 435–442.
- [13] Y. Tsujimoto, Apoptosis and necrosis: intracellular atp level as a determinant for cell death modes, *Cell Death Differ.* 4 (6) (1997) 429–434.
- [14] B.F. Trump, I.K. Berezsky, Calcium-mediated cell injury and cell death, *FASEB J.* 9 (2) (1995) 219–228.
- [15] K.A. Reimer, J.E. Lowe, R.B. Jennings, Effect of the calcium antagonist verapamil on necrosis following temporary coronary artery occlusion in dogs, *Circulation* 55 (4) (1977) 581–587.
- [16] G. Hajnoczky, E. Davies, M. Madesh, Calcium signaling and apoptosis, *Biochem. Biophys. Res. Commun.* 304 (3) (2003) 445–454.
- [17] G. Kroemer, L. Galluzzi, C. Brenner, Mitochondrial membrane permeabilization in cell death, *Physiol. Rev.* 87 (1) (2007) 99–163.
- [18] E. Jambirina, R. Alonso, M. Alcalde, M. del Carmen Rodriguez, A. Serrano, A.C. Martinez, J. Garcia-Sancho, M. Izquierdo, Calcium influx through receptor-operated channel induces mitochondria-triggered paraptotic cell death, *J. Biol. Chem.* 278 (16) (2003) 14134–14145.
- [19] C. De Duve, R. Wattiaux, Functions of lysosomes, *Annu. Rev. Physiol.* 28 (1966) 435–492.
- [20] T. Lindahl, Instability and decay of the primary structure of DNA, *Nature* 362 (6422) (1993) 709–715.
- [21] M. Hofreiter, V. Jaenicke, D. Serre, A. Haeseler, S. Paabo, DNA sequences from multiple amplifications reveal artifacts induced by cytosine deamination in ancient DNA, *Nucleic Acids Res.* 29 (23) (2001) 4793–4799.
- [22] K. Yuyama, H. Yamamoto, I. Nishizaki, T. Kato, I. Sora, T. Ya-mamoto, Caspase-independent cell death by low concentrations of nitric oxide in pc12 cells: involvement of cytochrome c oxidase inhibition and the production of reactive oxygen species in mitochondria, *J. Neurosci. Res.* 73 (3) (2003) 351–363.
- [23] M. Donovan, T.G. Cotter, Caspase-independent photoreceptor apoptosis in vivo and differential expression of apoptotic protease activating factor-1 and caspase-3 during retinal development, *Cell Death Differ.* 9 (11) (2002) 1220–1231.
- [24] M.F. Counis, A. Torriglia, Acid DNases and their interest among apoptotic endonucleases, *Biochimie* 88 (12) (2006) 1851–1858.
- [25] K.R. Chien, J. Abrams, A. Serroni, J.T. Martin, J.L. Far-ber, Accelerated phospholipid degradation and associated membrane dysfunction in irreversible, ischemic liver cell injury, *J. Biol. Chem.* 253 (13) (1978) 4809–4817.
- [26] S. Ayala-Torres, Y. Chen, T. Svoboda, J. Rosenblatt, B. Van, Houten, Analysis of gene-specific DNA damage and repair using quantitative polymerase chain reaction, *Methods* 22 (2) (2000) 135–147.
- [27] M. Hofreiter, D. Serre, H.N. Poinar, M. Kuch, S. Paabo, Ancient DNA, *Nat. Rev. Genet.* 2 (5) (2001) 353–359.
- [28] J. Filipinski, J. Leblanc, T. Youdale, M. Sikorska, P.R. Walker, Periodicity of DNA folding in higher order chromatin structures, *EMBO J.* 9 (4) (1990) 1319–1327.
- [29] A.M. Martelli, R. Bareggi, R. Bortul, V. Grill, P. Narducci, M. Zwyer, The nuclear matrix and apoptosis, *Histochem. Cell. Biol.* 108 (1) (1997) 1–10.
- [30] D.G. Brown, X.M. Sun, G.M. Cohen, Dexamethasone-induced apoptosis involves cleavage of DNA to large fragments prior to inter-nucleosomal fragmentation, *J. Biol. Chem.* 268 (5) (1993) 3037–3039.
- [31] N. Anzai, H. Kawabata, T. Hirama, H. Masutani, Y. Ueda, Y. Yoshida, M. Okuma, Types of nuclear endonuclease activity capable of inducing internucleosomal DNA fragmentation are completely different between human cd34+ cells and their granulocytic descendants, *Blood* 86 (3) (1995) 917–923.
- [32] H.S. Huang, A. Matevosian, Y. Jiang, S. Akbarian, Chromatin immunoprecipitation in postmortem brain, *J. Neurosci. Methods* 156 (1–2) (2006) 284–292.
- [33] P. Huang, K. Ballal, W. Plunkett, Biochemical characterization of the protein activity responsible for high molecular weight DNA fragmentation during drug-induced apoptosis, *Cancer Res.* 57 (16) (1997) 3407–3414.
- [34] P.R. Walker, S. Pandey, M. Sikorska, Degradation of chromatin in apoptotic cells, *Cell Death Differ.* 2 (2) (1995) 97–104.
- [35] X.M. Sun, G.M. Cohen, Mg(2+)-dependent cleavage of DNA into kilobase pair fragments is responsible for the initial degradation of DNA in apoptosis, *J. Biol. Chem.* 269 (21) (1994) 14857–14860.
- [36] M. Nakamura, Y. Sakaki, N. Watanabe, Y. Takagi, Purification and characterization of the Ca²⁺ plus Mg²⁺-dependent endodeoxyri-bonuclease from calf thymus chromatin, *J. Biochem.* 89 (1) (1981) 143–152.
- [37] M.A. Barry, A. Eastman, Identification of deoxyribonuclease ii as an endonuclease involved in apoptosis, *Arch. Biochem. Biophys.* 300 (1) (1993) 440–450.
- [38] G. Simpson, D. Roomes, B. Reeves, Successful treatment of empyema thoracis with human recombinant deoxyribonuclease, *Thorax* 58 (4) (2003) 365–366.
- [39] C. Giannakis, I.J. Forbes, P.D. Zalewski, Ca²⁺/Mg²⁺-dependent nuclease: tissue distribution, relationship to inter-nucleosomal DNA fragmentation and inhibition by Zn²⁺, *Biochem. Biophys. Res. Commun.* 181 (2) (1991) 915–920.
- [40] Y. Gavrieli, Y. Sherman, S.A. Ben-Sasson, Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation, *J. Cell. Biol.* 119 (3) (1992) 493–501.
- [41] D. Shiokawa, S. Tanuma, Characterization of human DNase i family endonucleases and activation of DNase gamma during apoptosis, *Biochemistry* 40 (1) (2001) 143–152.
- [42] S. Sunaga, A. Yoshimori, D. Shiokawa, S. Tanuma, Structure basis for the inhibitory mechanism of a novel DNase gamma-specific inhibitor, dr396, *Bioorg. Med. Chem.* 14 (12) (2006) 4217–4226.
- [43] A.G. Baranovskii, V.N. Buneva, G.A. Nevinsky, Human deoxyribonucleases, *Biochem. (Mosc)* 69 (6) (2004) 587–601.
- [44] N. Mukae, M. Enari, H. Sakahira, Y. Fukuda, J. Inazawa, H. Toh, S. Nagata, Molecular cloning and characterization of human caspase-activated DNase, *Proc. Natl. Acad. Sci. U.S.A.* 95 (16) (1998) 9123–9128.
- [45] J. Lieberman, Z. Fan, Nuclear war: the granzyme a-bomb, *Curr. Opin. Immunol.* 15 (5) (2003) 553–559.
- [46] P. Widlak, L.Y. Li, X. Wang, W.T. Garrard, Action of recombinant human apoptotic endonuclease g on naked DNA and chromatin substrates: cooperation with exonuclease and DNase i, *J. Biol. Chem.* 276 (51) (2001) 48404–48409.
- [47] I.A. Cymerman, I. Chung, B.M. Beckmann, J.M. Bujnicki, G. Meiss, Exog, a novel paralog of endonuclease g in higher eukaryotes, *Nucleic Acids Res.* 36 (4) (2008) 1369–1379.
- [48] J. Antheunisse, DNA decomposition by soil microorganisms, *Antonie Van Leeuwenhoek* 37 (2) (1971) 258–259.
- [49] J. Antheunisse, Decomposition of nucleic acids and some of their degradation products by microorganisms, *Antonie Van Leeuwenhoek* 38 (3) (1972) 311–327.
- [50] D.B. Nedwell, The input and mineralization of organic carbon in anaerobic sediments, in: K.C. Marshal (Ed.), *In Advances in Microbial Ecology*, vol. 7, Plenum Press, New York and London, 1984, pp. 93–131.
- [51] G. Eglinton, G.A. Logan, Molecular preservation, *Philos. Trans. R. Soc. Lond. B: Biol. Sci.* 333 (1268) (1991) 315–327.
- [52] A. Gotherstrom, M.J. Collins, A. Angerbjörn, K. Lidn, Bone preservation and DNA amplification, *Archaeometry* 44 (3) (2002) 395–404.
- [53] S. Paabo, H. Poinar, D. Serre, V. Jaenicke-Despres, J. Hebler, N. Roh-land, M. Kuch, J. Krause, L. Vigilant, M. Hofreiter, Genetic analyses from ancient DNA, *Annu. Rev. Genet.* 38 (2004) 645–679.
- [54] E.H. Serspersu, D. Shortle, A.S. Mildvan, Kinetic and magnetic resonance studies of active-site mutants of staphylococcal nuclease: factors contributing to catalysis, *Biochemistry* 26 (5) (1987) 1289–1300.
- [55] P. Crine, W.G. Verly, A study of DNA spontaneous degradation, *Biochim. Biophys. Acta* 442 (1) (1976) 50–57.
- [56] C. Tamm, H.S. Shapiro, R. Lipshitz, E. Chargaff, Distribution density of nucleotides within a desoxyribonucleic acid chain, *J. Biol. Chem.* 203 (2) (1953) 673–688.
- [57] B.P. Donald, M. Kaufmann, The degradation by ribonuclease of substrates other than ribonucleic acid, *J. Histochem. Cytochem.* 2 (5) (1954) 387–394.
- [58] T. Lindahl, A. Andersson, Rate of chain breakage at apurinic sites in double-stranded deoxyribonucleic acid, *Biochemistry* 11 (19) (1972) 3618–3623.
- [59] T. Lindahl, B. Nyberg, Rate of depurination of native deoxyribonucleic acid, *Biochemistry* 11 (19) (1972) 3610–3618.
- [60] E.M. Geigl, On the circumstances surrounding the preservation and analysis of very old DNA, *Archaeometry* 44 (2002) 337–342.
- [61] S. Paabo, Ancient DNA: extraction, characterization, molecular cloning, and enzymatic amplification, *Proc. Natl. Acad. Sci. U.S.A.* 86 (6) (1989) 1939–1943.
- [62] A. Sancar, L.A. Lindsey-Boltz, K. Unsal-Kacmaz, S. Linn, Molecular mechanisms of mammalian DNA repair and the DNA damage checkpoints, *Annu. Rev. Biochem.* 73 (2004) 39–85.
- [63] E. Freese, M. Cashel, Crosslinking of deoxyribonucleic acid by exposure to low pH, *Biochim. Biophys. Acta* 91 (1964) 67–77.
- [64] C. Goffin, W.G. Verly, Interstrand DNA crosslinks due to ap (apurinic/apyrimidinic) sites, *FEBS Lett.* 161 (1) (1983) 140–144.
- [65] U. Dutta, M.A. Cohenford, J.A. Dain, Nonyzymatic glycation of DNA nucleosides with reducing sugars, *Anal. Biochem.* 345 (2) (2005) 171–180.
- [66] W. Seidel, M. Pischetsrieder, DNA-glycation leads to depurination by the loss of n2-carboxyethylguanine in vitro, *Cell. Mol. Biol. (Noisy-le-grand)* 44 (7) (1998) 1165–1170.
- [67] R. Teoule, J. Cadet, in: A.J. Bertinchamps, J. Hutterman, W. Kohnlein, R. Teoule (Eds.), *Effects of Ionizing Radiation on DNA*, Springer, Berlin, 1978, pp. 171–203.
- [68] M. Hoss, P. Jaruga, T.H. Zastawny, M. Dizdargolu, S. Paabo, DNA damage and DNA sequence retrieval from ancient tissues, *Nucleic Acids Res.* 24 (7) (1996) 1304–1307.
- [69] A. Hall, J. Ballantyne, Characterization of uvc-induced DNA damage in blood-stains: forensic implications, *Anal. Bioanal. Chem.* 380 (1) (2004) 72–83.
- [70] J.M. Butler, *Forensic DNA Typing: Biology, Technology, and Genetics of STR Markers*, 2nd ed., Elsevier Academic Press, New York, 2005.
- [71] B. Sobrino, M. Brion, A. Carracedo, SNPs in forensic genetics: a review on SNP typing methodologies, *Forensic Sci. Int.* 154 (2–3) (2005) 181–194.

- [72] P. Fattorini, R. Ciofali, F. Cossutta, P. Giulianini, P. Edomi, M. Furlanut, C. Previdere, Fidelity of polymerase chain reaction-direct sequencing analysis of damaged forensic samples, *Electrophoresis* 20 (17) (1999) 3349–3357.
- [73] R. Sparkes, C. Kimpton, S. Gilbard, P. Carne, J. Andersen, N. Ol-droyd, D. Thomas, A. Urquhart, P. Gill, The validation of a 7-locus multiplex STR test for use in forensic casework. (ii). Artefacts, casework studies and success rates, *Int. J. Legal Med.* 109 (4) (1996) 195–204.
- [74] A. Alonso, P. Martin, C. Albarra, P. Garcia, D. Primorac, O. Garcia, L. Fernandez de Simon, J. Garcia-Hirschfeld, M. Sancho, J. Fernandez-Piqueras, Specific quantification of human genomes from low copy number DNA samples in forensic and ancient DNA studies, *Croat. Med. J.* 44 (3) (2003) 273–280.
- [75] B.E. Krenke, A. Tereba, S.J. Anderson, E. Buel, S. Culhane, C.J. Finis, C.S. Tomsey, J.M. Zachetti, A. Masibay, D.R. Rabbach, E.A. Amiot, C.J. Sprecher, Validation of a 16-locus fluorescent multiplex system, *J. Forensic Sci.* 47 (4) (2002) 773–785.
- [76] J.A. Irwin, M.D. Leney, O. Loreille, S.M. Barritt, A.F. Christensen, T.D. Holland, B.C. Smith, T.J. Parsons, Application of low copy number STR typing to the identification of aged, degraded skeletal remains, *J. Forensic Sci.* 52 (6) (2007) 1322–1327.
- [77] M. Takahashi, Y. Kato, H. Mukoyama, H. Kanaya, S. Kamiyama, Evaluation of five polymorphic microsatellite markers for typing DNA from decomposed human tissues-correlation between the size of the alleles and that of the template DNA, *Forensic Sci. Int.* 90 (1–2) (1997) 1–9.
- [78] D.T. Chung, J. Drabek, K.L. Opel, J.M. Butler, B.R. McCord, A study on the effects of degradation and template concentration on the amplification efficiency of the STR miniplex primer sets, *J. Forensic Sci.* 49 (4) (2004) 733–740.
- [79] J.M. Butler, Y. Shen, B.R. McCord, The development of reduced size STR amplicons as tools for analysis of degraded DNA, *J. Forensic Sci.* 48 (5) (2003) 1054–1064.
- [80] J.M. Butler, B.C. Levin, Forensic applications of mitochondrial DNA, *Trends Biotechnol.* 16 (4) (1998) 158–162.
- [81] J.M. Butler, C.H. Becker, Improved analysis of DNA short tandem repeats with time-of-flight mass spectrometry, Technical Report, National Institute of Justice, 2001.
- [82] D. Schmalzing, L. Koutny, A. Adourian, P. Belgrader, P. Matsudaira, D. Ehrlich, DNA typing in thirty seconds with a microfabricated device, *Proc. Natl. Acad. Sci. U.S.A.* 94 (19) (1997) 10273–10278.
- [83] P. Gill, An assessment of the utility of single nucleotide polymorphisms (SNPs) for forensic purposes, *Int. J. Legal Med.* 114 (4–5) (2001) 204–210.
- [84] N. Onori, V. Onofri, F. Alessandrini, L. Buscemi, M. Pesaresi, C. Turchi, A. Tagliabracchi, Post-mortem DNA damage: a comparative study of STRs and SNPs typing efficiency in simulated forensic samples, in: *International Congress Series*, volume 1288, 2006, 510–512.
- [85] D. Bogenhagen, D.A. Clayton, The number of mitochondrial deoxyribonucleic acid genomes in mouse l and human hela cells. quantitative isolation of mitochondrial deoxyribonucleic acid, *J. Biol. Chem.* 249 (24) (1974) 7991–7995.
- [86] E.D. Robin, R. Wong, Mitochondrial DNA molecules and virtual number of mitochondria per cell in mammalian cells, *J. Cell. Physiol.* 136 (3) (1988) 507–513.
- [87] S. Andelinovic, D. Sutlovic, I. Erceg Ivkovic, V. Skaro, A. Ivkovic, F. Paic, B. Rezić, M. Definis-Gojanovic, D. Primorac, Twelve-year experience in identification of skeletal remains from mass graves, *Croat. Med. J.* 46 (4) (2005) 530–539.
- [88] K.M. Sullivan, R. Hoggood, P. Gill, Identification of human remains by amplification and automated sequencing of mitochondrial DNA, *Int. J. Legal Med.* 105 (2) (1992) 83–86.
- [89] M.N. Gabriel, C.D. Calloway, R.L. Reynolds, D. Primorac, Identification of human remains by immobilized sequence-specific oligonucleotide probe analysis of mtDNA hypervariable regions i and ii, *Croat. Med. J.* 44 (3) (2003) 293–298.
- [90] Y. Seo, T. Uchiyama, K. Shimizu, K. Takahama, Identification of remains by sequencing of mitochondrial DNA control region, *Am. J. Forensic Med. Pathol.* 21 (2) (2000) 138–143.
- [91] S. Lutz, H.J. Weisser, J. Heizmann, S. Pollak, mtDNA as a tool for identification of human remains. identification using mtDNA, *Int. J. Legal Med.* 109 (4) (1996) 205–209.
- [92] M.M. Holland, D.L. Fisher, L.G. Mitchell, W.C. Rodriguez, J.J. Canik, C.R. Merril, V.W. Weedn, Mitochondrial DNA sequence analysis of human skeletal remains: identification of remains from the vietnam war, *J. Forensic Sci.* 38 (3) (1993) 542–553.
- [93] H. Mornstad, H. Pfeiffer, C. Yoon, A. Teivens, Demonstration and semi-quantification of mtDNA from human dentine and its relation to age, *Int. J. Legal Med.* 112 (2) (1999) 98–100.
- [94] Y. Yamada, H. Ohira, H. Iwase, T. Takatori, M. Nagao, S. Ohtani, Sequencing mitochondrial DNA from a tooth and application to forensic odontology, *J. Forensic Odontostomatol.* 15 (1) (1997) 13–16.
- [95] C. Ginther, L. Issel-Tarver, M.C. King, Identifying individuals by sequencing mitochondrial DNA from teeth, *Nat. Genet.* 2 (2) (1992) 135–138.
- [96] R.E. Cline, N.M. Laurent, D.R. Foran, The fingernails of mary sullivan: developing reliable methods for selectively isolating endogenous and exogenous DNA from evidence, *J. Forensic Sci.* 48 (2) (2003) 328–333.
- [97] T. Melton, K. Nelson, Forensic mitochondrial DNA analysis: two years of commercial casework experience in the united states, *Croat. Med. J.* 42 (3) (2001) 298–303.
- [98] B. Arezi, W. Xing, J.A. Sorge, H.H. Hogrefe, Amplification efficiency of thermostable DNA polymerases, *Anal. Biochem.* 321 (2) (2003) 226–235.
- [99] H. Utsuno, K. Minaguchi, Influence of template DNA degradation on the genotyping of SNPs and STR polymorphisms from forensic materials by PCR, *Bull. Tokyo Dent. Coll.* 45 (1) (2004) 33–46.
- [100] H.N. Poinar, The top 10 list: criteria of authenticity for DNA from ancient and forensic samples, *Prog. Forensics Genet.* 9 (2003) 575–579.
- [101] P. Gill, Application of low copy number DNA profiling, *Croat. Med. J.* 42 (3) (2001) 229–232.
- [102] P. Taberlet, S. Griffin, B. Goossens, S. Questiau, V. Manceau, N. Es-caravage, L.P. Waits, J. Bouvet, Reliable genotyping of samples with very low DNA quantities using PCR, *Nucleic Acids Res.* 24 (16) (1996) 3189–3194.
- [103] B. Budowle, D. Hobson, J. Smerick, J. Smith, Low copy number: consideration and caution, In *Twelfth International Symposium on Human Identification*, 2001. Available on: <http://www.promega.com>.
- [104] C.M. Strom, S. Rechitsky, Use of nested PCR to identify charred human remains and minute amounts of blood, *J. Forensic Sci.* 43 (3) (1998) 696–700.
- [105] E.K. Hanson, J. Ballantyne, Whole genome amplification strategy for forensic genetic analysis using single or few cell equivalents of genomic DNA, *Anal. Biochem.* 346 (2) (2005) 246–257.
- [106] P.J. Smith, J. Ballantyne, Simplified low-copy-number DNA analysis by post-PCR purification, *J. Forensic Sci.* 52 (4) (2007) 820–829.
- [107] G. Di Bernardo, S. Del Gaudio, M. Cammarota, U. Gaiderisi, A. Cas-cino, M. Cipollaro, Enzymatic repair of selected cross-linked ho-moduplex molecules enhances nuclear gene rescue from pompeii and herculaneum remains, *Nucleic Acids Res.* 30 (4) (2002) e16.
- [108] B.A. Budowle, J. Eisenberg, A. van Daal, Validity of low copy number typing and applications to forensic science, *Croat. Med. J.* 50 (3) (2009) 207–217.
- [109] O. Handt, M. Krings, R.H. Ward, S. Paabo, The retrieval of ancient human DNA sequences, *Am. J. Hum. Genet.* 59 (2) (1996) 368–376.
- [110] C.A. Heid, J. Stevens, K.J. Livak, P.M. Williams, Real time quantitative PCR, *Genome Res.* 6 (10) (1996) 986–994.
- [111] P.S. Walsh, H.A. Erlich, R. Higuchi, Preferential PCR amplification of alleles: mechanisms and solutions, *PCR Methods Appl.* 1 (4) (1992) 241–250.
- [112] J.P. Whitaker, T.M. Clayton, A.J. Urquhart, E.S. Millican, T.J. Downes, C.P. Kimpton, P. Gill, Short tandem repeat typing of bodies from a mass disaster: high success rate and characteristic amplification patterns in highly degraded samples, *Biotechniques* 18 (4) (1995) 670–677.
- [113] P. Wiegand, M. Kleiber, Less is more-length reduction of STR amplicons using redesigned primers, *Int. J. Legal Med.* 114 (4–5) (2001) 285–287.
- [114] L.A. Dixon, A.E. Dobbins, H.K. Pulker, J.M. Butler, P.M. Val-lone, M.D. Coble, W. Parson, B. Berger, P. Grubwieser, H.S. Mo-gensen, N. Morling, K. Nielsen, J.J. Sanchez, E. Petkovski, A. Car-racedo, P. Sanchez-Diz, E. Ramos-Luis, M. Brion, J.A. Irwin, R.S. Just, O. Loreille, T.J. Parsons, D. Syndercombe-Court, H. Schmit-ter, B. Stradmann-Bellinghausen, K. Bender, P. Gill, Analysis of artificially degraded DNA using STRs and SNPs—results of a collaborative european (eDNAp) exercise, *Forensic Sci. Int.* 164 (1) (2006) 33–44.
- [115] K. Yoshida, K. Sekiguchi, K. Kasai, H. Sato, S. Seta, G.F. Sens-abaugh, Evaluation of new primers for csf1po, *Int. J. Legal Med.* 110 (1) (1997) 36–38.
- [116] R.K. Saiki, D.H. Gelfand, S. Stoffel, S.J. Scharf, R. Higuchi, G.T. Horn, K.B. Mullis, H.A. Erlich, Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase, *Science* 239 (4839) (1988) 487–491.
- [117] P. Keohavong, W.G. Thilly, Fidelity of DNA polymerases in DNA amplification, *Proc. Natl. Acad. Sci. U.S.A.* 86 (23) (1989) 9253–9257.
- [118] K.A. Eckert, T.A. Kunkel, DNA polymerase fidelity and the polymerase chain reaction, *PCR Methods Appl.* 1 (1) (1991) 17–24.
- [119] M.T. Gilbert, J. Binladen, W. Miller, C. Wiuf, E. Willerslev, H. Poinar, J.E. Carlson, J.H. Leebens-Mack, S.C. Schuster, Recharacterization of ancient DNA miscoding lesions: insights in the era of sequencing-by-synthesis, *Nucleic Acids Res.* 35 (1) (2007) 1–10.
- [120] P. Hoff-Olsen, S. Jacobsen, B. Mevag, B. Olaisen, Microsatellite stability in human post-mortem tissues, *Forensic Sci. Int.* 119 (3) (2001) 273–278.
- [121] A. Hansen, E. Willerslev, C. Wiuf, T. Mourier, P. Arctander, Statistical evidence for miscoding lesions in ancient DNA templates, *Mol. Biol. Evol.* 18 (2) (2001) 262–265.
- [122] E. Willerslev, A.J. Hansen, B. Christensen, J.P. Steffensen, P. Arctander, Diversity of holocene life forms in fossil glacier ice, *Proc. Natl. Acad. Sci. U.S.A.* 96 (14) (1999) 8017–8021.
- [123] A. Cooper, C. Lalueza-Fox, S. Anderson, A. Rambaut, J. Austin, R. Ward, Complete mitochondrial genome sequences of two extinct moas clarify ratite evolution, *Nature* 409 (6821) (2001) 704–707.
- [124] C.M. Pusch, G.J. Nicholson, L. Bachmann, M. Scholz, Degenerate oligonucleotide-primed amplification of ancient DNA allows the retrieval of authentic DNA sequences, *Anal. Biochem.* 279 (1) (2000) 118–122.
- [125] L. Zhang, X. Cui, K. Schmitt, R. Hubert, W. Navidi, N. Arnheim, Whole genome amplification from a single cell: implications for genetic analysis, *Proc. Natl. Acad. Sci. U.S.A.* 89 (13) (1992) 5847–5851.
- [126] K. Satoh, K. Takahashi, Y. Itoh, R. Kobayashi, Typing of DNA using the primer extension preamplification (pep) method-studies of reliability of typing and detection limits, *Nihon Hoigaku Zasshi* 52 (3) (1998) 184–190.
- [127] D.A. Lawlor, C.D. Dickel, W.W. Hauswirth, P. Parham, Ancient hla genes from 7,500-year-old archaeological remains, *Nature* 349 (6312) (1991) 785–788.
- [128] O. Handt, M. Richards, M. Trommsdorff, C. Kilger, J. Simanainen, O. Georgiev, K. Bauer, A. Stone, R. Hedges, W. Schaffner, Molecular genetic analyses of the tyrolean ice man, *Science* 264 (5166) (1994) 1775–1778.
- [129] I.G. Wilson, Inhibition and facilitation of nucleic acid amplification, *Appl. Environ. Microbiol.* 63 (10) (1997) 3741–3751.
- [130] C.M. Pusch, I. Giddings, M. Scholz, Repair of degraded duplex DNA from pre-historic samples using *Escherichia coli* DNA polymerase i and t4 DNA ligase, *Nucleic Acids Res.* 26 (3) (1998) 857–859.
- [131] F.B. Dean, S. Hosono, L. Fang, X. Wu, A.F. Faruqi, P. Bray-Ward, Z. Sun, Q. Zong, Y. Du, J. Du, M. Driscoll, W. Song, S.F. Kingsmore, M. Egholm, R.S. Lasken, Comprehensive human genome amplification using multiple displacement amplification, *Proc. Natl. Acad. Sci. U.S.A.* 99 (8) (2002) 5261–5266.
- [132] S. Vasan, X. Zhang, A. Kapurniotu, J. Bernhagen, S. Teichberg, J. Basgen, D. Wagle, D. Shih, I. Terlecky, R. Bucala, A. Cerami, J. Egan, P. Ulrich, An agent cleaving

- glucose-derived protein crosslinks in vitro and in vivo, *Nature* 382 (6588) (1996) 275–278.
- [133] H.N. Poinar, M. Hofreiter, W.G. Spaulding, P.S. Martin, B.A. Stankiewicz, H. Bland, R.P. Evershed, G. Possnert, S. Paabo, Molecular coproscopy: dung and diet of the extinct ground sloth *nothrotheriops shastensis*, *Science* 281 (5375) (1998) 402–406.
- [134] B.M. Kemp, C. Monroe, D.G. Smith, Repeat silica extraction: a simple technique for the removal of PCR inhibitors from DNA extracts, *J. Archaeol. Sci.* 33 (12) (2006) 1680–1689.
- [135] J.L. Bada, X.S. Wang, H. Hamilton, Preservation of key biomolecules in the fossil record: current knowledge and future challenges, *Philos. Trans. R. Soc. Lond. B: Biol. Sci.* 354 (1379) (1999) 7787.
- [136] E. Willerslev, A.J. Hansen, H.N. Poinar, Isolation of nucleic acids and cultures from fossil ice and permafrost, *Trends Ecol. Evol.* 19 (3) (2004) 141–147.
- [137] E. Willerslev, A. Cooper, Ancient DNA, *Proc. Biol. Sci.* 272 (1558) (2005) 3–16.
- [138] M.A. Tabatabai, Soil enzymes, in: R.H. Miller, D.R. Keeney (Eds.), *Methods of Soil Analysis*, 2nd ed., American Society of Agronomy Soil Science Society of America, Madison, WI, 1982, pp. 903–947.
- [139] B. Ludes, H. Pfitzinger, P. Mangin, DNA fingerprinting from tissues after variable postmortem periods, *J. Forensic Sci.* 38 (3) (1993) 686–690.
- [140] T.J. Parsons, V.W. Weedn, Chemical and ultrastructural aspects of decomposition, in: W.D. Haglund, M.H. Sorg (Eds.), *Forensic Taphonomy: The Post-mortem Fate of Human Remains*, Robert B. Stern, CRC press, Inc., 1997, pp. 93–108.
- [141] P.F. van Bergen, H.A. Bland, M.C. Horton, R.P. Evershed, Chemical and morphological changes in archaeological seeds and fruits during preservation by desiccation, *Geochim. Cosmochim. Acta* 61 (1997) 1919–1930.
- [142] H.N. Poinar, B.A. Stankiewicz, Protein preservation and DNA retrieval from ancient tissues, *Proc. Natl. Acad. Sci. U.S.A.* 96 (15) (1999) 8426–8431.
- [143] G. Romanowski, M.G. Lorenz, W. Wackernagel, Adsorption of plasmid DNA to mineral surfaces and protection against DNase I, *Appl. Environ. Microbiol.* 57 (4) (1991) 1057–1061.
- [144] B. Rydberg, The rate of strand separation in alkali of DNA of irradiated mammalian cells, *Radiat. Res.* 61 (2) (1975) 274–287.
- [145] P.M. Kanter, H.S. Schwartz, A fluorescence enhancement assay for cellular DNA damage, *Mol. Pharmacol.* 22 (1) (1982) 145–151.
- [146] K.W. Kohn, L.C. Erickson, R.A. Ewig, C.A. Friedman, Fractionation of DNA from mammalian cells by alkaline elution, *Biochemistry* 15 (21) (1976) 4629–4637.
- [147] R. Batel, Z. Jakšić, N. Bihari, B. Hamer, M. Fafandel, C. Chauvin, H.C. Schroder, W.E. Muller, R.K. Zahn, A microplate assay for DNA damage determination (fast micromethod), *Anal. Biochem.* 270 (2) (1999) 195–200.
- [148] Z. Jakšić, R. Batel, DNA integrity determination in marine invertebrates by fast micromethod, *Aquat. Toxicol.* 65 (4) (2003) 361–376.
- [149] A. Dhawan, M. Bajpayee, D. Parmar, Comet assay: a reliable tool for the assessment of DNA damage in different models, *Cell Biol. Toxicol.* 25 (1) (2009) 5–32.
- [150] D.P. Kalinowski, S. Ilénye, B. Van Houten, Analysis of DNA damage and repair in murine leukemia I1210 cells using a quantitative polymerase chain reaction assay, *Nucleic Acids Res.* 20 (13) (1992) 3485–3494.
- [151] K.L. Swango, M.D. Timken, M.D. Chong, M.R. Buoncristiani, A quantitative PCR assay for the assessment of DNA degradation in forensic samples, *Forensic Sci. Int.* 158 (1) (2006) 14–26.
- [152] L.J. Dobbs, M.N. Madigan, A.B. Carter, L. Earls, Use of fta gene guard filter paper for the storage and transportation of tumor cells for molecular testing, *Arch. Pathol. Lab. Med.* 126 (1) (2002) 56–63.
- [153] A. Alonso, P. Martin, C. Albarran, P. Garcia, O. Garcia, L.F. de Simon, J. Garcia-Hirschfeld, M. Sancho, C. de La Rua, J. Fernandez-Piqueras, Real-time PCR designs to estimate nuclear and mitochondrial DNA copy number in forensic and ancient DNA studies, *Forensic Sci. Int.* 139 (2–3) (2004) 141–149.
- [154] L.A. Johnson, J.A. Ferris, Analysis of postmortem DNA degradation by single-cell gel electrophoresis, *Forensic Sci. Int.* 126 (1) (2002) 43–47.
- [155] J.L. Zhen, X.D. Zhang, Q.S. Niu, Relationship between the postmortem interval and nuclear DNA changes of heart muscular cells in mice, *Fa Yi Xue Za Zhi* 22 (3) (2006) 173–176.