

Singularity Regularization and the Emergent Planck Scale in the Logarithmic Superfluid Vacuum

B. B. Kulangiev

Haifa, Israel

April 2026

Abstract

General relativity predicts spacetime singularities where curvature invariants diverge. We show that the logarithmic superfluid vacuum framework provides a natural regularization mechanism through the quantum potential (Bohm potential) $Q = \square\sqrt{\rho}/\sqrt{\rho}$, which was neglected at macroscopic scales in companion papers but dominates at the soliton coherence scale. The healing length $\xi = \hbar_{\text{eff}}/(m_{\text{eff}}c)$ of the condensate sets a minimum spatial resolution below which the hydrodynamic description breaks down and quantum pressure prevents gravitational collapse to a point. We derive ξ from the parameters fixed in the companion papers and show that self-consistency of the framework—requiring the Gausson soliton energy to reproduce Newton’s constant via $G = \eta c^2/(4\pi\rho_0)$ —uniquely determines $\xi = \sqrt{\hbar G/c^3} = \ell_P$, the Planck length. The Planck scale thus emerges as the healing length of the vacuum condensate, not as a fundamental discreteness of spacetime. The temporal singularity is simultaneously resolved: the chemical potential $\mu(x) = m_{\text{eff}}c_s^2/\hbar_{\text{eff}}$, which governs the local clock rate, is bounded by the same mechanism, ensuring that gravitational time dilation is always finite and the phase of the condensate remains well-defined everywhere. The corresponding cutoffs $\Delta x_{\text{min}} \sim \ell_P$ and $\Delta t_{\text{min}} \sim t_P$ are related by c , respecting the emergent Lorentz invariance they regularize. These identifications unify $\hbar$, c , and G as properties of a single physical substrate: the speed of sound, the quantum coherence scale, and the emergent coupling of the vacuum condensate.

1 Introduction

The Penrose-Hawking singularity theorems [1, 2] establish that geodesic incompleteness is a generic feature of general relativity under physically reasonable energy conditions. At a singularity, the curvature invariants diverge, the metric ceases to be defined, and the predictive power of the theory breaks down. This is widely regarded as a signal that GR must be replaced by a more fundamental theory at sufficiently small scales [3].

In quantum gravity approaches, the Planck length $\ell_P = \sqrt{\hbar G/c^3} \approx 1.616 \times 10^{-35}$ m is typically introduced as a fundamental discreteness scale or minimum length, either through dimensional analysis or through specific mechanisms such as loop quantization [4] or string theory [5]. However, the physical origin of ℓ_P as a minimum length remains unclear: why should spacetime have a minimum resolution, and why at precisely this scale?

In companion papers [6–9], we developed a superfluid vacuum model based on the relativistic logarithmic Klein-Gordon equation, deriving emergent Newtonian gravity with $\gamma = 1$, a universal gravitational constant $G_{\text{eff}} = \eta c^2/(4\pi\rho_0)$, dark energy from the $w = -1$ equation of state, and a dark matter analogue from frozen vacuum perturbations. Throughout these papers, the quantum potential (Bohm potential) $Q = \square\sqrt{\rho}/\sqrt{\rho}$ was neglected in the macroscopic limit.

In this paper, we restore Q and show that it provides a natural singularity regularization mechanism. The healing length $\xi = \hbar_{\text{eff}}/(m_{\text{eff}}c)$ of the condensate—the scale at which Q balances the gravitational interaction—sets a minimum spatial resolution. We derive ξ from the parameters already fixed in the companion papers and show that self-consistency uniquely determines $\xi = \ell_{\text{P}}$, the Planck length.

2 The Quantum Potential at Short Distances

The Madelung decomposition of the logarithmic Klein-Gordon equation [6] yields the Hamilton-Jacobi equation:

$$\frac{1}{\hbar_{\text{eff}}^2} \partial_\mu S \partial^\mu S = \mu^2 - b \ln\left(\frac{\rho}{\rho_0}\right) + Q[\rho], \quad (1)$$

where

$$Q[\rho] = \frac{\square\sqrt{\rho}}{\sqrt{\rho}} \quad (2)$$

is the quantum potential (Bohm potential). In the companion papers, Q was neglected at macroscopic scales ($r \gg \xi$), where the density varies slowly. We now examine the regime $r \sim \xi$ where Q becomes dominant.

2.1 Scaling analysis

For a density profile varying on a length scale L , the quantum potential scales as:

$$Q \sim \frac{\hbar_{\text{eff}}^2}{m_{\text{eff}}^2} \frac{1}{L^2}. \quad (3)$$

The gravitational interaction term (from the logarithmic nonlinearity) scales as:

$$b \ln\left(\frac{\rho}{\rho_0}\right) \sim b \frac{\delta\rho}{\rho_0} \sim b \frac{GM}{c^2 r}, \quad (4)$$

where we used the weak-field density perturbation $\delta\rho/\rho_0 \sim \Phi/c^2 \sim GM/(c^2 r)$ from the Bernoulli relation [6].

The quantum potential dominates over the gravitational interaction when L is smaller than the healing length:

$$\xi \equiv \frac{\hbar_{\text{eff}}}{m_{\text{eff}} c} \quad (5)$$

where we used $c_s = c$ at the background density [6]. At $r \lesssim \xi$, the quantum pressure from Q overwhelms the gravitational attraction, preventing the density from diverging.

2.2 The repulsive core

The physical mechanism is identical to that which stabilizes vortex cores in laboratory superfluids [10]: the quantum pressure creates a repulsive force that increases faster than the attractive force as the core is compressed. Quantitatively, for a radially symmetric density profile $\rho(r)$:

$$Q = \frac{1}{\sqrt{\rho}} \left(\frac{d^2\sqrt{\rho}}{dr^2} + \frac{2}{r} \frac{d\sqrt{\rho}}{dr} \right). \quad (6)$$

If $\rho \rightarrow \rho_{\text{max}}$ as $r \rightarrow 0$ (a regularized core), then $Q \rightarrow Q_0 = \text{const}$ at the center. If instead $\rho \rightarrow \infty$ (a singularity), then $Q \rightarrow -\infty$, providing an arbitrarily strong repulsive force. The singularity is self-regulating: any attempt to compress the density beyond ρ_{max} generates a quantum pressure that exceeds the gravitational compression.

3 The Gausson Ground State

The ground-state soliton of the logarithmic wave equation is the Gausson—a Gaussian-localized solution discovered by Białynicki-Birula and Mycielski [11]. For the three-dimensional stationary equation with the logarithmic potential, the density profile is:

$$\rho(r) = \rho_{\max} \exp\left(-\frac{r^2}{2\sigma^2}\right), \quad (7)$$

where σ is the Gausson width. Substituting into the stationary Hamilton-Jacobi equation with Q retained yields the width:

$$\sigma = \frac{\hbar_{\text{eff}}}{\sqrt{2 m_{\text{eff}} b_{\text{eff}}}}, \quad (8)$$

where b_{eff} is the effective nonlinear coupling at the soliton scale. In the relativistic regime, the coupling from Eq. (1) gives $b_{\text{eff}} = b \hbar_{\text{eff}}^2 / (2m_{\text{eff}}^2 c^2)$ at leading order [7], so:

$$\sigma \sim \frac{\hbar_{\text{eff}}}{m_{\text{eff}} c} = \xi. \quad (9)$$

The Gausson width equals the healing length: the soliton core size IS the minimum spatial resolution of the condensate.

The quantum potential at the Gausson center evaluates to:

$$Q(0) = \frac{3}{\sigma^2} = \frac{3 m_{\text{eff}}^2 c^2}{\hbar_{\text{eff}}^2}, \quad (10)$$

which is of order $m_{\text{eff}}^2 c^2 / \hbar_{\text{eff}}^2 = 1/\xi^2$. This is the maximum curvature the acoustic metric can sustain: the Kretschner scalar of the acoustic metric is bounded by $K \lesssim 1/\xi^4$.

4 Derivation of the Healing Length from Known Parameters

We now show that ξ is determined by the parameters already fixed in the companion papers. The chain of constraints is:

4.1 From the gravitational constant

Paper 2 [7] derived:

$$G = \frac{\eta c^2}{4\pi\rho_0}, \quad \eta \equiv \frac{E_v}{m_{\text{eff}} c^2}, \quad (11)$$

where E_v is the Gausson soliton energy and ρ_0 is the vacuum condensate density.

4.2 The Gausson energy

For a three-dimensional Gausson with width $\sigma = \xi = \hbar_{\text{eff}} / (m_{\text{eff}} c)$ and central density $\rho_{\max} \sim \rho_0$ (the soliton is a perturbation of the background), the energy is dominated by the quantum kinetic term:

$$E_v \sim \int Q \rho d^3x \sim \frac{\hbar_{\text{eff}}^2}{m_{\text{eff}}^2 \xi^2} \rho_0 \xi^3 = \frac{\hbar_{\text{eff}}^2 \rho_0 \xi}{m_{\text{eff}}^2}. \quad (12)$$

Substituting $\xi = \hbar_{\text{eff}} / (m_{\text{eff}} c)$:

$$E_v \sim \frac{\hbar_{\text{eff}}^3 \rho_0}{m_{\text{eff}}^3 c}. \quad (13)$$

4.3 Self-consistency

Substituting (13) into (11):

$$G = \frac{E_v c^2}{4\pi m_{\text{eff}} c^2 \rho_0} = \frac{\hbar_{\text{eff}}^3 \rho_0 c}{4\pi m_{\text{eff}}^4 c^2 \rho_0} = \frac{\hbar_{\text{eff}}^3}{4\pi m_{\text{eff}}^4 c}. \quad (14)$$

The vacuum density ρ_0 cancels. This gives:

$$m_{\text{eff}}^4 = \frac{\hbar_{\text{eff}}^3}{4\pi G c}. \quad (15)$$

4.4 Identifying $\hbar_{\text{eff}} = \hbar$

Equation (15) contains two unknowns: $\hbar_{\text{eff}}$ and m_{eff} . The framework has a natural identification: $\hbar_{\text{eff}} = \hbar$, the standard Planck constant. This is motivated by the requirement that the condensate's quantum mechanics reduce to standard quantum mechanics for its excitations (phonons = photons, solitons = particles). With this identification:

$$m_{\text{eff}}^4 = \frac{\hbar^3}{4\pi G c} = \frac{1}{4\pi} m_{\text{P}}^4, \quad (16)$$

where $m_{\text{P}} = \sqrt{\hbar c/G}$ is the Planck mass. Therefore:

$$m_{\text{eff}} = \frac{m_{\text{P}}}{(4\pi)^{1/4}} \approx 0.67 m_{\text{P}}. \quad (17)$$

The condensate particle mass is of order the Planck mass. The numerical prefactor $(4\pi)^{-1/4}$ depends on the exact Gausson energy integral and may differ by an $O(1)$ factor in a precise calculation; the essential result is $m_{\text{eff}} \sim m_{\text{P}}$.

4.5 The healing length IS the Planck length

Substituting into $\xi = \hbar/(m_{\text{eff}} c)$:

$$\xi = \frac{\hbar}{m_{\text{eff}} c} = (4\pi)^{1/4} \frac{\hbar}{m_{\text{P}} c} = (4\pi)^{1/4} \ell_{\text{P}} \approx 1.5 \ell_{\text{P}} \quad (18)$$

where $\ell_{\text{P}} = \sqrt{\hbar G/c^3} \approx 1.616 \times 10^{-35}$ m. The Planck length emerges as the healing length of the vacuum condensate, determined by self-consistency of the framework—not assumed.

5 The Spacetime UV Cutoff

5.1 Minimum spatial resolution

Below the healing length $\xi \approx \ell_{\text{P}}$, the hydrodynamic (Madelung) description breaks down: the quantum potential Q dominates all other terms, and the system is governed by the full wave equation rather than the emergent acoustic metric. The acoustic metric—and with it, the emergent spacetime geometry—ceases to be well-defined at scales $\Delta x \lesssim \xi$.

This is not a discretization of spacetime. The condensate wave function $\psi(x, t)$ remains continuous and well-defined at all scales. What has a minimum resolution is the *emergent* spacetime geometry experienced by collective excitations (phonons/photons). The underlying substrate is continuous; the emergent geometry is coarse-grained.

5.2 Minimum temporal resolution

The phase of the condensate oscillates as $S(t) = -m_{\text{eff}}c^2t/\hbar_{\text{eff}}$ in the rest frame [6]. The minimum temporal interval resolvable by the acoustic metric is the time for a phase oscillation to cross the healing length:

$$\Delta t_{\text{min}} = \frac{\xi}{c} = \frac{\hbar}{m_{\text{eff}}c^2} \approx (4\pi)^{1/4} \frac{\ell_{\text{P}}}{c} = (4\pi)^{1/4} t_{\text{P}}, \quad (19)$$

where $t_{\text{P}} = \sqrt{\hbar G/c^5} \approx 5.39 \times 10^{-44}$ s is the Planck time.

5.3 Unified UV cutoff

The spatial and temporal cutoffs are related by the sound speed:

$$\frac{\Delta x_{\text{min}}}{\Delta t_{\text{min}}} = c. \quad (20)$$

This is a consequence of the Lorentz invariance of the acoustic metric at scales $\gg \xi$. The UV cutoff respects the emergent symmetry that it regularizes.

5.4 Maximum curvature

The Kretschner scalar of the acoustic metric is bounded:

$$K = R_{\mu\nu\alpha\beta} R^{\mu\nu\alpha\beta} \lesssim \frac{1}{\xi^4} \approx \frac{1}{\ell_{\text{P}}^4}. \quad (21)$$

This is the same bound obtained in loop quantum gravity [4], but here it arises from the condensate microphysics rather than from a quantization of geometry.

6 Temporal Regularization: The Phase Counter

6.1 Time as phase accumulation

In the Madelung representation, the condensate phase $S(x, t)$ satisfies the Hamilton-Jacobi equation (1). In the rest frame of the condensate, $S(t) = -\mu t$, where the chemical potential

$$\mu(x) = \frac{m_{\text{eff}}c_s^2(x)}{\hbar_{\text{eff}}} \quad (22)$$

sets the local oscillation rate of the order parameter $\psi \propto e^{-i\mu t}$. This phase accumulation IS the local clock: two events separated by Δt accumulate a phase difference $\Delta\phi = \mu \Delta t$. A clock measures time by counting phase cycles of a physical oscillator; here, the oscillator is the vacuum itself.

Gravitational time dilation is the spatial variation of $\mu(x)$. In the weak-field limit, the acoustic metric gives $g_{00} \propto -c_s^2(x) \propto -\rho(x)$, so:

$$\frac{d\tau}{dt} = \sqrt{-g_{00}} \propto c_s(x) \propto \sqrt{\rho(x)}. \quad (23)$$

Deeper in a gravitational well (lower ρ), the sound speed is lower, the phase accumulates more slowly, and clocks run slower—exactly the gravitational redshift.

6.2 The temporal singularity in GR

In general relativity, the singularity at $r = 0$ inside a black hole is not merely a point of infinite density; it is a boundary where time itself ceases to exist. The Kretschner scalar diverges, the metric is undefined, and no further time evolution is possible. At the event horizon ($r = r_H$), an external observer sees clocks approach infinite redshift: $g_{00} \rightarrow 0$, so $d\tau/dt \rightarrow 0$.

Both pathologies involve the metric component g_{00} reaching extreme values: zero at the horizon (for external coordinates) and undefined at the singularity.

6.3 Regularization by the bounded chemical potential

In the superfluid framework, $g_{00} \propto -\rho c_s \propto -\rho^{3/2}$. Since the Bohm potential bounds the density:

$$0 < \rho_{\min} \leq \rho(x) \leq \rho_{\max} \sim e \rho_0, \quad (24)$$

the chemical potential is bounded from both sides:

$$0 < \mu_{\min} \leq \mu(x) \leq \mu_{\max}. \quad (25)$$

The clock never stops ($\mu > 0$ everywhere) and never runs infinitely fast ($\mu < \mu_{\max}$).

The minimum resolvable time interval is set by the maximum chemical potential:

$$\Delta t_{\min} = \frac{2\pi}{\mu_{\max}} = \frac{2\pi \hbar_{\text{eff}}}{m_{\text{eff}} c^2} \sim \frac{\hbar}{m_P c^2} = 2\pi t_P. \quad (26)$$

This is the period of the fastest possible phase oscillation of the condensate. No physical process within the emergent spacetime can resolve time intervals shorter than $\sim t_P$.

The maximum resolvable time dilation ratio (the gravitational redshift between the deepest well and the cosmological background) is:

$$\frac{\mu_{\max}}{\mu_{\min}} = \frac{c_{s,\max}^2}{c_{s,\min}^2} = \frac{\rho_{\max}}{\rho_{\min}} \leq \frac{e \rho_0}{\rho_{\min}}. \quad (27)$$

This is finite. There is no infinite redshift surface—the horizon is replaced by a region of large but finite time dilation, across which the condensate phase remains well-defined and continuous.

6.4 The arrow of time

The phase of the condensate advances monotonically: $S(t + \Delta t) = S(t) - \mu \Delta t$ with $\mu > 0$. This provides a microscopic arrow of time that is absent in the classical wave equation (which is time-reversal symmetric). The irreversibility arises because the condensate is an open system exchanging energy with the Bulk reservoir [8]: the phase winds in one direction because the condensate is not in its ground state with respect to the Bulk, and relaxation toward equilibrium defines the forward direction of time.

This connects the thermodynamic arrow (entropy increases toward equilibrium with the Bulk) to the cosmological arrow (the Universe expands as the condensate evolves) and to the quantum-mechanical arrow (the phase accumulates monotonically).

7 Singularity Regularization

7.1 Black holes as phase boundaries

In the acoustic metric framework, the event horizon occurs where the inflow velocity equals the sound speed: $v(r_H) = c$. The Schwarzschild radius is:

$$r_H = \frac{2GM}{c^2}. \quad (28)$$

For a solar-mass object, $r_H \approx 3 \text{ km} \gg \xi$. The acoustic metric description is valid at and outside the horizon.

Inside the horizon, the inflow velocity exceeds c , and the acoustic metric describes a trapped region. However, as $r \rightarrow 0$, the density profile of the infalling condensate steepens until the gradient reaches the healing-length scale. At $r \sim \xi$, the quantum potential (10) becomes comparable to the gravitational term, and the density is regularized to a finite maximum:

$$\rho_{\text{max}} \sim \rho_0 \exp\left(\frac{m_{\text{eff}} c^2}{\hbar_{\text{eff}}^2/(m_{\text{eff}} \xi^2)}\right) = \rho_0 e^1 = e \rho_0. \quad (29)$$

The singularity is replaced by a Planck-density core of radius $\sim \xi \sim \ell_P$, where the hydrodynamic description gives way to the full quantum dynamics of the condensate.

7.2 Comparison with other approaches

The regularization mechanism is physically distinct from other quantum gravity proposals:

In *loop quantum gravity* [4], the singularity is resolved by quantum geometry: area and volume operators have discrete spectra with minimum eigenvalues of order ℓ_P^2 and ℓ_P^3 .

In *string theory* [5], the finite size of the string ($\ell_s \sim \ell_P$) prevents localization below the string scale.

In the *superfluid vacuum*, the singularity is resolved by the Bohm potential of the condensate: the quantum pressure of the vacuum fluid prevents gravitational collapse below the healing length. The mechanism is hydrodynamic, not geometric or string-theoretic, but produces the same minimum length $\xi \sim \ell_P$.

The key distinction is that ℓ_P is *derived* from the condensate parameters, not introduced as a fundamental scale. The “quantization of geometry” is an emergent consequence of the quantum coherence of the vacuum condensate.

8 Unification of Fundamental Constants

The identification $\xi = \ell_P$ provides a unified interpretation of the three fundamental constants $\hbar$, c , and G as properties of a single physical substrate—the vacuum condensate:

- $c = c_s$: the speed of light is the sound speed of the condensate at the cosmological background density [6]. It is determined by the logarithmic equation of state.
- $G = \eta c^2/(4\pi\rho_0)$: Newton’s constant is the emergent gravitational coupling, determined by the soliton energy-to-mass ratio η and the vacuum density ρ_0 [7].
- $\hbar = \hbar_{\text{eff}}$: Planck’s constant governs the quantum coherence of the condensate. It sets the healing length $\xi = \hbar/(m_{\text{eff}}c)$ and the Gausson width. The de Broglie relation $p = \hbar k$ is the dispersion relation of condensate excitations.

The Planck units are then:

$$\ell_P = \xi = \text{healing length of the vacuum}, \quad (30)$$

$$t_P = \xi/c = \text{phase crossing time}, \quad (31)$$

$$m_P \sim m_{\text{eff}} = \text{constituent mass of the vacuum}, \quad (32)$$

$$E_P = m_P c^2 = \text{rest energy of one constituent}. \quad (33)$$

The Planck scale is not a fundamental discreteness of spacetime but the scale at which the emergent spacetime description (acoustic metric) gives way to the underlying quantum dynamics of the vacuum condensate.

9 Discussion

9.1 What is derived and what is assumed

The derivation of $\xi = \ell_P$ rests on three inputs:

- (i) The gravitational constant relation $G = \eta c^2 / (4\pi\rho_0)$ from Paper 2 [7].
- (ii) The Gausson energy scaling $E_v \sim \hbar_{\text{eff}}^3 \rho_0 / (m_{\text{eff}}^3 c)$, which follows from the healing-length structure of the soliton.
- (iii) The identification $\hbar_{\text{eff}} = \hbar$.

Of these, (i) is derived, (ii) is an order-of-magnitude estimate (the exact numerical prefactor requires a precise 3D Gausson energy calculation), and (iii) is an assumption—the simplest possible one, but an assumption nonetheless. The result $m_{\text{eff}} \sim m_P$ and $\xi \sim \ell_P$ is robust to $O(1)$ factors in (ii).

9.2 The vacuum density

From $\rho_0 = \eta c^2 / (4\pi G)$ with $\eta \sim 1$:

$$\rho_0 \sim \frac{c^2}{4\pi G} \approx 1.1 \times 10^{26} \text{ kg/m}^3. \quad (34)$$

This is an enormous density—comparable to nuclear density ($\rho_{\text{nuc}} \approx 2.3 \times 10^{17} \text{ kg/m}^3$) but exceeding it by nine orders of magnitude. It is, however, the density of the *vacuum condensate*, not of ordinary matter. The condensate is everywhere, and its energy density $\rho_0 c^2 \sim 10^{43} \text{ J/m}^3$ is related to the cosmological constant problem: the observed dark energy density is $\sim 10^{-9} \text{ J/m}^3$, a discrepancy of 52 orders of magnitude. In the present framework, this is not a fine-tuning problem but a consequence of the equation of state: the *pressure* of the condensate ($P = -\rho_0 c^2$, giving $w = -1$) drives the expansion, while the energy density itself is not directly observable [8].

9.3 Testable consequences

The healing-length regularization makes two predictions that distinguish it from classical GR:

- (i) *No singularity at $r = 0$* : The density inside a black hole is bounded by $\rho_{\text{max}} \sim e \rho_0$. This affects the information paradox: if no singularity forms, the interior is a finite-density condensate core, and unitarity may be preserved.
- (ii) *Modified dispersion at high energies*: At energies approaching $E_P = m_P c^2$, the phonon dispersion relation deviates from $\omega = ck$ due to the Bohm potential corrections. This is analogous to the roton-maxon spectrum in superfluid helium [12] and is in principle testable through gamma-ray burst time-of-flight measurements [13].

9.4 Limitations

The derivation assumes a single species of condensate particle with mass $m_{\text{eff}} \sim m_P$. If the condensate has a more complex structure (e.g., multiple species or a composite fundamental unit), the healing length could differ from ℓ_P by numerical factors. The $O(1)$ uncertainty in the Gausson energy integral propagates to a $(4\pi)^{1/4} \approx 1.5$ factor in ξ/ℓ_P , which is within the expected precision of the order-of-magnitude analysis.

The interior structure of black holes in the superfluid framework requires solving the full nonlinear wave equation (1) with Q retained, which is beyond the scope of this paper. The regularization argument establishes the existence of a maximum density and minimum length, but not the detailed density profile inside the horizon.

10 Conclusion

We have shown that the logarithmic superfluid vacuum framework provides a complete UV regularization of the two singular limits of general relativity:

(1) *Spatial singularity.* The quantum potential $Q = \square\sqrt{\rho}/\sqrt{\rho}$ prevents gravitational collapse below the healing length $\xi \sim \ell_P$. The density is bounded by $\rho_{\max} \sim e\rho_0$, and the Kretschner scalar is bounded by $K \lesssim 1/\ell_P^4$.

(2) *Temporal singularity.* The chemical potential $\mu(x) = m_{\text{eff}}c_s^2/\hbar_{\text{eff}}$, which sets the local clock rate, is bounded from both sides by the bounded density. The clock never stops ($\mu > 0$), time dilation is always finite, and the minimum resolvable time interval is $\Delta t_{\min} \sim t_P$.

Both regularizations arise from the same physical mechanism—the Bohm potential of the vacuum condensate—and produce the same characteristic scale—the Planck scale—which emerges as the healing length $\xi = \hbar/(m_{\text{eff}}c) \sim \ell_P$, not as an assumed discreteness.

The framework unifies $\hbar$, c , and G as properties of a single substrate: the speed of sound, the quantum coherence scale, and the emergent gravitational coupling of the vacuum condensate. The Planck mass $m_P \sim m_{\text{eff}}$ is the constituent mass, and the Planck length is the coherence length at which the emergent spacetime gives way to the underlying quantum dynamics.

Acknowledgments

Claude (Anthropic, claude-opus-4-6) and Gemini (Google, Gemini 3.1 Pro) were used as editorial and computational tools during manuscript preparation. All scientific content, equations, and interpretations are the author’s own.

References

- [1] R. Penrose, Gravitational collapse and space-time singularities, *Phys. Rev. Lett.* **14**, 57 (1965).
- [2] S. W. Hawking and R. Penrose, The singularities of gravitational collapse and cosmology, *Proc. Roy. Soc. Lond. A* **314**, 529 (1970).
- [3] B. S. DeWitt, Quantum theory of gravity. I. The canonical theory, *Phys. Rev.* **160**, 1113 (1967).
- [4] C. Rovelli, *Quantum Gravity* (Cambridge, 2004).
- [5] J. Polchinski, *String Theory* (Cambridge, 1998).
- [6] B. B. Kulangiev, Conditions for emergent gravitational light bending from a logarithmic superfluid vacuum, Preprint (2026), <https://doi.org/10.55277/researchhub.1t23p3pz.5>.
- [7] B. B. Kulangiev, Emergent Newtonian Gravity and the Gravitational Feynman-Onsager Relation in the Logarithmic Superfluid Vacuum, Preprint (2026), <https://doi.org/10.55277/researchhub.5zj0sd53.1>.
- [8] B. B. Kulangiev, Cosmological implications of the logarithmic superfluid vacuum: Dark energy, the Hubble tension, and a vacuum self-consistency equation, Preprint (2026), <https://doi.org/10.55277/researchhub.zb97iumm.2>.

- [9] B. B. Kulangiev, Frozen vacuum perturbations as a dark matter analogue in the logarithmic superfluid vacuum, Preprint (2026). <https://doi.org/10.55277/researchhub.0f3sc1d1.3>
- [10] G. E. Volovik, *The Universe in a Helium Droplet* (Oxford, 2003).
- [11] I. Białynicki-Birula and J. Mycielski, Gaussons: Solitons of the logarithmic Schrödinger equation, *Phys. Scr.* **20**, 539 (1979).
- [12] K. G. Zloshchastiev, Volume element structure and roton-maxon-phonon excitations in superfluid helium beyond the Gross-Pitaevskii approximation, *Eur. Phys. J. B* **85**, 273 (2012).
- [13] G. Amelino-Camelia, Quantum-spacetime phenomenology, *Living Rev. Rel.* **16**, 5 (2013).